

It will be seen from these tables that the effect of ammonia gas in solution upon the water absorbing it is to increase greatly the co-efficient of expansion and to lower very rapidly both the points of maximum density and of freezing.

In these respects the gas acts just as a salt in solution would do. Gas solution and salt solution would seem to be closely related phenomena, each resulting in the formation of a mixed liquid, viz: of a liquid composed of two sets of independently moving molecules.

The effect of ammonia gas upon the volume of the water absorbing it is expressed by the following law:

When it is absorbed by water, the increase in volume for a constant temperature is directly proportional to the amount of gas absorbed.

This may be shown to be for NH_3 gas in water by plotting a set of curves with the volumes given in the above tables as ordinates and percentages of gas as abscissae. These curves, whatever temperatures be chosen, resolve themselves into straight lines. Since for the case of CO_2 gas in water the same law had been already found true by direct measurement of the change of volume due to the absorption of the gas at constant temperatures, we are warranted in suspecting the law to be a general one.

THE ENDOCRANIUM AND MAXILLARY SUSPENSORIUM OF THE BEE.

PROF. GEORGE MACLOSIE, OF PRINCETON, N. J.

The endocranium of insects is produced by infoldings of the cranial wall, and although several groups (as Diptera, Hemiptera, Coleoptera, Lepidoptera,) have been represented as devoid of such structures, Prof. Macloskie finds an endocranium present in all these orders. The posterior or epicranial part of the skull has no internal processes. The clypeus, or "face," has a thick posterior ridge (just in advance of the antennae). From this ridge descend, in bees and allied insects, two meso-cephalic pillars, reaching to the floor of the cranium, in front of the great foramen. These two pillars support the roof of the skull. They occur, with variations, in squash-bug, gaddy, mosquito, butterfly, and dragonfly. In the cockroach they take the form of a perforated plate, being united anteriorly by a cross-bar (which binds the mandibles together), and being webbed excepting at the centre. (Huxley's description of this in his *Anatomy of the Invertebrates* is inaccurate.)

The maxillae and labium of the bee are supported by a long framework with elbows and hinges. This suspensorium is incorrectly represented in published figures. It is, in part correctly figured by Wolff, who misinterprets it (as if it were on the type of the mammalian skull). Its basal or posterior rods are attached close to the great foramen and to the base of the meso-cephalic pillars, and they are united by a thick web to the base of the skull. The mid-segment, consisting of a pair of bars, supports the maxillae, and upon it is an anterior pair of bars supporting the labium. In its working, this frame-work embodies the principle of a recent patent for producing steady motion.

The methods by which the maxillae and labium are protruded and withdrawn were described, also the relations and mode of working of the pharynx-parts in the mouth. The discovery of a double set of salivary glands was reported; a cephalic set supplying the inner tongue on the floor of the mouth, and the thoracic glands, sending their long duct forward to the labium. The inner structure of the bee's head was shown to be of the same pattern as in other insects, though varied in details. The paper was illustrated by diagrams and microscopic preparations.

NEW PLANETARY NEBULÆ.

BY PROFESSOR PICKERING.

He described the observations of the planetary nebulae, are now in progress at the Harvard College Observatory. Besides measuring the light of these bodies, the

spectrum of each has been examined by inserting a prism between the objective and eyepiece of the large telescope. A star is converted into a colored line of light, but the nebulae, being nearly monochromatic, appears as a bright point. The difference is so marked that the idea suggested itself that by this means planetary nebulae might be discovered, whose disks are so small that they can not otherwise be distinguished from stars. A search was accordingly undertaken on the evening of July 13th, by sweeping or moving the telescope so that a great number of stars could be examined in a short time. In a few minutes such a nebula was found, which with an ordinary eye-piece might readily be mistaken for a twelfth magnitude star. A similar object was also detected on the next evening. After this, sweeps on several evenings failed to reveal any new nebulae, although it is estimated that the spectra of over a hundred thousand stars were examined.

On night before last, while continuing this work, an object with a remarkable spectrum entered the field. The light appeared to consist mainly of a band in the green, a line in the red and probably a fainter band in the yellow, the whole being superposed on a faint continuous spectrum. The new stars which blazed out in Corona in 1863 and in Cygnus in 1876, presented for a short time a similar spectrum, but with this exception the star noted above appears to be unique. It is too soon to form a theory regarding the nature of this body, as clouds interrupted the observations and barely allowed time for its identification. It proved to be the star known as Oeltzen 17681, and must therefore have had nearly its present brightness forty years ago.

The field for discovery by the method here given is far from being exhausted since, less than one hundredth part of the heavens has as yet been examined.

ON LAND SNAILS OF THE PALÆOZOIC PERIOD.

By DR. DAWSON, F. R. S., Principal of McGill University, Montreal.

The land snails occurring in the carboniferous and Devonian systems, of which six species are known, were noticed in detail. Two of these, *Pupa Bigsbyi* from the coal formation of Nova Scotia, and *Strophites grandæva* from the Erian (Devonian) of St. Johns, New Brunswick, were described for the first time. Four of the known species belong to the different subdivisions of the old genus *Pupa*, and two are helicoid or snail-like in form. They constitute a very isolated group of fossils, as none are known in older formations, and there are none newer till we reach the early Tertiary. Though all of somewhat distinct types, they all belong to one great family or sub-order of the *Pulmonifera*, and are all closely allied to types still living. All the species hitherto found are American, four being found in Nova Scotia and New Brunswick, and two in Illinois. The latter were discovered and described by the late Mr. Bradley. *Pupa vetusta*, the earliest known, was found in the material filling a hollow *Sigillaria*, by Sir Charles Lyell and Dr. Dawson in 1851. In the paper, which will probably appear in full in the *AMERICAN JOURNAL OF SCIENCE*, figures and descriptions of all the species are given, and their affinities and mode of occurrence are discussed.

FURTHER NOTES ON THE POLLINATION OF YUCCA AND ON PRONUBA AND PRODOXUS.

BY C. V. RILEY.

The author refers to the original paper on the Fructification of *Yucca* read at the Dubuque (1872) meeting of the Association and notices various criticisms since made upon its conclusions. The paper shows that none of these criticisms were warranted, and verifies the original observations and conclusions by subsequent experience. It points out the causes of error in that other writers have confounded related moths having similar general appearance but great structural differences and different habits. The characters of the Bogus *Yucca* Moth (*Prodoxus decipiens*), are given, and five new