

with the absolute size of the bird, some flight-hindering element not yet therein contained, increases. We might therefore put the question, whether equally rapid, and (comparatively) equally great contraction in a small bird. In fact, too, it is chiefly the larger birds that present the phenomenon of scaring, a condition in which, the body being maintained at the same height for a certain time, muscular work is saved by special arrangements. If soaring be an advantage, it must, in alternation with periods of active rise by means of rudder-like mechanism, be extensively utilized for the problem of a flying machine.

COLOR RELATIONS OF METALS.

In a paper on the color relations of copper, nickel, cobalt, iron, manganese, and chromium, lately read before the Chemical Society, Mr. T. Bayley records some remarkable relations between solutions of these metals. It appears that iron, cobalt, and copper form a natural color group, for if solutions of their sulphates are mixed together in the proportions of 20 parts of copper, 7 of iron, and 6 of cobalt, the resulting liquid is free from color, but is gray, and partially opaque. It follows from this that a mixture of any two of these elements is complementary to the third, if the above proportions are maintained. Thus a solution of cobalt (pink) is complementary to a mixture of iron and copper (bluish green); a solution of iron (yellow) to a mixture of copper and cobalt (violet); and a solution of copper (blue) to a mixture of iron and cobalt (red). But, as Mr. Bayley shows, a solution of copper is exactly complementary to the red reflection from copper, and a polished plate of this metal, viewed through a solution of copper salt of a certain thickness, is silver-white. As a further consequence, it follows that a mixture of iron (7 parts) and cobalt (6 parts) is identical in color with a plate of copper. The resemblance is so striking that a silver or platinum vessel covered to the proper depth with such a solution is indistinguishable from copper.

There is a curious fact regarding nickel also worthy of attention. This metal forms solutions, which can be exactly simulated by a mixture of iron and copper solutions; but this mixture contains more iron than that which is complementary to cobalt. Nickel solutions are almost complementary to cobalt solutions; but they transmit an excess of very yellow light. Now, the atomic weight of nickel is nearly the mean of the atomic weight of iron and copper; but it is a little lower, that is, nearer to iron. There is thus a perfect analogy between the atomic weights and the color properties in this case. This analogy is even more general, for Mr. Bayley states that in the case of iron, cobalt, and copper, the mean wave length of the light absorbed is proportional to the atomic weight. The specific chromatic power of the metals varies, being least for copper. The specific chromatic power increases with the affinity of the metal for oxygen. Chromium forms three kinds of salts. Pink salts, identical in color with the cobalt salts; blue salts, identical in color with copper salts; and green salts, complementary to the red salts.

Manganese, in like manner, forms more than one kind of salt. The red salts of manganese are identical in color with the cobalt salts, and with the red chromium salts. The salts of chromium and manganese, according to the author, are with difficulty attainable in a state of chromatic purity. He thinks these properties of the metals lead up to some very interesting considerations.

FIRE AND WATER-PROOF PAPER.—A mixture is made of two-thirds ordinary paper pulp, and one-third asbestos. The whole is then steeped in a solution of common salt and alum, and after being made into paper is coated with an alcoholic solution of shellac.

DETECTION OF STARCH-SUGAR MECHANICALLY MIXED WITH COMMERCIAL CANE-SUGAR.*

By P. CASAMAJOR.

In a previous communication on the same subject,† read before the American Chemical Society at the meeting of March, 1880, I gave several processes for the detection of starch-sugar in commercial sugars. One of these consisted in adding to the suspected sugar a quantity of cold water, somewhat less than its own weight, and stirring the mixture for a few seconds. If starch-sugar is present, it will be seen in the shape of white chalky specks.

Quite lately a sample of yellow refined sugar was given to me which was supposed to be adulterated by being mixed with starch glucose. By applying the test just mentioned, there seemed to be left a few small chalky specks, which dissolved after standing a minute or two, making it very uncertain whether any starch glucose was present. Upon repeatedly trying the same test the result was always doubtful.

I was then lead to treat the suspected sugar by a liquid capable of dissolving sugar, but without any solvent action on starch-glucose. After many trials, I found that methylic alcohol of such density as to mark 50° by Gay-Lussac's alcohometer answered the purpose very well, if previously saturated with starch-sugar, as this solution dissolves cane-sugar, either white or yellow, very readily, but does not dissolve starch-glucose.

Methylic alcohol at 50°, saturated with starch-sugar, gives a solution of specific gravity = 1.25. 100 c.c. of methylic alcohol at 50° dissolves 57 grms. of dry starch-sugar, the volume of the solution being 133 c.c. A solution of starch-sugar in ethylic alcohol does not answer so well, because ethylic alcohol does not dissolve so readily the gummy matters found in soft sugars, which are those generally chosen for adulteration with glucose.

To test the presence of starch-sugar in a commercial cane-sugar, the suspected sugar should, in the first place, be thoroughly dried, as otherwise any water present will weaken the alcohol, and enable it to dissolve more starch-sugar. It should then be stirred for about two minutes with the saturated solution of starch-sugar in methylic alcohol. After this, the residue is allowed to settle, and the clear solution poured off. The residue may then be washed with a fresh quantity of the same solution. After stirring again and allowing the residue to settle, there will remain, if any starch-sugar is present, a certain quantity of chalky white specks, accompanied by a fine deposit, formed by the starch-sugar present in power of fine grains. These finer particles are never seen when water is used for detecting the presence of starch-sugar, as they dissolve in water very readily. It seems probable that by using this solution of starch-sugar in weak methylic alcohol, the starch-sugar in an adulterated sample could be estimated quantitatively by a process somewhat analogous to that of Payen for estimating cane-sugar.

Not having had any occasion for such a process I have not determined experimentally the degree of approximation obtainable in this way.

The methylic solution of starch-sugar should be poured on a filter, after it has dissolved all it can from a commercial sugar, and the residue should be washed out with the same solution, and everything poured on a weighed filter. After all the liquid has run off, the filter and the residue may be rapidly washed with the strongest methylic alcohol found in commerce, which tests 92½° by Gay-Lussac's alcohometer, and which dissolves starch-sugar with great difficulty.

By a dexterous use of this process it seems probable that very approximate results may be obtained, although what is said here is merely in the nature of a suggestion to those who may have use for quantitative results.

* A paper read before the American Chemical Society, Nov. 4, 1880.

† *Chemical News*, vol. xli., p. 221; *Journal of the American Chemical Society*, vol. ii., p. 111; *Sugar-Cane*, vol. xii., p. 283.