Herrick has remarked, "some of them have doubtless been concealed by clouds, and others witnessed only by barbarians.

But between the great display of B. C. 687 and A. D. 1803, Professor Newton gives the following list of show-ers at or near the epoch of April  $20,^{7}$  viz.: B. C, 15, A. D. 582, 1093, 1094,<sup>8</sup> 1095, 1096, 1122 and 1123. The appearance of 582 ought probably to be rejected. It was two days from the epoch, and the record as quoted by Outcleft much house no reference to choosing sterr<sup>8</sup>. The Quetelet may have no reference to shooting stars.9 The three remaining returns, B. C. 15, A, D. 1093-1096, and 1122-3, indicate a period of about 27 years. Now it is obvious that, at every close approach of meteors to the earth, many must be thrown into new orbits, all of which will pass through the point at which the perturbation occurred. It seems probable, therefore, that at some remote epoch a considerable cluster of this meteoric stream was thrown by perturbation into a new orbit corresponding to a period of 27 years. The change may have been produced by the earth during the passage of the meteoric swarm.

The facts which we have considered apparently indicate that the first comet of 1861, and the April meteors, formed a system in space before entering the solar domain; the latter moving in advance of the former at a distance comparable to the diameter of Neptune's orbit. By planetary perturbation the orbits were transformed into ellipses. If, as supposed by Professor Forbes, the disturbing body was an ultra-Neptunian planet in the vicinity of the present aphelion of the comet's orbit, said planet would probably describe less than 20° of its circuit during the interval between the nearest approach of the two bodies. But in aphelion the comet 1861 I, is too remote from the plane of the ecliptic to be sensibly disturbed by a planet moving in that plane. It seems more probable that the comet, as well as the meteoric group, owes the transformation of its orbit to one of the known major planets. Its radius vector when at its ascending node is about 10. In other words, its orbit approaches very near that of Saturn in longitude 30°. Now, it is remarkable that the interval between the perihelion passages of the meteors and the comet is almost exactly equal to two periods of Saturn. The meteors and Saturn were in the same longitude and in close proximity about B. C. 683, and the comet approached very near the planet at the same point about B. C. 625. The orbits may have been transformed into ellipses by Saturn's influence at these respective epochs. It may be worthy of remark that II times the period of the comet are equal to 155 times that of Saturn.

## CHEMICAL NOTES.

ON BALLO'S SUPPOSED ADIPIC ACID OBTAINED FROM CAMPHOR.-On oxidation with chromic acid camphor does not yield adipic acid, but the same oxidation-products as with nitric acid. Chromic acid, however, converts the camphoric acid first formed completely into members poorer in carbon.—J. KACHLER.

ON THE REMARKABLE REDUCING PROPERTIES OF POTAS-SIUM FERROUS OXALATE, AND ON SOME OF THE REACTIONS THUS PRODUCED .- Ferrous oxalate is very permanent on exposure to the air, both in a wet and a dry state, and pos-sesses very feeble reducing properties. The solution of ferrous oxalate in potassium oxalate, as well as the solid double salt, takes up oxygen greedily, and passes into po-tassium ferric oxalate. Its affinity for oxygen is equal to that of an alkaline ferrous hydrate, or of ammoniacal cuprous chloride, or of pyrogallic acid in an alkaline solution. The double oxalate exerts its reducing powers, not merely in alkaline, but in neutral, and even acid solutions. The solution quickly reduces platinum chloride and silver nitrate to metal. Silver chloride, bromide, and iodide are reduced completely, but more slowly. Copper acetate is reduced very slowly to cuprous oxide, and even to metal With the aid of heat mercuric chloride is reduced to metal. Recently precipitated Prussian blue is reduced to white ferro-cyanide of potassium. Indigo blue is reduced to white indigo, and solutions of sulphindigotic acid are rapidly decolorised.—J. M. EDER.

On the Acids  $C_8H_{14}O_4$  Formed from Butyric Acid. Besides a volatile oily acid, probably identical with isocro-tonic acid, there are formed by the reaction of suberic and bromobutyric acid, two acids agreeing in composition with suberic acid, but distinctly different from each other, and from the two isomeric acids produced by a corresponding reaction with brom-isobutyric acid. There exist, therefore, five isomeric suberic acids .- CARL HELL AND O. MUL-HAUSER.

A NEW SYNTHESIS OF PHOSPHENYL SULPHO-CHLORIDE.-Twenty parts phosphenyl-chloride are placed in a small flask with a reflux condenser, and five parts sulphur-chloride are slowly added by means of a dropping-funnel. After the reaction is over, the flask is set in a freezing mixture of Glauber's salt and hydrochloric acid. Pale yellow crystals of phosphenyl-tetra-chloride are formed, from which the liquid is separated by decantation, then shaken with water, dried and rectified. The yield is almost quantitative.-H. KOEHLER.

More Particular Observations on the Action of Pot-ASSIUM CARBONATE UPON ISOBUTYL-ALDEHYDE.-F. Urech places about 3 grms. pure isobutyl-aldehyde in a narrow test tube graduated in half millimetres. With a lens it is pos sible to read accurately quarter millimetres. After 3 decigrms. of finely-powdered recently-ignited potassium carbonate have been added, the tube is closed, set in a horizontal position, and the level is read off every five minutes for forty-eight hours. The liquid will be found to have sunk from 21.50 to 14.50 degrees.

AT a meeting of the Société Industrielle of Mulhouse, it was stated that tin sulphocyanide, formed by the double decomposition of calcium sulphocyanide and tin oxalate, is found very useful in calico printing.

For printing cotton with the azo-colors, Dr. Allrich proposes to dissolve 100 grms. of the color in five times is weight of water; then to make up a solution of solutin stannate or aluminate at  $15^{\circ}$  B, to every litre of which are added 20 grms, alizarin oil. Of this mixture 150 grms, are incorporated with the color, which is then thickened with starch and printed. After printing the pieces are steeped for an hour in lead or barium acetate or barium chloride at  $5^{\circ}$  to  $10^{\circ}$  B., and washed in cold water.

## CORRESPONDENCE.

[The Editor does not hold himself responsible for opinions expressed by his correspondents. No notice is taken of anonymous communi-cations.]

To the Editor of "SCIENCE:"

In reference to the correction of one of my statements made in your issue of the 29th inst. by Dr. Burt G. Wilder, I would say that I accept the criticism in all its bearings. The view which Dr. Wilder expresses regarding the upper wall of the third ventricle being constituted by the *ependyma* stretched across between the *habenulæ* of the pineal gland, was once entertained by myself (in accordance with the orthodox view of embryologists since the time of Rathke), and was the one which Dr. Wilder may year. I return to that view again. My abandonment of it was due to the confounding of two distinct questions, *i. e.*, the question of the true inner boundary of the floor at the lateral ventricle and the true upper and outer boun-dary of the third. The view I should have credited to Wilder and Hadlich, is that the *lateral ventricle does not* extend over the thalamus. My misapprehension of Wilder's statement is based on the fact that it rested on a verbal communication. That I mentioned it at all was

<sup>&</sup>lt;sup>7</sup> Am. Jour. of Science, July, 1863. <sup>8</sup> "At this period, so many stars fell from heaven that they could not be counted. In France the inhabitants were amazed to see one o themsfi great size fall to the earth, and they poured water on the spot, we hish their exceeding astonishment smoke issued from the ground with aoot f-ing noise."—Herrick's Catalogue. This record is of great interest as indr cating the fall of an aerolite during the shower of meteors. <sup>9</sup> "A Soissons, on voit le ciel en feu. Une pluie de sang tombe sur Paris."

due to my unwillingness to adopt even unpublished views of others without at least giving formal credit. If my good intentions have in this instance had rather the effect of doing an injustice, that was not a matter of intent, but of error, and one I may claim an excuse for under the circumstances above explained.

Dr. Wilder refers to the absence of literary references in my short paper. This is in accordance with the fact that the paper was a provisional communication of results purely my own, and that quotations of the views of others were limited to such relating to collateral details or of works like those of Luys and Meynert, which are so much household words among cerebral anatomists that I thought a mention of their exact titles a useless affectation.

On reviewing the matter, I find that the relations are in every respect as Dr. Wilder gives them, viz.: Ist. That the lateral and third ventricles only communicate through the Foramen of Monro. 2d. That the medial aspect of the Thalamus only is in the third ventricle. 3d. That no part of the Thalamusis in the lateral ventricles. 4th. That the dorsal aspect of the Thalamus is neither in the third nor in the lateral, nor in any other ventricle. My description of the third ventricle as a T shaped slit,

My description of the third ventricle as a T shaped slit, requires this modification, that the vertical branch alone is ventricular, the horizontal being neither a part of the lateral nor of the third ventricle, but a *fissure* extending between the thalamus and the cerebral hemispheres.

While thus embryologically and morphologically the thalamus must be looked on as a distinct cerebral segment, yet in the higher mammalia (as I claimed some years ago) it has become practically a subsidiary ganglionic depot of the cerebral hemispheres, and thus assumes a position not without its analogies to that of the *Corpus* striatum; a view in full accord with that of Meynert.

In conclusion I would offer a formal apology for the erroneous interpretation I made of Dr. Wilder's views, and state that the communication I made was merely provisional, and intended to be followed up by more full essays on the different subjects mentioned therein, in which the necessary literary references will not be wanting.

I also trust that Dr. Wilder will, in his publication, enlighten us on the anterior termination of the "extraventricular" slit which extends over the thalamus and under the Fornix. I am unable to determine this question from my sections, which are chiefly made in the transverse direction, and in which the ependymal roof of the third ventricle was either absent or removed by the knife. This feature was in part responsible for my misapprehension of the shape of the third ventricle. Respectfully,

NEW YORK, February 3d, 1881.

E. C. Spitzka.

# RECURRING PERIODS IN THE WEATHER.

#### To the Editor of "SCIENCE:"

Last year I published in the New York Tribune a table for the month of February showing the dates of rain and snow in that month for twenty years. I also pointed out that there seemed to be two well marked storm periods, about the 3rd and 20th of that month, when a storm might be expected. Since then I have made some corrections in the dates as then published and have extended my investigation to all the rest of the months. The result will be found in the table below; the table gives the dates of rain or snow as observed at Newburgh, N. Y. In all cases the blanks indicate that I had no observations for those dates. I would be very much obliged to any of the readers of "SCIENCE" who may have observations for these missing dates if they would publish them in "SCIENCE." For example, did it rain about 3rd of August 1862, May 1877, August 1873, August 1874, August 1876 and 1877, or about September 3, 1869? The missing dates from the 20th I would also be very much pleased to have supplied. An examination of the table will show that there seems to be two well marked storm periods in each month, and that a storm may be expected about the 3rd and 20th of every month. The crucial test for any theory concerning the weather is to see how accurately it predicts for any one place. It will readily be comprehended that the larger the area we predict for, greater is the probability of our predictions coming true, for if it does not storm at one place it may at another. Had I felt justified in using observations made in the vicinity of Newburgh, the table would have agreed much closer with theory, but I have only used the dates on which storms occurred at Newburgh. Besides, when the storm was very light near the 3rd or 20th, I have in some cases omitted it and recorded the date of the next storm. Thus, for example, it snowed at Newburgh De-cember 21, 1880, but it did not last long, and I have recorded the heavier storm of December 25th; and in September, 1864, it rained before the 30th, but as it was very light, I omitted it and recorded the first rain of any amount after the 20th. However, my object will be ac-complished if this article should direct the attention of observers to the importance of paying more attention to the recurring periods in the weather, on which Vennor has founded his predictions which have proved so true in many instances.

# JAMES H. GARDINER.

NEWBURGH, N. Y.

Year.	Jan.	Feb.	March.	April.	May.	June.	July.	August.	Sept.	Oct.	Nov.	Dec.
1860	3 20 5 20 2 21 1 19 4 19	3 18 3 20 3 19 1 20 1 24 5 26 2 19 3 20 9 19 3 18 5 20 9 19 3 18 5 20 9 19 3 20 3 21 3 20 3 21 1 21 5 26 3 21 3 20 3 21 3 20 3 21 3 20 3 21 3 19 4 19 5 26 5 26 9 19 9 19 3 10 9 19 3 10 9 19 3 10 9 19 3 10 9 19 3 20 3 20 3 20 3 21 3 20 3 20 3 21 3 21 3 20 3 21 3 21 3 21 3 20 3 21 3 21 3 21 3 20 3 21 3 21 3 21 3 21 3 20 3 21 3 21	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	I 21 3 24 2 20 4 23 2 25 6 20 7 22 2 20 I 20 4 21 4 20 I 21 3 21 4 19 3 18 3 20 2 20 2 3 19 3 20	3 19 1 25 2 19 4 17 3 21 1 18 2 20 6 19 2 21 3 19 7 24 19 3 21 2 19 4 17 3 21 2 19 4 17 3 21 2 19 4 17 3 21 1 18 2 20 6 19 2 21 1 18 2 20 6 19 2 21 3 19 2 21 3 19 4 17 3 21 1 18 2 20 6 19 2 21 3 21 2 19 2 21 3 19 2 21 3 21 2 19 2 21 3 19 2 21 3 21 2 19 2 21 2 21 2 19 2 21 2 20 5 19 2 22 2 2 2 22 2 2 2	I 20 2 19 2 21 I 20 9 19 3 22 2 18 2 21 I 20 7 .20 4 25 6 23 3 20 4 25 6 23 3 20 5 21 4 22 2 20	3 21 6 20 5 20 2 25 1 21 4 20 4 20 4 20 3 24 3 25 3 26 3 28 4 18 2 19 4 18 2 20 4 18 2 20 4 18 2 20 1 17 4 22 3 20 2 20 2 20 2 20 2 20 2 20 2 20 2	4 21 8 22 8 22 3 21 2 21 3 22 4 21 4 25 5 6 21  1 20 4 19 3 20 3 20 3 20 3 20 3 20 3 20 3 20 4 21 4 21 5 6 21 1 9	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	2 I4 5 I9 4 I9 2 28 4 I9 I4 22 5 21 I 21 3 28 II 22 2 18 3 20 2 21 I 16 5 21 I 16 5 21 4 20 2 21 I 16 5 21 4 21 6 I9 9 19 9 4 22	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$