

reply to Mr. WARINGTON the speaker said that the acetic acid fermentation went on in the presence of chloroform.

Mr. KINGZETT called attention to the fact that the oxygen was completely used up when the meat infusion was placed in contact with air. He did not think the experiments represented the action of bacteria on gases or of gases on bacteria, but rather the effects of various gases on the mode and extent of ordinary putrefaction.

Dr. FRANKLAND expressed his satisfaction with the results obtained by the author in his laborious research. He must confess that these results had surprised him not a little. The fact that bacteria, which were real organisms and could not be shielded under the term putrefaction, lived and flourished in SO_2 , CO , CN , &c., seemed to him very extraordinary, and the question arose whether the germs to which infectious diseases were probably due were not similarly endowed with a power of great resistance to ordinary influences.

Mr. F. J. M. PAGE said that Dr. Baxter had proved that with some fever-producing liquids, their virulence was destroyed by chlorine and sulphuric acid, and that he had seen some experiments at the Brown Institution which led to the same conclusion; so it seemed that, at all events in some cases, the virulence of infective liquids was due to organic matter, essentially different from the bacteria observed by Mr. Hatton.

NOTES ON CHICKEN CHOLERA.

We observe in a recent number of the *Chemical News* that C. T. Kingzett, F. C. S., points out, that, in explaining the protective influence of repeated inoculations with the attenuated virus of chicken cholera, against the more virulent forms of this disease, Pasteur finds it "impossible to resist the idea that the microscopic germ which causes the disease, finds in the body of the animal conditions suitable to its development, and that to satisfy the necessities of its life the germ alters certain substances, or destroys them, which comes to the same thing, whether it assimilates them or whether it consumes them with oxygen borrowed from the blood."

So, again, in cases where complete immunity has been attained, the birds "no longer contain food for the germ."

More striking still is the following passage in reference to chickens which are born proof against cholera:—"Animals in this condition may be said to be born vaccinated for this disease, because the foetal evolution has not placed in their bodies the proper food of the parasite, or because substances which would serve as such food have disappeared while they were yet young."

Now whether or not we may be prepared to regard the said parasite as the direct cause of the disease, it is remarkable that the reasoning of Pasteur should have culminated in the conclusion upon which Liebig insisted with considerable power.

If we turn to Gregory's (3rd) edition of Liebig's "Animal Chemistry" (p. 205) we find the following passage:—"The condition which determines, in a second individual, his liability to the contagion, is the presence in his body of a substance which by itself, or by means of the vital force acting in the organism, offers no resistance to the cause of change in form and composition operating on it. If this substance be a necessary constituent of the body, then the disease must be communicable to all persons; if it be an accidental constituent, then only those persons will be attacked by the disease in whom it is present in the proper quantity and of the proper composition. The course of the disease is the destruction and removal of this substance: it is the establishment of an equilibrium between the cause acting in the organism which determines the normal performance of its functions and a foreign power by whose influence these functions are altered."

I repeat that to me it seems somewhat remarkable that the investigations and reasoning of two such eminent (and

in many matters diametrically opposed) thinkers should have culminated in the same conclusion as regards the conditions of the living body which subject it to, or protect it from, infection.

While, however, it can be readily understood how a profuse growth of parasites could quickly alter or destroy a comparatively large amount of substance—as, for instance, happens in ordinary putrefaction—it does not appear to me so easy to accept Pasteur's reasoning as to his so-called vaccination.

In this inflicted process an attenuated virus is introduced into the body of a chicken which becomes ill but does not die. It does not die because, if Pasteur be correct, the parasites do not sufficiently multiply. Why do they not multiply? It cannot be on account of the insufficiency of the pabulum, for in the large majority of cases where death results this seems to arise from the very profusion of the growth of the parasite when more freely introduced.

Can it be expected, therefore, that even, say, in three successive inoculations the substance which I have here spoken of as pabulum can be entirely removed or destroyed by the very limited number of parasites which are introduced by the inoculations, and which so soon perish in the body? I think this cannot be expected; but if it may be, then the particular substance or substances upon which the parasites prey must be extremely limited in quantity. After all, we are faced with the enormous difficulty of ascertaining the nature of such substance, and the further equally great difficulty of understanding why an undiscovered and undetermined substance should be entirely absent from the bodies in some animals and present in varying proportions in others.

Here we come in contact with the weakest point in the parasitic theory. The immunity from a second attack of an infectious disease of the class in question is simply inexplicable under the parasitic theory. We are forced back to an alternative theory, and that is one of which we at present only recognize the beginnings.

A NEW CORTICAL CENTRE.*

By GRAEME M. HAMMOND, M.D., NEW YORK.

Physician to the Department for Diseases of the Nervous System in the Metropolitan Throat Hospital.

Some six years ago there appeared in the *Centralblatt*, Nos. 37 and 38, a short communication by Betz, embodying an account of certain nerve-cells found by him in the cortex of a region of the brain which he newly named the paracentral lobule. This paper has probably aroused more general attention among neurologists than any other paper of recent times dealing with the structure of the cerebral hemispheres, and this, on account of the anatomical confirmation which the discovery seemed to furnish, of the localization doctrine based on the electrical stimulation of the cortex carried out by Hitzig and Fritzsche.

After localizing these cells chiefly in the paracentral lobule and the upper ends of the pre- and post-central gyri of man, stating them to be very few in number in the lower halves of these gyri, Betz proceeds to say, "the constancy of the occurrence of these cells, not only as regards the cortical layer, but also the special convolutions in which they are found, led me to direct my attention to that portion of the brain of animals, and particularly of the dog, on which latter Hitzig and Fritzsche obtained such brilliant physiological results. I refer to that lobule which bounds the sulcus cruciatus. Now I found in this very lobule in the dog, cells in similar nests and of a similar shape. With the dog as in man they are distributed in the fourth layer."

Engaged in a study of the ganglionic masses of the forebrain of the cat, an animal on which the experiments of Hitzig and Fritzsche have been repeated, and in which

* Read before the New York Neurological Society, February 1, 1881.