with special precautions, viz., washing the respired gases, and performing parallel experiments, in which, for the breath I substituted a rapid current of air, and lastly raising the latter to a temperature of  $40^{\circ}$  C. The result was to prove conclusively my original statement that the decomposition is brought about by a constituent of the respired air, and therefore by its carbonic acid. In performing this experiment it is only necessary to secure the neutrality of the solution; this being done, the development of a full purple color occupies from two to three minutes.

It is evident that this demonstration of the presence of *some* acid body precedes the lime-water test in the logical development of the complete proof of the presence of *carbonic* acid.—*Chemical News*.

## THE BEST METHOD OF MOUNTING WHOLE CHICK EMBRYO.\*

#### By Dr. CHARLES S. MINOT.

The blastoderm is removed and cleaned in the usual manner, and then floated out on a glass slide, where it remains permanently. It is carefully spread out and allowed to dry until the edges become glued to the slide. It is then treated with a 0.5 per cent osmic acid solution, until a slight browning occurs. Stain with picro-carmine. The next step is particularly important, because it prevents the further darkening by the osmium, which otherwise injures or ruins the specimen. Pour Müller's fluid, or 0.5 per cent chromic acid solution, on the slide, and leave it over night. The next morning the blastoderm is ready for dehydration by alcohol, and mounting in the usual manner in balsam or dammar lac. Embryos prepared in this manner make particulary beautiful specimens.

# ON THE ALLEGED DECOMPOSITION OF THE ELEMENTS.†

#### BY PROF. DEWAR, M.A., F.R.S.

In his remarks Prof. Dewar dealt chiefly with the spectroscopic work from which Mr. Norman Lockyer had drawn conclusions very different from those of Professors Liveing and Dewar, especially concerning the value of evidence on the subject. Prof. Dewar argued that Mr. Lockyer's views regarding the existence of carbon vapor in the corona of the sun would not bear scientific investigation, and that his views regarding the modification of the spectrum of magnesium were equally illusory, and gave no proof of the decomposition of elementary sub-stances. Finally he discussed Mr. Lockyer's theory of "basic lines," and addressed himself to a refutation of the same. The results recorded, he said, strongly con-firmed Young's observations, and left little doubt that the few as yet unresolved coincidences either would yield to a higher dispersion, or were merely accidental. It would indeed be strange it amongst all the variety of chemical elements and the still greater variety of vibrations which some of them were capable of taking up, there were no two which could take up vibrations of the same period. They certainly should have supposed that substances like iron and titanium, with such a large number of lines, must each consist of more than one kind of molecule, and that not single lines, but several lines of each, would be found repeated with the spectra of some other chemi-cal elements. The fact that hardly a single coincidence could be established was a strong argument that the materials of iron and titanium, even if they be not homo-geneous, were still different from those of other chemical elements. The supposition that the different elements might be resolved into simple constituents and even into a single substance had long been a favorite speculation

with chemists; but however probable that hypothesis might appear *a priori*, it must be acknowledged, according to Prof. Dewar, that the facts derived from the most powerful method of analytical investigation yet devised, gave it but scant support.

#### ASTRONOMY.

### ELEMENTS AND EPHEMERIS OF COMET (d'), 1881.— BARNARD.

Mr. S. C. Chandler, Jr., has computed the following elements and ephemeris of Comet (d), 1881—Barnard—which are published, by permission of Prof. E. C. Pickering, of Harvard College Observatory. The observations upon which the computation is based are the following: Washington Mean Time being given with the Nashville observation, which was obtained at Vanderbilt University, by Prof. O. H. Landreth, and Cambridge Mean Time with the two others:

								R. A	. –	·	Decl	
1881.	Sept.	d. 20 21 25	h. 7 7	111. 46 34	s. 43	Nashville Harvard Obs. Harvard Obs.	h. 13 13	111. 28 30 26	5. 2 20 20.62	+3	/ 47 54 6	n 42.7
				~ ~	1.44						_	

The observation of the 20th was received by telegraph, and that of the 21st depends on only two comparisons, taken when the comet was but one degree and a half above the horizon.

#### ELEMENTS.

T = 1881, September, 14.785. Washington Mean Time.

$$\begin{array}{c} \pi = 271 & 22 \\ \Omega = 260 & 43 \\ i = 107 & 27 \\ \log \cdot q = 9.7053 \end{array}$$
 Mean Eq., 1881. o.

EPHEMERIS.

Wash	n. midnight,	_	R. A	. –	Dec	:l.—			
	1881.	h.	112.	<i>s</i> .	0	1	Log. $r$ .	Log. $\Delta$ .	Light
Sept.	29	13	41	36	+13	4	9.7894	0.1350	1.00
Oct.	3	τ3	45	28	16	26	9.8270	0.1467	.80
	7	13	48	40	19	29	9.86.48	0.1569	.65
	II	13	51	32	22	18	9.9014	0.1628	.52

The light of the comet on September 29 is taken as unity, and in this scale its light at discovery, on September 17, was 1.85. The orbit does not resemble that of any known comet. The comet is circular, not over one minute of arc in

The comet is circular, not over one minute of arc in diameter, with a very decided central condensation. Its collective brightness is not more than equivalent to that of an  $8\frac{1}{2}$  mag. star. The comet is rapidly decreasing in light, and the moon is advancing, so that observations of it at once are very desirable. So far as is known, positions have been obtained only at Nashville and Cambridge, the early setting of the comet, and clouds, having greatly interfered. Under the circumstances, the orbit cannot be other than a rough one, and considerable latitude for error had better be allowed in searching for it.

#### MICROSCOPY.

The following method of hardening the spinal cord for microscopic sections has been highly recommended by Dr. M. Debove :

Place the cord in a 4 per cent solution of bichromate of ammonia for three weeks, then in a solution of phenic gum for three days, and for three days more in alcohol. Sections may then be cut with great facility. They should be placed in water to prevent curling. They are then immersed in a saturated solution of picric acid for twenty-four hours, and colored with carmine for about twenty minutes, the picric acid acting as a mordant.— *Archives de Neurologie*.

An era of microtomes appears to be approaching, and numerous are such devices which are advertised by the opticians. Mr. Thomas Taylor of the Agricultural de-

<sup>\*</sup> Read before the A. A. A. S., Cincinnati, 1881.

<sup>†</sup> British Association, 1881.