

The second, more portable form, does not allow such an accurate quantitative air analysis. The instrument is attached to a weathercock, and thus is always directed against the wind, which traverses it, and deposits, as in the other permanent form, its solid matter on a glycerine plate. An anemometer placed in the vicinity serves to give an approximate idea of the quantity of air which has passed through the apparatus. These instruments have been called *aërosopes* by their inventor. It is likely that the second form given to the apparatus will be best fitted for the purpose which the Committee has in view.

#### THE NEW ASTRONOMER ROYAL.

Mr. William Henry Mahony Christie, who has succeeded Sir George Airy in the office of Astronomer Royal at the Royal Observatory, Greenwich Park, was born on October 1, 1845, at Woolwich. He is a younger son of the late Professor S. H. Christie, of the Royal Military Academy, Woolwich, and formerly Secretary to the Royal Society. Mr. W. H. M. Christie was educated at Kings College School, London; and at Trinity College, Cambridge, which he entered in 1864, having won a Minor's Scholarship of that College; he subsequently gained a Foundation Scholarship and was afterwards elected a Fellow of Trinity College. He took his degree of B.A. in 1868, as fourth wrangler in the Mathematical Tripos, and in 1871 proceeded to the M. A. degree. In 1870, Mr. Christie was appointed Chief Assistant at the Royal Observatory; and he has, during the past ten years, done special good service by contriving and introducing several valuable improvements in the scientific apparatus there in use; a new form of spectroscope, an instrument for determining the colors and brightness of the stars, a recording micrometer, and a polarising solar eye-piece, are to be mentioned as his inventions. In the recent address of the President of the British Association, at York, a passing reference was made to Mr. Christie's work in verifying the results obtained by Dr. Huggins, with regard to the motions of stars, as inferred from spectroscopic observations. The new Astronomer Royal has directed particular attention, at the Royal Observatory, both to spectroscopy and to photography, as a means of recording the observations. He is a fellow of the Royal Society, and was elected Secretary of the Royal Astronomical Society last year. He contributed to the proceedings of the Royal Society, in March, 1877, a paper "on the magnifying power of the half-prism, as a means of obtaining great dispersion, and on the general theory of the half-prism spectroscope." To the monthly notices of the Royal Astronomical Society, he has furnished these: in June, 1873, a paper on the recording micrometer; in January, 1874, on the color and brightness of stars, as measured with a new photometer; in May, 1875, on the determination of the scale in photographs of the Transit of Venus; in 1876, (January) on a new form of solar eye-piece; (May) on the displacement of lines in the spectra of stars; (November) on the effect of wear in the micrometer screws of the Greenwich Transit Circle; same year (December) on the gradation of light on the disk of Venus; in 1878 (January) on specular reflection from Venus; (June) on the existence of bright lines in the solar spectrum; in 1879 (January) on a phenomē-

seen in the occultation of a star by the moon's bright limb; in 1880, November, on the spectrum of Hartwig's comet of that year; in 1881 (January) on Mr. Stone's alterations of Bessel's refractions; (May) on the flexure of the Greenwich transit circle, and some further remarks on Mr. Stone's alterations of Bessel's refractions; besides various papers on the Greenwich spectroscopic and photographic observations, communicated by the late Astronomer Royal; and a paper which will be found in the Memoirs of the Royal Astronomical Society, published in January, 1880, on the systematic errors of the Greenwich North Polar distances. Mr. Christie is also the founder and editor of a journal entitled "*The Observatory*, a Monthly Review of Astronomy," which has been published during the past four years; and he is author of the "*Manual of Elementary Astronomy*," published in 1875 by the Society for Promoting Christian Knowledge.

#### ON THE ELECTRIC CONDUCTIVITY AND DICHROIC ABSORPTION OF TOURMALINE.\*

By Prof. SILVANUS P. THOMPSON.



WILLIAM H. M. CHRISTIE.

The electric conductivity of tourmaline differs in different directions; being, according to the author's experiments, a minimum along the optic axis. Tourmaline also possesses the optical property of dichroism, its absorption being a maximum for rays parallel to the axis, and greater for blue rays than for red, equal thicknesses of crystal being considered. According to the electro-magnetic theory of light, bodies which are good conductors of electricity should be opaque to light. The author has in the August number of the *Philosophical Magazine* rewritten the equations of Maxwell's electro-magnetic theory for the case of crystalline media possessing different conductivities in different directions. From these equations it appears that in tourmaline and negative uniaxial crystals

electric displacements at right angles to the axis will be more absorbed than electric displacements parallel to the axis. This accounts for the well-known greater absorption of the ordinary ray, provided the views of Stokes and Fresnel are correct, that these displacements are at right angles to the so-called plane of polarization. The difference of velocity between the rays of different color accounts for the difference of absorption being greater in that direction in which the conductivity is a minimum. It was also pointed out that in positive uniaxial crystals, in which the electric conductivity is a maximum along the axis, there will be maximum absorption of the extraordinary ray, and there will be least opacity along the axis. Smoky quartz and magnesian platinocyanide fulfil the latter condition. Specimens of tourmaline cut into cubes to show the colors in different directions were shown, and also specimens of magnesian platinocyanide and of herapathite. Mechanico-optical models were also shown illustrating the theory; a tourmaline being represented by a cube built up of layers of glass and wire-gauze. In conclusion it was shown that crystals in which the electric conductivity differs in three different directions will exhibit *trichroism*; and that di- or trichroic absorption is a general property of all colored crystals other than those of the cubical system.

\* British Association, 188r.