

known by their true scientific names. This is the case with *Eschscholtzia*, *Romneya*, *Clematis*, *Isomeris*, *Silena*, *Malva*, *Ceanothus*, *Hosackia*, *Ribes*, *Phacelia*, *Gilia*, and many others for which the generic name has become a popular name.

This is owing to various causes, one being the difficulty of applying the old familiar garden names; which are used, however, when any resemblance can be traced, as is the case with larkspur, honeysuckle, columbine, etc. Many of the settlers have also become familiar with the true names of these flowers by having received them from parties that have introduced them to cultivation, for which the greatest credit is due to the late James Vick.

Many visitors, as well as settlers, seek to learn the names of the many strange and beautiful flowers, that, by massing, become such a feature in the scenery, and find the 'dry' scientific names as easy to learn, and as sensible, as the old Spanish names, but few of which survive in the popular mind. Thanks to the little botany of Volney Rattan, largely supplemented by visiting and amateur botanists, all are enabled to learn the more common species with comparative ease.

C. R. ORCUTT.

The use of the method of rates in mathematical teaching.

In the case of the question, "Does change in the rate of motion take place at an instant, or during an interval?" I am surprised to find that Professor Wood (*Science*, May 16) regards my amendment as only increasing the difficulty. It may be that I have been misunderstood: permit me, therefore, to answer the questions which the professor goes on to ask in illustration of this difficulty. Assuming (of course, correctly) that my answer to the question is, that it takes time to produce a change in the rate of motion, he asks, "How long is this interval?" I answer, "As long as you please usually: of course, the longer the interval, the greater the change." — "If ever so small, is the rate variable during the interval?" — "Certainly." — "If variable, the original question arises, and we wish to know if *change* involves a *part* of the interval." — "Of course, a part of the change takes place in a part of the interval, and the rest of the change takes place in the rest of the interval." — "Does change in the rate take place at 'a point' in the path, or during 'a space' of the path?" — "During 'a space' of the path; that is, while the point is passing over a space of the path." — "If at 'a point,' is it not equivalent to asserting that a change takes place in no time? [Most certainly it would be, but we do not assert this at all.] And if an interval is necessary, must it not be conceived as infinitesimal?" — "By no means: if you want a finite change, and that is what is usually meant by a change, you must take a finite interval of time; but, if you insist on introducing the conception of an infinitesimal change, you must admit also an infinitesimal interval of time." Let us put precisely parallel questions with respect to the *position* of a moving point. Does change of position take place at an instant, or during an interval? During an interval. How long is this interval? That depends upon the amount of change of position you desire to produce. If ever so small, is the position of the point variable during the interval? Certainly, if the point moves. Does change of position take place at a point in the path? Certainly not: a point has position, but no magnitude.

If there is any difficulty in conceiving the velocity of a point to be continuously variable, there is precisely the same difficulty in conceiving the abscissa of a point moving on the axis of x to be continuously

variable; in other words, in conceiving the possibility of motion itself. It should be remembered that the definition of the measure of a variable velocity, presupposed in this discussion, is simply that which we find in such treatises as Tait and Steel's *Dynamics* of a particle: "Velocity is said to be variable when the moving point does not describe equal spaces in equal times. *The velocity at any instant is then measured by the space which would have been described in a unit of time, if the point had moved on uniformly for that interval with the velocity which it had at the instant contemplated.*"

WM. WOOLSEY JOHNSON.

Annapolis, May 19.

Pleuracanthus and Didymodus.

In your issue of April 11, my friend Professor Gill communicates his views on the relationships of *Pleuracanthus* and *Chlamydoselachus*, and endeavors to correct some of my opinions and statements. On some points I stand corrected, thanks to Professor Gill's superior knowledge of the literature of the subject. However, as Professor Gill has not seen my material, nor the paper which I read before the Philosophical society upon it, I may, in turn, enlighten him on some important aspects of the case.

Professor Gill objects to the identification of the genera *Didymodus* and *Chlamydoselachus* on the sole ground of the diversity in the form of the teeth. He probably has other reasons for objecting; but, with his usual magnanimity, he has not used his most effective weapons. He doubts the pertinence of the recent and extinct genera to the same order. He points out that the oldest name of the genus called *Diplodus* is *Pleuracanthus*, and that my order *Ichthyotomi* has been already defined and named by Lütken as the *Xenacanthini*.

On these positions, I make the following comments:—

1. There is no generic difference to be detected, in my opinion, between the teeth which are typical of *Diplodus* Agass. and *Thrinacodus* St. J. and W. and the recent *Chlamydoselachus*. Differences there are, but apparently not of generic value. The identification of the recent and extinct genera rests, as far as this point goes, on the same basis as that of the recent and extinct *Ceratodus*.

2. At the time of my proposal of the name *Didymodus*, I was not convinced that fishes of this type bore the spines referred to the genus *Pleuracanthus* Agass. None of the authors cited figure any specimens which present both tricuspidate teeth and a nuchal spine. None of my ten specimens possess a spine. However, Kner describes two specimens as exhibiting both tricuspidate teeth and a spine, and Sir P. Egerton's statements (*l.c.*) on this point are positive. So we must regard *Pleuracanthus* as the name of this genus, with *Diplodus* as a synonyme.

3. *Diplodus* being regarded as a synonyme of *Pleuracanthus*, it follows that *Chlamydoselachus* Garm. is distinct on account of the *different structure of the dorsal fin*, which is single and elongate in *Pleuracanthus*, according to Geinitz and Kner. The presence of the nuchal spine in *Pleuracanthus* is also, probably, a character of distinction, although we do not yet know whether such a spine is concealed in *Chlamydoselachus* or not.

4. The identity of *Didymodus* (type, *Diplodus compressus* Newberry) and *Pleuracanthus* may now be questioned. None of the specimens are figured and described by the authors above cited, as displaying an occipital condyle, or posterior frontal cornua. My specimens of *Didymodus compressus* do not exhibit