

early period in the life of the time-service, the telegraphic lines have been in charge of the electricians, Messrs. Stearns and George, and their successor, Mr. C. L. Bly.

EDWARD C. PICKERING,
Director of the Astronomical
Observatory of Harvard College.

Cambridge, Mass.

NOTES AND NEWS.

MR. J. L. KIPLING says of the monkeys of India: "They have a game like the English boys' cock of the dung-hill or king of the castle, but instead of pushing each other from the top of a knoll or dust-heap, the castle is a pendant branch of a tree. The game is to keep a place on the bough, which swings with their weight as with a cluster of fruit, while the players struggle to dislodge one another, each, as he drops, running round and climbing up again to begin anew. This sport is kept up for an hour at a time with keen enjoyment, and when one is nimble as a monkey it must be splendid fun."

— In 1890 was published the important discovery by Behring and Kitasato that blood serum taken from animals that had been rendered immune to tetanus and diphtheria was capable of curing other animals suffering from those diseases. Drs. G. and F. Klemperer (*Berliner klinische Wochenschrift*, Aug. 24 and 31, 1891) publish a research carried out in regard to pneumonia, with the object of discovering how immunity against the pneumococcus could be best produced, whether recovery from the disease rendered an animal immune, and whether it was possible to cure pneumonia by the blood serum of animals that have recovered from the disease. Their experiments, which were confined to rabbits, revealed that every nutrient medium in which the pneumococcus has been cultivated will, if inoculated, render an animal immune against pneumonic septicæmia, even after the cocci have been removed by filtration. The power of producing immunity is more speedily acquired, and is increased if the infected nutrient medium (before or after removal of the cocci) is exposed to a temperature of between 41° and 42° C. for two or three days, or of 60° for an hour or two. In every case, however, it was found necessary that some interval (varying from three to fourteen days) should elapse between the inoculation and the production of immunity. Hence it was too late to cure a diseased animal or even to prevent the onset of an attack if the injection was given simultaneously with the outbreak of the disease. On the other hand, serum taken from animals enjoying immunity was found able, especially when introduced directly into the circulation, to cure pneumonic septicæmia. The serum was injected twenty-four hours after infection, while the animals had a febrile temperature of between 105° and 106.5° F. Eight cubic centimetres were injected, with the result that the temperature gradually sank during the next twenty-four hours. In twelve successive cases a successful result was obtained. This research therefore confirms, in regard to pneumonia in rabbits, what Behring and Kitasato did for tetanus and diphtheria. Drs. Klemperer next studied the question how the blood serum of an immune animal cures an attack of pneumonic septicæmia, and discovered that when the pneumococcus is introduced into the body of an animal it generates a poisonous substance which can be isolated, and to which the name of "pneumotoxin" has been given. This pneumotoxin sets up a febrile condition which lasts several days, after which another substance is found to have been produced called "antipneumotoxin," which is able to neutralize the pneumotoxin. The serum taken from an immune animal contains this antipneumotoxin, and it is by means of this substance that it cures an attack of pneumonic septicæmia in other animals. The relation of pneumonia as seen in rabbits with that met with in man was next investigated, and the conclusion arrived at that the disease in both cases is produced by the pneumococcus, but that the human body is much less susceptible to the latter than the rabbit is. Thus it was found that serum taken from pneumonic patients after the crisis could cure pneumonia in rabbits; moreover, pneumotoxin and antipneumotoxin were found to be present in human serum as in that taken from rabbits. The crisis of pneumo-

nia, according to Drs. Klemperer, takes place as soon as a antipneumotoxin is produced in sufficient quantity to neutralize the pneumotoxin. Why immunity against further attacks lasts so short a time in man is still uncertain, but possibly less antipneumotoxin is formed in man than in rabbits in proportion to the pneumotoxin. Some attempts have already been made to cure patients suffering from pneumonia with the help of antipneumotoxin, but further observations are necessary.

— It is a well known fact that, with the same temperature by the thermometer, one may have, at different times, a very different feeling of heat and cold. This varies with the temperature of the skin, which is chiefly influenced (according to M. Vincent of Uccle Observatory, Belgium), by four things: air-temperature, air-moisture, solar radiation, and force of wind. M. Vincent recently made a large number of observations of skin-temperature in the ball of the left hand, and constructed a formula by means of which the skin-temperature may be approximately deduced from those four elements. He experimented by keeping three of the four constant, while the fourth was varied, and a relation could thus be determined between the latter and skin-temperature. One fact which soon appeared was, that the relative moisture of the air has but little influence on skin-temperature. It was also found that for every 1° C. of the actinometric difference (excess of black bulb thermometer) the skin-temperature rises about 0.2°; and with small wind-velocities, every metre per second depresses the skin-temperature about 1.2°. In testing his formula M. Vincent found, with cold or very cold sensation, considerably greater differences between the calculated and observed values than in other cases. This he attributes to the great cooling of the relatively small mass of the hand. Taking the cheek or eyelid the results were better, says *Nature*.

— Last winter, in December and January, M. Chaix made a number of observations of the temperature of the air, the snow, and the ground at Geneva, of which he has given an account to the Physical Society there. He observed the air at four different heights; granular, pulverulent, and bedded snow, on the surface and at different depths; and the surface of bare ground as well as of ground covered with snow. There was no difference in mean temperature between the air at one and two metres; and very little between the former and that on the snow surface. The surface of the ground was 4.265° C. warmer than the surface of the snow (0.13 m. above), through arrest of radiation. But the bare ground was not cooled so much as the snow surface, and it was only 2.04° colder than the snow-clad ground. This shows the frigorific influence of snow on climate. Air passing over bare ground would have been 2° warmer than if it passed over the snow. The snow surface was sometimes warmer, sometimes colder than the air one or two metres above. In the dry winters of Siberia and Sweden, the snow-surface is generally (according to Woeikof) much colder than the air. M. Chaix explains the variations observed at Geneva by fluctuations in the relative humidity, involving alternate vaporization and condensation at the snow-surface. In two-thirds of the cases, indeed, abnormal cooling of the snow corresponded with a low humidity, and heating with a high humidity, and often formation of hoar frost at the surface, according to *Nature*.

— An illustration of the height of breaking waves is afforded by the following paragraph, which we take from the *San Francisco Chronicle* of Jan. 6: "Portland, Jan. 5 The lighthouse tender 'Manzanita' reached Tillamook Rock Sunday for the first time in six weeks, and brought away the keeper, George Hunt, who has been on the rock for four years, and has been transferred to the Cape Mars Light. He says, in the storm of Dec. 7 the waves swept clear over the house, washing away their boats, and tearing loose and carrying away the landing platform and tramway, which were bolted to the rock. On the 29th the waves were still higher, and streams of water poured into the lantern through the ventilators in the balloon top of the dome, 157 feet above the sea-level. The lighthouse was shaken to its foundation by the impact of seas against it, and the water found its way into the house. Men were on duty all night to keep the lamp burning, and but for the wire screen the shutters of the lantern would have been demolished."