SCIENCE.

EDITORIAL COMMITTEE: S. NEWCOMB, Mathematics; R. S. WOODWARD, Mechanics; E. C. PICKERING, Astronomy; T. C. MENDENHALL, Physics; R. H. THURSTON, Engineering; IRA REMSEN, Chemistry; J. LE CONTE, Geology; W. M. DAVIS, Physiography; O. C. Marsh, Paleontology; W. K. Brooks, Invertebrate Zoölogy; C. Hart Merriam, Vertebrate Zoölogy; S. H. Scudder, Entomology; N. L. Britton, Botany; Henry F. Osborn, General Biology; H. P. Bowditch, Physiology; J. S. Billings, Hygiene; J. McKeen Cattell, Psychology; Daniel G. Brinton, J. W. Powell, Anthropology.

FRIDAY, JUNE 28, 1895.

CONTENTS:

Argon: LORD RAYLEIGH	701
Lloyd Morgan Upon Instinct: H. F. O	712
Some Meandering Rivers of Wisconsin: Henry Kümmel	B. 714
Correspondence: —	716
Scientific Literature:—	C.
Notes and News:— The Remedy for Pear Blight: M. B. WAITE. N York Botanic Garden; The Helmholtz Memoria General.	ew
Societies and Academies:— Biological Society of Washington; Entomologic Society of Washington; The New York Academ of Science; The Wisconsin Academy of Science Arts and Letters; The Texas Academy of Science	cal ny es,
Non Rooks	798

MSS. intended for publication and books, etc., intended for review should be sent to the responsible editor, Prof. J. McKeen Cattell, Garrison on Hudson, N. Y. Subscriptions and advertisements should be sent to Science, 41 N. Queen St., Lancaster, Pa., or 41 East 49th St., New York.

ARGON.*

It is some three or four years since I had the honour of lecturing here one Friday evening upon the densities of oxygen and hydrogen gases, and upon the conclusions that might be drawn from the results. It is not necessary, therefore, that I should

*A Lecture given by Lord Rayleigh before the Royal Institution of Great Britain, on Friday, April 5, 1895. Reprinted from the official report.

trouble you to-night with any detail as to the method by which gases can be accurately weighed. I must take that as known, merely mentioning that it is substantially the same as is used by all investigators nowadays, and introduced more than fifty years ago by Regnault. It was not until after that lecture that I turned my attention to nitrogen; and in the first instance I employed a method of preparing the gas which originated with Mr. Vernon Harcourt, of Oxford. In this method the oxygen of ordinary atmospheric air is got rid of with the aid of ammonia. Air is bubbled through liquid ammonia, and then passed through a red-hot tube. In its passage the oxygen of the air combines with the hydrogen of the ammonia, all the oxygen being in that way burnt up and converted into water. The excess of ammonia is subsequently absorbed with acid, and the water by ordinary desiccating agents. That method is very convenient; and, when I had obtained a few concordant results by means of it, I thought that the work was complete, and that the weight of nitrogen was satisfactorily determined. But then I reflected that it is always advisable to employ more than one method, and that the method that I had used—Mr. Vernon Harcourt's method —was not that which had been used by any of those who had preceded me in weighing The usual method consists in absorbing the oxygen of air by means of