

SCIENCE

EDITORIAL COMMITTEE : S. NEWCOMB, Mathematics ; R. S. WOODWARD, Mechanics ; E. C. PICKERING, Astronomy ; T. C. MENDENHALL, Physics ; R. H. THURSTON, Engineering ; IRA REMSEN, Chemistry ; JOSEPH LE CONTE, Geology ; W. M. DAVIS, Physiography ; HENRY F. OSBORN, Paleontology ; W. K. BROOKS, C. HART MERRIAM, Zoology ; S. H. SCUDDER, Entomology ; C. E. BESSEY, N. L. BRITTON, Botany ; C. S. MINOT, Embryology, Histology ; H. P. BOWDITCH, Physiology ; J. S. BILLINGS, Hygiene ; WILLIAM H. WELCH, Pathology ; J. McKEEN CATTELL, Psychology ; J. W. POWELL, Anthropology.

FRIDAY, SEPTEMBER 21, 1900.

CONTENTS :

<i>The Address of the President of the Section of Mathematics and Physics of the British Association for the Advancement of Science : DR. JOSEPH LARMOR.....</i>	417
<i>Inland Biological Laboratories</i>	436
<i>The Colorado Potato Beetle : DR. W. L. TOWER... ..</i>	438
<i>The Eighth International Geological Congress at Paris : H. F. O.....</i>	440
<i>Scientific Books :—</i>	
<i>The Davenports' Introduction to Zoology : DR. MAURICE A. BIGELOW. Herdman and Boyce on Oysters and Disease: DR. H. F. MOORE. Belzung's Anatomie et physiologie végétale : DR. D. T. MACDOUGAL. Report of Competitive Tests of Street Car Brakes : PROFESSOR R. H. THURSTON. Zenker's Lehrbuch der Photochromie : PROFESSOR R. W. WOOD. Books Received</i>	442
<i>Societies and Academies :</i>	
<i>Section of Geology and Mineralogy of the New York Academy of Sciences : DR. THEODORE G. WHITE.....</i>	446
<i>Discussion and Correspondence :—</i>	
<i>Mr. Tesla and the Universe.....</i>	447
<i>Botanical Notes :—</i>	
<i>A New Laboratory Manual ; Origin of the Higher Fungi ; Supplement to Nicholson's Dictionary of Gardening ; New Edition of Prantl's Lehrbuch : PROFESSOR CHARLES E. BESSEY.....</i>	451
<i>The Coal Fields of China</i>	452
<i>Scientific Notes and News.....</i>	453
<i>University and Educational News.....</i>	456

ADDRESS OF THE PRESIDENT OF THE MATHEMATICAL AND PHYSICAL SECTION OF THE BRITISH ASSOCIATION FOR THE ADVANCEMENT OF SCIENCE.

It is fitting that before entering upon the business of the Section we should pause to take note of the losses which our department of science has recently sustained. The fame of Bertrand, apart from his official position as Secretary of the French Academy of Sciences, was long ago universally established by his classical treatise on the 'Infinitesimal Calculus' : it has been of late years sustained by the luminous exposition and searching criticism of his books on the 'Theory of Probability' and 'Thermodynamics' and 'Electricity.' The debt which we owe to that other veteran, G. Wiedemann, both on account of his own researches, which take us back to the modern revival of experimental physics, and for his great and indispensable thesaurus of the science of electricity, cannot easily be overstated. By the death of Sophus Lie, following soon after his return to a chair in his native country Norway, we have lost one of the great constructive mathematicians of the century, who has in various directions fundamentally expanded the methods and conceptions of analysis by reverting to the fountain of direct geometrical intuition. In Italy the death of Beltrami has removed

an investigator whose influence has been equally marked on the theories of transcendental geometry and on the progress of mathematical physics. In our own country we have lost in D. E. Hughes one of the great scientific inventors of the age ; while we specially deplore the removal in his early prime, of one who has recently been well known at these meetings, Thomas Preston, whose experimental investigations on the relations between magnetism and light, combined with his great powers of lucid exposition, marked out for him a brilliant future.

Perhaps the most important event of general scientific interest during the past year has been the definite undertaking of the great task of the international coordination of scientific literature ; and it may be in some measure in the prolonged conferences that were necessitated by that object that the recently announced international federation of scientific academies has had its origin. In the important task of rendering accessible the stores of scientific knowledge, the British Association, and in particular this Section of it, has played the part of pioneer. Our annual volumes have long been classical, through the splendid reports of progress of the different branches of knowledge that have been from time to time contributed to them by the foremost British men of science ; and our work in this direction has received the compliment of successful imitation by the sister Associations on the Continent.

The usual conferences connected with our department of scientific activity have been this year notably augmented by the very successful international congresses of mathematicians and of physicists which met a few weeks ago in Paris. The three volumes of reports on the progress of physical science during the last ten years, for which we are indebted to the initiative of the French Physical Society, will provide an

admirable conspectus of the present trend of activity, and form a permanent record for the history of our subject.

Another very powerful auxiliary to progress is now being rapidly provided by the republication, in suitable form and within reasonable time, of the collected works of the masters of our science. We have quite recently received, in a large quarto volume, the mass of most important unpublished work that was left behind him by the late Professor J. C. Adams ; the zealous care of Professor Sampson has worked up into order the more purely astronomical part of the volume ; while the great undertaking, spread over many years, of the complete determination of the secular change of the magnetic condition of the earth, for which the practical preparations had been set on foot by Gauss himself, has been prepared for the press by Professor W. G. Adams. By the publication of the first volume of Lord Rayleigh's papers a series of memoirs which have formed a main stimulus to the progress of mathematical physics in this country during the past twenty years has become generally accessible. The completed series will form a landmark for the end of the century that may be compared with Young's 'Lectures on Natural Philosophy' for its beginning.

The recent reconstruction of the University of London and the foundation of the University of Birmingham will, it is to be hoped, give greater freedom to the work of our University Colleges. The system of examinations has formed an admirable stimulus to the effective acquisition of that general knowledge which is a necessary part of all education. So long as the examiner recognizes that his function is a responsible and influential one, which is to be taken seriously from the point of view of moulding the teaching in places where external guidance is helpful, test by examination will remain a most valuable means of

extending the area of higher education. Except for workers in rapidly progressive branches of technical science, a broad education seems better adapted to the purposes of life than special training over a narrow range; and it is difficult to see how a reasonably elastic examination test can be considered as a hardship. But the case is changed when preparation for a specialized scientific profession, or mastery of the lines of attack in an unsolved problem, is the object. The general education has then been presumably finished; in expanding departments of knowledge, variety rather than uniformity of training should be the aim, and the genius of a great teacher should be allowed free play without external trammels. It would appear that in this country we have recently been liable to unduly mix up two methods. We have been starting students on the special and lengthy, though very instructive, processes which are known as original research at an age when their time would be more profitably employed in rapidly acquiring a broad basis of knowledge. As a result, we have been extending the examination test from the general knowledge to which it is admirably suited into the specialized activity which is best left to the stimulus of personal interest. Informal contact with competent advisers, themselves imbued with the scientific spirit, who can point the way towards direct appreciation of the works of the masters of the science, is far more effective than detailed instruction at second hand, as regards growing subjects that have not yet taken on an authoritative form of exposition. Fortunately there seems to be now no lack of such teachers to meet the requirements of the technical colleges that are being established throughout the country.

The famous treatise which opened the modern era by treating magnetism and electricity on a scientific basis appeared just 300 years ago. The author, William Gil-

bert, M.D., of Colchester, passed from the Grammar School of his native town to St. John's College, Cambridge; soon after taking his first degree, in 1560, he became a Fellow of the College, and seems to have remained in residence, and taken part in its affairs, for about ten years. All through his subsequent career, both at Colchester and afterwards at London, where he attained the highest position in his profession, he was an exact and diligent explorer, first of chemical and then of magnetic and electric phenomena. In the words of the historian Hallam, writing in 1839, 'in his Latin treatise on the *'Magnet,'* he not only collected all the knowledge which others had possessed, but he became at once the father of experimental philosophy in this island'; and no demur would be raised if Hallam's restriction to this country were removed. Working nearly a century before the time when the astronomical discoveries of Newton had originated the idea of attraction at a distance, he established a complete formulation of the interaction of magnets by what we now call the exploration of their fields of force. His analysis of the facts of magnetic influence, and incidentally of the points in which it differs from electric influence, is virtually the one which Faraday reintroduced. A cardinal advance was achieved, at a time when the Copernican Astronomy had still largely to make its way by assigning the behavior of the compass and the dip needle to the fact that the earth itself is a great magnet, by whose field of influence they are controlled. His book passed through many editions on the Continent within forty years; it won the high praise of Galileo. Gilbert has been called 'the father of modern electricity' by Priestley, and 'the Galileo of magnetism' by Poggendorff.

When the British Association last met at Bradford in 1873, the modern theory which largely reverts to Gilbert's way of formula-

tion, and refers electric and magnetic phenomena to the activity of the æther instead of attractions at a distance, was of recent growth; it had received its classical exposition only two years before by the publication of Clerk Maxwell's treatise. The new doctrine was already widely received in England on its own independent merits. On the Continent it was engaging the strenuous attention of Helmholtz, whose series of memoirs, deeply probing the new ideas in their relation to the prevalent and fairly successful theories of direct action across space, had begun to appear in 1870. During many years the search for crucial experiments that would go beyond the results equally explained by both views, met with small success; it was not until 1887 that Hertz, by the discovery of the æthereal radiation of long wave-length emitted from electric oscillators, verified the hypothesis of Faraday and Maxwell and initiated a new era in the practical development of physical science. The experimental field thus opened up was soon fully occupied both in this country and abroad; and the borderland between the sciences of optics and electricity is now being rapidly explored. The extension of experimental knowledge was simultaneous with increased attention to directness of explanation; the expositions of Heaviside and Hertz and other writers fixed attention in a manner already briefly exemplified by Maxwell himself, on the inherent simplicity of the completed æthereal scheme, when once the theoretical scaffolding employed in its construction and dynamical consolidation is removed; while Poynting's beautiful corollary specifying the path of the transmission of energy through the æther has brought the theory into simple relations with the applications of electrodynamics.

Equally striking has been the great mastery obtained during the last twenty years over the practical manipulation of electric

power. The installation of electric wires as the nerves connecting different regions of the earth had attained the rank of accomplished fact so long ago as 1857, when the first Atlantic cable was laid. It was largely the theoretical and practical difficulties, many of them unforeseen, encountered in carrying that great undertaking to a successful issue, that necessitated the elaboration by Lord Kelvin and his coadjutors, of convenient methods and instruments for the exact measurement of electric quantities, and thus prepared the foundation for the more recent practical developments in other directions. On the other hand, the methods of theoretical explanation have been in turn improved and simplified through the new ways of considering the phenomena which have been evolved in the course of practical advances on a large scale, such as the improvement of dynamo armatures, the conception and utilization of magnetic circuits, and the transmission of power by alternating currents. In our time the relations of civilized life have been already perhaps more profoundly altered than ever before, owing to the establishment of practically instantaneous electric communication between all parts of the world. The employment of the same subtle agency is now rapidly superseding the artificial reciprocating engines and other contrivances for the manipulation of mechanical power that were introduced with the employment of steam. The possibilities of transmitting power to great distances at enormous tension, and therefore with very slight waste, along lines merely suspended in the air, are being practically realized; and the advantages thence derived are increased many fold by the almost automatic manner in which the electric power can be transformed into mechanical rotation at the very point where it is desired to apply it. The energy is transmitted at such lightning speed that at a given instant only an exceedingly minute

portion of it is in actual transit. When the tension of the alternations is high, the amount of electricity that has to oscillate backwards and forwards on the guiding wires is proportionately diminished, and the frictional waste reduced. At the terminals the direct transmission from one armature of the motor to the other, across the intervening empty space, at once takes us beyond the province of the pushing and rubbing contacts that are unavoidable in mechanical transmission; while the perfect symmetry and reversibility of the arrangement by which power is delivered from a rotatory alternator at one end, guided by the wires to another place many miles away, where it is absorbed by another alternator with precise reversal of the initial stages, makes this process of distribution of energy resemble the automatic operations of nature rather than the imperfect material connections previously in use. We are here dealing primarily with the flawless continuous medium which is the transmitter of radiant energy across the celestial spaces; the part played by the coarsely constituted material conductor is only that of a more or less imperfect guide which directs the current of æthereal energy. The wonderful nature of this theoretically perfect, though of course practically only approximate, method of abolishing limitations of locality with regard to mechanical power is not diminished by the circumstance that its principle must have been in some manner present to the mind of the first person who fully realized the character of the reversibility of a gramme armature.

In theoretical knowledge a new domain, to which the theory as expounded twenty years ago had little to say, has recently been acquired through the experimental scrutiny of the electric discharge in rarefied gaseous media. The very varied electric phenomena of vacuum tubes, whose electrolytic character was first practically estab-

lished by Schuster, have been largely reduced to order through the employment of the high exhaustions introduced and first utilized by Crookes. Their study under these circumstances, in which the material molecules are so sparsely distributed as but rarely to interfere with each other, has conduced to enlarged knowledge and verification of the fundamental relations in which the individual molecules stand to all electric phenomena, culminating recently in the actual determination, by J. J. Thomson and others following in his track, of the masses and velocities of the particles that carry the electric discharge across the exhausted space. The recent investigations of the circumstances of the electric dissociation produced in the atmosphere and in other gases by ultra-violet light, the Röntgen radiation, and other agencies, constitute one of the most striking developments in experimental molecular physics since Graham determined the molecular relations of gaseous diffusion and transpiration more than half a century ago. This advance in experimental knowledge of molecular phenomena, assisted by the discovery of the precise and rational effect of magnetism on the spectrum, has brought into prominence a modification or rather development of Maxwell's exposition of electric theory, which was dictated primarily by the requirements of the abstract theory itself; the atoms or ions are now definitely introduced as the carriers of those electric charges which interact across the æther, and so produce the electric fields whose transformations were the main subject of the original theory.

We are thus inevitably led, in electric and æthereal theory, as in the chemistry and dynamics of the gaseous state which is the department of abstract physics next in order of simplicity, to the consideration of the individual molecules of matter. The theoretical problems which

had come clearly into view a quarter of a century ago, under Maxwell's lead, whether in the exact dynamical relations of æthereal transmission or in the more fortuitous domain of the statistics of interacting molecules, are those around which attention is still mainly concentrated; but as the result of the progress in each, they are now tending towards consolidation into one subject. I propose—leaving further review of the scientific aspect of the recent enormous development of the applications of physical science for hands more competent to deal with the practical side of that subject—to offer some remarks on the scope and validity of this molecular order of ideas, to which the trend of physical explanation and development is now setting in so pronounced a manner.

If it is necessary to offer an apology for detaining the attention of the Section on so abstract a topic, I can plead its intrinsic philosophical importance. The hesitation so long felt on the Continent in regard to discarding the highly developed theories which analyzed all physical actions into direct attractions between the separate elements of the bodies concerned, in favor of a new method in which our ideas are carried into regions deeper than the phenomena, has now given place to eager discussion of the potentialities of the new standpoint. There has even appeared a disposition to consider that the Newtonian dynamical principles, which have formed the basis of physical explanation for nearly two centuries, must be replaced in these deeper subjects by a method of direct description of the mere course of phenomena, apart from any attempt to establish causal relations; the initiation of this method being traced, like that of the Newtonian dynamics itself, to this country. The question has arisen as to how far the new methods of æthereal physics are to be considered as an independent departure, how far they form

the natural development of existing dynamical science. In England, whence the innovation came, it is the more conservative position that has all along been occupied. Maxwell was himself trained in the school of physics established in this country by Sir George Stokes and Lord Kelvin, in which the dominating idea has been that of the strictly dynamical foundation of all physical action. Although the pupil's imagination bridged over dynamical chasms, across which the master was not always able to follow, yet the most striking feature of Maxwell's scheme was still the dynamical framework into which it was built. The more advanced reformers have now thrown overboard the apparatus of potential functions which Maxwell found necessary for the dynamical consolidation of his theory, retaining only the final result as a verified descriptive basis for the phenomena. In this way all difficulties relating to dynamical development and indeed consistency are avoided, but the question remains as to how much is thereby lost. In practical electromagnetics the transmission of power is now the most prominent phenomenon; if formal dynamics is put aside in the general theory, its guidance must here be replaced by some more empirical and tentative method of describing the course of transmission and transformation of mechanical energy in the system.

The direct recognition in some form, either explicitly or tacitly, of the part played by the æther has become indispensable to the development and exposition of general physics ever since the discoveries of Hertz left no further room for doubt that this physical scheme of Maxwell was not merely a brilliant speculation, but constituted, in spite of outstanding gaps and difficulties, a real formulation of the underlying unity in physical dynamics. The domain of abstract physics is in fact roughly divisible into two regions. In one of them we are

mainly concerned with interactions between one portion of matter and another portion occupying a different position in space; such interactions have very uniform and comparatively simple relations; and the reason is traceable to the simple and uniform constitution of the intervening medium in which they have their seat. The other province is that in which the distribution of the material molecules comes into account. Setting aside the ordinary dynamics of matter in bulk, which is founded on the uniformity of the properties of the bodies concerned and their experimental determination, we must assign to this region all phenomena which are concerned with the uncoordinated motions of the molecules, including the range of thermal and in part of radiant actions; the only possible basis for detailed theory is the statistical dynamics of the distribution of the molecules. The far more deep-seated and mysterious processes which are involved in changes in the constitution of the individual molecules themselves are mainly outside the province of physics, which is competent to reason only about permanent material systems; they must be left to the sciences of chemistry and physiology. Yet the chemist proclaims that he can determine only the results of his reactions and the physical conditions under which they occur; the character of the bonds which hold atoms in their chemical combinations is at present unknown, although a large domain of very precise knowledge relating, in some diagrammatic manner, to the topography of the more complex molecules has been attained. The vast structure which chemical science has in this way raised on the narrow foundation of the atomic theory is perhaps the most wonderful existing illustration both of the rationality of natural processes and of the analytical powers of the human mind. In a word, the complication of the material world is referable to

the vast range of structure and of states of aggregation in the material atoms; while the possibility of a science of physics is largely due to the simplicity of constitution of the universal medium through which the individual atoms interact on each other.

The reference of the uniformity in the interactions at a distance between material bodies to the part played by the æther is a step towards the elimination of extraneous and random hypotheses about laws of attraction between atoms. It also places that medium on a different basis from matter, in that its mode of activity is simple and regular, whereas intimate material interactions must be of illimitable complexity. This gives strong ground for the view that we should not be tempted towards explaining the simple group of relations which have been found to define the activity of the æther, by treating them as mechanical consequences of concealed structure in that medium; we should rather rest satisfied with having attained to their exact dynamical correlation, just as geometry explores or correlates, without explaining, the descriptive and metric properties of space. On the other hand, a view is upheld which considers the pressures and thrusts of the engineer, and the strains and stresses in the material structures by which he transmits them from one place to another, to be the archetype of the processes by which all mechanical effect is transmitted in nature. This doctrine implies an expectation that we may ultimately discover something analogous to structure in the celestial spaces, by means of which the transmission of physical effect will be brought into line with the transmission of mechanical effect by material frame work.

At a time when the only definitely ascertained function of the æther was the undulatory propagation of radiant energy across space, Lord Kelvin pointed out that, by reason of the very great velocity of prop-

agation, the density of the radiant energy in the medium at any place must be extremely small in comparison with the amount of energy that is transmitted in a second of time: this easily led him to the very striking conclusion that, on the hypothesis that the æther is like material elastic media, it is not necessary to assume its density to be more than 10^{-18} of that of water, or its optical rigidity to be more than ten 10^{-8} of that of steel or glass. Thus far the æther would be merely an impalpable material atmosphere for the transference of energy by radiation, at extremely small densities but with very great speed, while ordinary matter would be the seat of practically all this energy. But this way of explaining the absence of sensible influence of the æther on the phenomena of material dynamics lost much of its basis as soon as it was recognized that the same medium must be the receptacle of very high densities of energy in the electric fields around currents and magnets.* The other mode of explanation is to consider the æther to be of the very essence of all physical actions, and to correlate the absence of obvious mechanical evidence of its intervention with its regularity and universality.

On this plan of making the æther the essential factor is the transformation of energy as well as its transmission across space, the material atom must be some

*We can here only allude to Lord Kelvin's recent most interesting mechanical illustrations of a solid æther interacting with material molecules and with itself by attraction at a distance: unlike the generalized dynamical methods expounded in the text, which can leave the intimate structure of the material molecule outside the problem, a definite working constitution is there assigned to the molecular nucleus. It is pointed out in a continuation that is to appear in the *Philosophical Magazine* for September, that a density of æther of the order of only 10^{-9} , which would not appreciably affect the inertia of matter, would involve rigidity comparable with that of steel, and thus permit transmission of magnetic forces by stress; this solid æther is, however, as usual, taken to be freely permeable to the molecules of matter.

kind of permanent nucleus that retains around itself an æthereal field of physical influence, such as, for example, a field of strain. We can recognize the atom only through its interactions with other atoms that are so far away from it as to be practically independent systems; thus our direct knowledge of the atom will be confined to this field of force which belongs to it. Just as the exploration of the distant field of magnetic influence of a steel magnet, itself concealed from view, cannot tell us anything about the magnet except the amount and direction of its moment, so a practically complete knowledge of the field of physical influence of an atom might be expressible in terms of the numerical values of a limited number of physical moments associated with it, without any revelation as to its essential structure or constitution being involved. This will at any rate be the case for ultimate atoms if, as is most likely, the distances at which they are kept apart are large compared with the diameters of the atomic nuclei; it in fact forms our only chance for penetrating to definite dynamical views of molecular structure. So long as we cannot isolate a single molecule, but must deal observationally with an innumerable distribution of them, even this kind of knowledge will be largely confined to average values. But the last half-century has witnessed the successful application of a new instrument of research, which has removed in various directions the limitations that had previously been placed on the knowledge to which it was possible for human effort to look forward. The spectroscope has created a new astronomy by revealing the constitutions and the unseen internal motions of the stars. Its power lies in the fact that it does take hold of the internal relations of the individual molecule of matter, and provides a very definite and detailed, though far from complete, analysis of the vibratory motions

that are going on in it; these vibrations being in their normal state characteristic of its dynamical constitution, and in their deviations from the normal giving indications of the velocity of its movement and the physical state of its environment. Maxwell long ago laid emphasis on the fact that a physical atomic theory is not competent even to contemplate the vast mass of potentialities and correlations of the past and the future, that biological theory has to consider as latent in a single organic germ containing at most only a few million molecules. On our present view we can accept his position that the properties of such a body cannot be those of a 'purely material system,' provided, however, we restrict this phrase to apply to physical properties as here defined. But an exhaustive discovery of the intimate nature of the atom is beyond the scope of physics; questions as to whether it must not necessarily involve in itself some image of the complexity of the organic structures of which it can form a correlated part must remain a subject of speculation outside the domain of that science. It might be held that this conception of discrete atoms and continuous æther really stands, like those of space and time, in intimate relation with our modes of mental apprehension, into which any consistent picture of the external world must of necessity be fitted. In any case it would involve abandonment of all the successful traditions of our subject if we ceased to hold that our analysis can be formulated in a consistent and complete manner, so far as it goes, without being necessarily an exhaustive account of phenomena that are beyond our range of experiment. Such phenomena may be more closely defined as those connected with the processes of intimate combination of the molecules: they include the activities of organic beings which all seem to depend on change of molecular structure.

If, then, we have so small a hold on the intimate nature of matter, it will appear all the more striking that physicists have been able precisely to divine the mode of operation of the intangible æther, and to some extent explore in it the fields of physical influence of the molecules. On consideration we recognize that this knowledge of fundamental physical interaction has been reached by a comparative process. The mechanism of the propagation of light could never have been studied in the free æther of space alone. It was possible, however, to determine the way in which the characteristics of optical propagation are modified, but not wholly transformed, when it takes place in a transparent material body instead of empty space. The change in fact arises on account of the æther being entangled with the network of material molecules; but inasmuch as the length of a single wave of radiation covers thousands of these molecules the wave-motion still remains uniform and does not lose its general type. A wider variation of the experimental conditions has been provided for our examination in the case of those substances in which the phenomenon of double refraction pointed to a change of the æthereal properties which varied in different directions; and minute study of this modification has proved sufficient to guide to a consistent appreciation of the nature of this change, and therefore of the mode of æthereal propagation that is thus altered. In the same way, it was the study and development of the manner in which the laws of electric phenomena in material bodies had been unraveled by Ampère and Faraday, that guided Faraday himself and Maxwell—who were impressed with the view that the æther was at the bottom of it all—in their progress towards an application of similar laws to æther devoid of matter, such as would complete a scheme of continuous action by consist-

ently interconnecting the material bodies and banishing all untraced interaction across empty space. Maxwell in fact chose to finally expound the theory by ascribing to the æther of free space a dielectric constant and a magnetic constant of the same type as had been found to express the properties of material media, thus extending the seat of the phenomena to all space on the plan of describing the activity of the æther in terms of the ordinary electric ideas. The converse mode of development, starting with the free æther under the directly dynamical form which has been usual in physical optics, and introducing the influence of the material atoms through the electric charges which are involved in their constitution,* was hardly employed by him; in part, perhaps, because, owing to the necessity of correlating his theory with existing electric knowledge and the mode of its expression, he seems never to have reached the stage of moulding it into a completely deductive form.

The dynamics of the æther, in fact the recognition of the existence of an æther, has thus, as a matter of history, been reached through study of the dynamical phenomena of matter. When the dynamics of a material system is worked up to its purest and most general form, it becomes a formulation of the relations between the succession of the configurations and states of motion of the system, the assistance of an independent idea of force not being usually required. We can, however, only attain such a compact statement when the system is self-contained, when its motion is not being dissipated by agencies of fric-

tional type, and when its connections can be directly specified by purely geometrical relations between the co-ordinates, thus excluding such mechanisms as rolling contacts. The course of the system is then in all cases determined by some form or other of a single fundamental property, that any alteration in any small portion of its actual course must produce an increase in the total 'Action' of the motion. It is to be observed that in employing this law of minimum as regards the Action expressed as an integral over the whole time of the motion, we no more introduce the future course as a determining influence on the present state of motion than we do in drawing a straight line from any point in any direction, although the length of the line is the minimum distance between its ends. In drawing the line piece by piece we have to make tentative excursions into the immediate future in order to adjust each element into straightness with the previous element; so in tracing the next stage of the motion of a material system we have similarly to secure that it is not given any such directions as would unduly increase the Action. But whatever views may be held as to the ultimate significance of this principle of action, its importance, not only for mathematical analysis, but as a guide to physical exploration, remains fundamental. When the principles of the dynamics of material systems are refined down to their ultimate common basis, this principle of minimum is what remains. Hertz preferred to express its contents in the form of a principle of straightness of course or path. It will be recognized, on the lines already indicated, that this is another mode of statement of the same fundamental idea; and the general equivalence is worked out by Hertz on the basis of Hamilton's development of the principles of dynamics. The latter mode of statement may be adaptable so as to avoid

* In 1870 Maxwell, while admiring the breadth of the theory of Weber, which is virtually based on atomic charges combined with action at a distance, still regarded it as irreconcilable with his own theory, and left to the future the question as to why 'theories apparently so fundamentally opposed should have so large a field of truth common to both.'—*Scientific Papers*, II., p. 228.

the limitations which restrict the connections of the system, at the expense, however, of introducing new variables; if, indeed, it does not introduce gratuitous complexity for purposes of physics to attempt to do this. However these questions may stand, this principle of straightness or directness of path forms, whenever it applies, the most general and comprehensive formulation of purely dynamical action: it involves in itself the complete course of events. In so far as we are given the algebraic formula for the time-integral which constitutes the Action, expressed in terms of any suitable coordinates, we know implicitly the whole dynamical constitution and history of the system to which it applies. Two systems in which the Action is expressed by the same formula are mathematically identical, are physically precisely correlated, so that they have all dynamical properties in common. When the structure of a dynamical system is largely concealed from view, the safest and most direct way towards an exploration of its essential relations and connections, and in fact towards answering the prior question as to whether it is a purely dynamical system at all, is through this order of ideas. The ultimate test that a system is a dynamical one is not that we shall be able to trace mechanical stresses throughout it, but that its relations can be in some way or other consolidated into accordance with this principle of minimum Action. This definition of a dynamical system in terms of the simple principle of directness of path may conceivably be subject to objection as too wide; it is certainly not too narrow; and it is the conception which has naturally been evolved from two centuries of study of the dynamics of material bodies. Its very great generality may lead to the objection that we might completely formulate the future course of a system in its terms, without having obtained a working famili-

arity with its details of the kind to which we have become accustomed in the analysis of simple material systems; but our choice is at present between this kind of formulation, which is a real and essential one, and an empirical description of the course of phenomena combined with explanations relating to more or less isolated groups. The list of great names, including Kelvin, Maxwell, Helmholtz, that have been associated with the employment of the principle for the elucidation of the relations of deep-seated dynamical phenomena, is a strong guarantee that we shall do well by making the most of this clue.

Are we then justified in treating the material molecule, so far as revealed by the spectroscope, as a dynamical system coming under this specification? Its intrinsic energy is certainly permanent and not subject to dissipation; otherwise the molecule would gradually fade out of existence. The extreme precision and regularity of detail in the spectrum shows that the vibrations which produce it are exactly synchronous whatever be their amplitude, and in so far resemble the vibrations of small amplitude in material systems. As all indications point to the molecule being a system in a state of intrinsic motion, like a vortex ring, or a stellar system in astronomy, we must consider these radiating vibrations to take place around a steady state of motion which does not itself radiate, not around a state of rest. Now not the least of the advantages possessed by the Action principle, as a foundation for theoretical physics, is the fact that its statement can be adapted to systems involving in their constitution permanent steady motions of this kind, in such a way that only the variable motions superposed on them come into consideration. The possibilities as regards physical correlation of thus introducing permanent motional states as well as permanent structure into the constitution of

our dynamical systems have long been emphasized by Lord Kelvin;* the effective adaptation of abstract dynamics to such systems was made independently by Kelvin and Routh about 1877; the more recent exposition of the theory by Helmholtz has directed general attention to what is undoubtedly the most significant extension of dynamical analysis which has taken place since the time of Lagrange.

Returning to the molecules, it is now verified that the Action principle forms a valid foundation throughout electrodynamics and optics; the introduction of the æther into the system has not affected its application. It is therefore a reasonable hypothesis that the principle forms an allowable foundation for the dynamical analysis of the radiant vibrations in the system formed by a single molecule and surrounding æther; and the knowledge which is now accumulating, both of the orderly grouping of the lines of the spectrum and of the modifications impressed on these lines by a magnetic field or by the density of the matter immediately surrounding the vibrating molecule, can hardly fail to be fruitful for the dynamical analysis of its constitution. But let it be repeated that this analysis would be complete when a formula for the dynamical energy of the molecule is obtained, and would go no deeper. Starting from our definitely limited definition of the nature of a dynamical system, the problem is merely to correlate the observed relations of the periods of vibration in a molecule, when it has come into a steady state as regards constitution and is not under the influence of intimate encounter with other molecules.

It may be recalled incidentally that the generalized Maxwell-Boltzmann principle

* For a classical exposition see his *Brit. Assoc. Address* of 1884 on 'Steps towards a Kinetic Theory of Matter,' reprinted in 'Popular Lectures and Addresses,' vol i.

of the equable distribution of the acquired store of kinetic energy of the molecule, among its various possible independent types of motion, is based directly on the validity of the Action principle for its dynamics. In the demonstrations usually offered the molecule is considered to have no permanent or constitutive energy of internal motion. It can, however, be shown, by use of the generalization aforesaid of the Action principle, that no discrepancy will arise on that account. Such intrinsic kinetic energy virtually adds on to the potential energy of the system; and the remaining or acquired part of the kinetic energy of the molecule may be made the subject of the same train of reasoning as before.

Let us now return to the general question whether our definition of a dynamical system may not be too wide. As a case in point, the single principle of Action has been shown to provide a definite and sufficient basis for electrodynamics; yet when, for example, one armature of an electric motor pulls the other after it without material contact, and so transmits mechanical power, no connection between them is indicated by the principle such as could by virtue of internal stress transmit the pull. The essential feature of the transmission of a pull by stress across a medium is that each element of volume of the medium acts by itself, independently of the other elements. The stress excited in any element depends on the strain or other displacement occurring in that element alone; and the mechanical effect that is transmitted is considered as an extraneous force applied at one place in the medium, and passed on from element to element through these internal pressures and tractions until it reaches another place. We have, however, to consider two atomic electric charges as being themselves some kind of strain configurations in the æther; each of them already involves an atmosphere of

strain in the surrounding æther which is part of its essence, and cannot be considered apart from it; each of them essentially pervades the entire space, though on account of its invariable character we consider it as a unit. Thus we appear to be debarred from imagining the æther to act as an elastic connection which is merely the agent of transmission of a pull from the one nucleus to the other, because there are already stresses belonging to and constituting an intrinsic part of the terminal electrons, which are distributed all along the medium. Our action criterion of a dynamical system, in fact, allows us to reason about an electron as a single thing, notwithstanding that its field of energy is spread over the whole medium; it is only in material solid bodies, and in problems in which the actual sphere of physical action of the molecule is small compared with the smallest element of volume that our analysis considers, that the familiar idea of transmission of force by simple stress can apply. Whatever view may ultimately command itself, this question is one that urgently demands decision. A very large amount of effort has been expended by Maxwell, Helmholtz, Heaviside, Hertz and other authorities in the attempt to express the mechanical phenomena of electrical action in terms of a transmitting stress. The analytical results up to a certain point have been promising, most strikingly so at the beginning, when Maxwell established the mathematical validity of the way in which Faraday was accustomed to represent to himself the mechanical interactions across space, in terms of a tension along the lines of force equilibrated by an equal pressure preventing their expansion sideways. According to the views here developed, that ideal is an impossible one; if this could be established to general satisfaction the field of theoretical discussion would be much simplified.

This view that the atom of matter is, so far as regards physical actions, of the nature of a structure in the æther involving an atmosphere of æthereal strain all around it, not a small body which exerts direct actions at a distance on other atoms according to extraneous laws of force, was practically foreign to the eighteenth century, when mathematical physics was modelled on the Newtonian astronomy and dominated by its splendid success. The scheme of material dynamics, as finally compactly systematized by Lagrange, had therefore no direct relation to such a view, although it has proved wide enough to include it. The remark has often been made that it is probably owing to Faraday's mathematical instinct, combined with his want of acquaintance with the existing analysis, that the modern theory of the æther obtained a start from the electric side. Through his teaching and the weight of his authority, the notion of two electric currents exerting their mutual forces by means of an intervening medium, instead of by direct attraction across space, was at an early period firmly grasped in this country. In 1845 Lord Kelvin was already mathematically formulating, with most suggestive success, continuous elastic connections, by whose strain the fields of activity of electric currents or of electric distributions could be illustrated; while the exposition of Maxwell's interconnected scheme, in the earlier form in which it relied on concrete models of the electric action, goes back almost to 1860. Corresponding to the two physical ideals of isolated atoms exerting attraction at a distance, and atoms operating by atmospheres of æthereal strain, there are, as already indicated, two different developments of dynamical theory. The original Newtonian equations of motion determined the course of a system by expressing the rates at which the velocity of each of its small parts or elements is changing. This

method is still fully applicable to those problems of gravitational astronomy in which dynamical explanation was first successful on a grand scale, the planets being treated as point-masses, each subject to the gravitational attraction of the other bodies. But the more recent development of the dynamics of complex systems depends on the fact that analysis has been able to reduce within manageable limits the number of varying quantities whose course is to be explicitly traced, through taking advantage of those internal relations of the parts of the system that are invariable, either geometrically or dynamically. Thus, to take the simplest case, the dynamics of a solid body can be confined to a discussion of its three components of translation and its three components of rotation, instead of the motion of each element of its mass. With the number of independent co-ordinates thus diminished when the initial state of the motion is specified the subsequent course of the complete system can be traced; but the course of the changes in any part of it can only be treated in relation to the motion of the system as a whole. It is just this mode of treatment of a system as a whole that is the main characteristic of modern physical analysis. The way in which Maxwell analyzed the interactions of a system of linear electric currents, previously treated as if each were made up of small independent pieces or elements, and accumulated the evidence that they formed a single dynamical system, is a trenchant example. The interactions of vortices in fluid form a very similar problem, which is of special note in that the constitution of the system is there completely known in advance, so that the two modes of dynamical exposition can be compared. In this case the older method forms independent equations for the motion of each material element of the fluid, and so requires the introduction of the

stress—here the fluid pressure—by which dynamical effect is passed on to it from the surrounding elements: it corresponds to a method of contact action. But Helmholtz opened up new ground in the abstract dynamics of continuous media when he recognized (after Stokes) that, if the distribution of the velocity of spin at those places in the fluid where the motion is vortical be assigned, the motion in every part of the fluid is therein kinematically involved. This, combined with the theorem of Lagrange and Cauchy, that the spin is always confined to the same portions of the fluid, formed a starting-point for his theory of vortices, which showed how the subsequent course of the motion can be ascertained without consideration of pressure or other stress.

The recognition of the permanent state of motion constituting a vortex ring as a determining agent as regards the future course of the system was in fact justly considered by Helmholtz as one of his greatest achievements. The principle had entirely eluded the attention of Lagrange and Cauchy and Stokes, who were the pioneers in this fundamental branch of dynamics, and had virtually prepared all the necessary analytical material for Helmholtz's use. The main import of this advance lay, not in the assistance which is afforded to the development of the complete solution of special problems in fluid motion, but in the fact that it constituted the discovery of the types of permanent motion of the system, which could combine and interact with each other without losing their individuality,* though each of them pervaded the whole field. This rendered possible an entirely new mode of treatment; and mathematicians who were accustomed, as in astronomy, to aim directly at the determina-

* We may compare G. W. Hill's more recent introduction of the idea of permanent orbits into physical astronomy.

tion of all the details of the special case of motion, were occasionally slow to apprehend the advantages of a procedure which stopped at formulating a description of the nature of the interaction between various typical groups of motions into which the whole disturbance could be resolved.

The new train of ideas introduced into physics by Faraday was thus consolidated and emphasized by Helmholtz's investigations of 1858 in the special domain of hydrodynamics. In illustration let us consider the fluid medium to be pervaded by permanent vortices circulating round solid rings as cores; the older method of analysis would form equations of motion for each element of the fluid, involving the fluid pressure, and by their integration would determine the distribution of pressure on each solid ring, and thence the way it moves. This method is hardly feasible even in the simplest cases. The natural plan is to make use of existing simplifications by regarding each vortex as a permanent reality, and directly attacking the problem of its interactions with the other vortices. The energy of the fluid arising from the vortex motion can be expressed in terms of the positions and strengths of the vortices alone; and then the principle of Action, in the generalized form which includes steady motional configurations as well as constant material configurations, affords a method of deducing the motions of the cores and the interactions between them. If the cores are thin they in fact interact mechanically, as Lord Kelvin and Kirchhoff proved, in the same manner as linear electric currents would do; though the impulse thence derived towards a direct hydro-kinetic explanation of electro-magnetics was damped by the fact that repulsion and attraction have to be interchanged in the analogy. The conception of vortices, once it has been arrived at, forms the natural physical basis of investigation, al-

though the older method of determining a distribution of pressure-stress throughout the fluid and examining how it affects the cores is still possible; that stress, however, is not simply transmitted, as it has to maintain the changes of velocity of the various portions of the fluid. But if the vortices have no solid cores we are at a loss to know where even this pressure can be considered as applied to them; if we follow up the stress, we lose the vortex; yet a fluid vortex can nevertheless illustrate an atom of matter, and we can consider such atoms as exerting mutual forces, only these forces cannot be considered as transmitted through the agency of fluid pressure. The reason is that the vortex cannot now be identified with a mere core bounded by a definite surface, but is essentially a configuration of motion extending throughout the medium.

Thus we are again in face of the fundamental question whether all attempts to represent the mechanical interactions of electro-dynamic systems, as transmitted from point to point by means of simple stress, are not doomed to failure; whether they do not, in fact, introduce unnecessary and insurmountable difficulty into the theory. The idea of identifying an atom with a state of strain or motion, pervading the region of the æther around its nucleus, appears to demand wider views as to what constitutes dynamical transmission. The idea that any small portion of the primordial medium can be isolated, by merely introducing tractions acting over its surface and transmitted from the surrounding parts, is no longer appropriate or consistent; a part of the dynamical disturbance in that element of the medium is on this hypothesis already classified as belonging to, and carried along with, atoms that are outside it but in its neighborhood—and this part must not be counted twice over. The law of Poynting relating to the paths of the

transmission of energy is known to hold in its simple form only when the electric charges or currents are in a steady state; when they are changing their positions or configurations their own fields of intrinsic energy are carried along with them.

It is not surprising, considering the previous British familiarity with this order of ideas, that the significance for general physics of Helmholtz's doctrine of vortices was eagerly developed in this country, in the form in which it became embodied through Lord Kelvin's famous illustration of the constitution of the matter, as consisting of atoms with separate existence and mutual interactions. This vortex atom theory has been a main source of physical suggestion because it presents, on a simple basis, a dynamical picture of an ideal material system, atomically constituted, which could go on automatically without extraneous support. The value of such a picture may be held to lie, not in any supposition that this is the mechanism of the actual world laid bare, but in the vivid illustration it affords of the fundamental postulate of physical science, that mechanical phenomena are not parts of a scheme too involved for us to explore, but rather present themselves in definite and consistent correlations, which we are able to disentangle and apprehend with continuously increasing precision.

It would be an interesting question to trace the origin of our preference for a theory of transmission of physical action over one of direct action at a distance. It may be held that it rests on the same order of ideas as supplies our conception of force; that the notion of effort which we associate with change of the motion of a body involves the idea of a mechanical connection through which that effort is applied. The mere idea of a transmitting medium would then be no more an ultimate foundation for physical explanation than that of force itself. Our choice between direct distance

action and mediate transmission would thus be dictated by the relative simplicity and coherence of the accounts they give of the phenomena: this is, in fact, the basis on which Maxwell's theory had to be judged until Hertz detected the actual working of the medium. Instantaneous transmission is to all intents action at a distance, except in so far as the law of action may be more easily formulated in terms of the medium than in a direct geometrical statement.

In connection with these questions it may be permitted to refer to the eloquent and weighty address recently delivered by M. Poincaré to the International Congress of Physics. M. Poincaré accepts the principle of Least Action as a reliable basis for the formulation of physical theory, but he imposes the condition that the results must satisfy the Newtonian law of equality of action and reaction between each pair of bodies concerned, considered by themselves; this, however, he would allow to be satisfied indirectly, if the effects could be traced across the intervening æther by stress, so that the tractions on the two sides of each ideal interface are equal and opposite.* As above argued, this view appears to exclude *ab initio* all atomic theories of the general type of vortex atoms, in which the energy of the atom is distributed throughout the medium instead of being concentrated in a nucleus; and this remark seems to go to the root of the question. On the other hand, the position here asserted is that recent dynamical developments have permitted the extension of the principle of Action to systems involving permanent motions, whether obvious or latent, as part of their constitution; that on this wider basis the

* Cf. also Hertz on the electro-magnetic equations, § 12, *Wied. Ann.*, 1890. The problem of merely replacing a system of forces by a statical stress is widely indeterminate, and therefore by itself unreal; the actual question is whether any such representation can be coordinated with existing dynamics.

atom may itself involve a state of steady disturbance extending through the medium, instead of being only a local structure acting by push and pull. The possibilities of dynamical explanation are thus enlarged. The most definite type of model yet imagined of the physical interaction of atoms through the æther is, perhaps, that which takes the æther to be a rotationally elastic medium after the manner of MacCullagh and Rankine, and makes the ultimate atom include the nucleus of a permanent rotational strain-configuration, which as a whole may be called an electron. The question how far this is a legitimate and effective model stands by itself, apart from the dynamics which it illustrates; like all representations it can only cover a limited ground. For instance, it cannot claim to include the internal structure of the nucleus of an atom or even of an electron; for purposes of physical theory that problem can be put aside, it may even be treated as inscrutable. All that is needed is a postulate of free mobility of this nucleus through the æther. This is definitely hypothetical, but it is not an unreasonable postulate because a rotational æther has the properties of a perfect fluid medium except where differentially rotational motions are concerned, and so would not react on the motion of any structure moving through it except after the manner of an apparent change of inertia. It thus seems possible to hold that such a model forms an allowable representation of the dynamical activity of the æther, as distinguished from the complete constitution of the material nuclei between which that medium establishes connection.

At any rate, models of this nature have certainly been most helpful in Maxwell's hands toward the effective intuitive grasp of a scheme of relations as a whole, which might have proved too complex for abstract unravelment in detail. When a physical model of concealed dynamical processes has

served this kind of purpose, when its content has been explored and estimated, and has become familiar through the introduction of new terms and ideas, then the ladder by which we have ascended may be kicked away, and the scheme of relations which the model embodied can stand forth in severely abstract form. Indeed many of the most fruitful branches of abstract mathematical analysis itself have owed their start in this way to concrete physical conceptions. This gradual transition into abstract statement of physical relations in fact amounts to retaining the essentials of our working models while eliminating the accidental elements involved in them; elements of the latter kind must always be present because otherwise the model would be identical with the thing which it represents, whereas we cannot expect to mentally grasp all aspects of the content of even the simplest phenomena. Yet the abstract standpoint is always attained through the concrete; and for purposes of instruction such models, properly guarded, do not perhaps ever lose their value; they are just as legitimate aids as geometrical diagrams, and they have the same kind of limitations. In Maxwell's words, 'for the sake of persons of these different types scientific truth should be presented in different forms, and should be regarded as equally scientific whether it appear in the robust form and the vivid coloring of a physical illustration, or in the tenuity and paleness of a symbolical expression.' The other side of the picture, the necessary incompleteness of even our legitimate images and modes of representation, comes out in the despairing opinion of Young ('*Chromatics*,' 1817), at a time when his faith in the undulatory theory of light had been eclipsed by Malus's discovery of the phenomena of polarization by reflection, that this difficulty 'will probably long remain, to mortify the van-

ity of an ambitious philosophy, completely unresolved by any theory': not many years afterwards the mystery was solved by Fresnel.

This process of removing the intellectual scaffolding by which our knowledge is reached, and preserving only the final formulæ which express the correlations of the directly observable things, may moreover readily be pushed too far. It asserts the conception that the universe is like an enclosed clock that it wound up to go, and that accordingly we can observe that it is going, and can see some of its more superficial movements, but not much of them; that thus, by patient observation and use of analogy, we can compile, in merely tabular form, information as to the manner in which it works and is likely to go on working, at any rate for some time to come; but that any attempt to probe the underlying connection is illusory or illegitimate. As a theoretical precept this is admirable. It minimizes the danger of our ignoring or forgetting the limitations of human faculty, which can only utilize the imperfect representations that the external world impresses on our senses. On the other hand such a reminder has rarely been required by the master minds of modern science, from Descartes and Newton onwards, whatever their theories may have been. Its danger as a dogma lies in its application. Who is to decide without risk of error, what is essential fact and what is intellectual scaffolding? To which class does the atomic theory of matter belong? That is, indeed, one of the intangible things which it is suggested may be thrown overboard, in sorting out and classifying our scientific possessions. Is the mental idea or image, which suggests, and alone can suggest, the experiment that adds to our concrete knowledge, less real than the bare phenomenal uniformity which it has revealed? Is it not, perhaps, more real in

that the uniformities might not have been there in the absence of the mind to perceive them?

No time is now left for review of the methods of molecular dynamics. Here our knowledge is entirely confined to steady states of the molecular system: it is purely statical. In ordinary statics and the dynamics of undisturbed steady motions, the form of the energy function is the sufficient basis of the whole subject. This method is extended to thermo-dynamics by making use of the mechanically available energy of Rankine and Kelvin, which is a function of the bodily configuration and chemical constitution and temperature of the system, whose value cannot under any circumstances spontaneously increase, while it will diminish in any operation which is not reversible. In the statics of systems in equilibrium or in steady motion, this method of energy is a particular case of the method of Action; but in its extension to thermal statics it is made to include chemical as well as configurational changes, and a new point appears to arise. Whether we do or do not take it to be possible to trace the application of the principle of Action throughout the process of chemical combination of two molecules, we certainly here postulate that the static case of that principle, which applies to steady systems, can be extended across chemical combinations. The question is suggested whether extension would also be valid to transformations which involve vital processes. This seems to be still considered an open question by the best authorities. If it be decided in the negative a distinction is involved between vital and merely chemical processes.

It is now taken as established that vital activity cannot create energy, at any rate in the long run which is all that can from the nature of the case be tested. It seems not unreasonable to follow the anal-

ogy of chemical actions, and assert that it cannot in the long run increase the mechanical availability of energy—that is, considering the organism as an apparatus for transforming energy without being itself in the long run changed. But we cannot establish a Carnot cycle for a portion of an organism, nor can we do so for a limited period of time; there might be creation of availability accompanied by changes in the organism itself, but compensated by destruction and the inverse changes a long time afterwards. This amounts to asserting that where, as in a vital system or even in a simple molecular combination, we are unable to trace or even assert complete dynamical sequence, exact thermodynamic statements should be mainly confined to the activity of the existing organism as a whole; it may transform inorganic material without change of energy and without gain of availability, although any such statements would be inappropriate and unmeaning as regards the details of the processes that take place inside the organism itself.

In any case it would appear that there is small chance of reducing these questions to direct dynamics; we should rather regard Carnot's principle, which includes the law of uniformity of temperature and is the basis of the whole theory, as a property of statistical type confined to stable or permanent aggregations of matter. Thus no dynamical proof from molecular considerations could be regarded as valid unless it explicitly restricted the argument to permanent systems; yet the conditions of permanency are unknown except in the simpler cases. The only mode of discussion that is yet possible is the method of dynamical statistics of molecules introduced by Maxwell. Now statistics is a method of arrangement rather than of demonstration. Every statistical argument requires to be verified by comparison with

the facts, because it is of the essence of this method to take things as fortuitously distributed except in so far as we know the contrary; and we simply may not know essential facts to the contrary. For example, if the interaction of the æther or other cause produces no influence to the contrary, the presumption would be that the kinetic energy acquired by a molecule is, on the average, equally distributed among its various independent modes of motion, whether vibrational or translational. Assuming this type of distribution to be once established in a gaseous system, the dynamics of Boltzmann and Maxwell show that it must be permanent. But its assumption in the first instance is a result rather of the absence than of the presence of knowledge of the circumstances, and can be accepted only so far as it agrees with the facts; our knowledge of the facts of specific heat shows that it must be restricted to modes of motion that are homologous. In the words of Maxwell, when he first discovered in 1860, to his great surprise, that in a system of colliding rigid atoms the energy would always be equally divided between translatory and rotatory motion, it is only necessary to assume, in order to evade this unwelcome conclusion, that 'something essential to the complete statement of the physical theory of molecular encounters must have hitherto escaped us.'

Our survey thus tends to the result, that as regards the simple and uniform phenomena which involve activity of finite regions of the universal æther, theoretical physics can lay claim to constructive functions, and can build up a definite scheme; but in the domain of matter the most that it can do is to accept the existence of such permanent molecular systems as present themselves to our notice, and fit together an outline plan of the more general and universal features in their activity. Our

well-founded belief in the rationality of natural processes asserts the possibility of this, while admitting that the intimate details of atomic constitution are beyond our scrutiny and provide plenty of room for processes that transcend finite dynamical correlation.

JOSEPH LARMOR.

INLAND BIOLOGICAL LABORATORIES.

THE following informal notes have been received concerning the season's work in various summer laboratories and experiment stations:

Of the research work carried out on the Great Lakes under the auspices of the United States Fish Commission, Professor Reighard says: The work has been purely research work and it was understood from the start that it should be of a fundamental scientific character rather than directed toward the immediate solution of questions of supposed practical importance.

The funds available have not permitted of carrying on the work for more than two months of each summer. During the summers of 1898 and 1899 it was carried on chiefly at Put in Bay, Ohio, (an island in the western end of Lake Erie, at which there is a hatchery of the Commission). By removing the internal fittings of the hatchery it was temporarily converted into a laboratory for each summer's use. This laboratory has been in every way amply equipped. There is gas and water, a small steamer and a supply of other boats. It is intended that work should begin on the first of July, but owing to delay in appropriation bills and to other causes it may happen, as it did this year, that no authorization for the commencement of the work can be issued until the end of June or even the early part of July. Supplies must then be ordered, arrangements made with workers and the hatchery converted into a laboratory. The difficulty involved in under-

taking to do this after the first of July for work which is to continue only two months, led this year to the trial of a different plan. Instead of opening the Put in Bay laboratory an effort is being made to carry on the work by means of individual investigators or small parties working independently. It is hoped that work carried on in this way can be continued over a longer period, even during a part of the college year.

The investigations carried on at the laboratory (and elsewhere during the present summer) are as follows:

BOTANICAL WORK.

1. *The Algæ of Lake Erie*.—Dr. Julia W. Snow has been engaged during each of the three seasons and is now engaged in the determination of the algæ of the Lake and in working out their life histories by means of cultures. As many of them assume different forms under different conditions, it is necessary to cultivate them and no final identifications are possible until the life history of each is known. This is of course a labor of years and involves a consideration of the relation of the various algæ groups to the nutritive substances contained in the water, that is, it leads into bio-chemistry. It is expected that results already obtained will be made ready for publication during the coming year.

2. *The larger Aquatic Plants*.—During the first season Mr. A. J. Pieters of the Department of Agriculture at Washington undertook a study of the larger aquatic plants with the purpose of determining whether they are wholly dependent on the water for nutrition or partly on the soil. Mr. Pieters' results are now in press. He did not get much further than a determination of the various soils present on the Lake bottom and the relation of the plants to them. During the second season and during the present season Mr. R. H. Pond, an assistant in Botany at the University, has car-