

the recognized rules of priority. When for any group such diagnoses of the different subdivisions shall have been published, and, after discussion, so modified as to be acceptable to the majority of students of the group, forms subsequently described should be accompanied by similar diagnoses and similar designations of a type which will render them strictly comparable to forms already known.

The Hydracarina form a sharply limited and very homogeneous group in which the application of such a scheme as proposed above seems practicable. Accordingly, it is suggested here, and in the complete paper it is expected that there will be given for each family and genus: first, the name having priority; second the author of the same, together with the date and exact reference; third, a diagnosis in Latin and English; fourth, the type, with reference to the author and exact date, together with the reasons for selection of the same.

*Southeastern United States as a Center of Geographical Distribution of Fauna and Flora:* CHARLES C. ADAMS. (Read by title only.)

In general the geographical relationship of the fauna and flora of the northern United States, east of the Great Plains, is with that of the Southeast, and points to an origin in that direction, except in the case of the distinctly boreal forms. The abundance and diversity of life in the Southeast indicate that it has been, and now is, a center of dispersal. The relicts indicate that it has been a center of preservation of ancient types, and the endemism shows that it has been a center of origin of types. There are two distinct southern centers of dispersal in temperate United States, one in the moist Southeast and the other in the arid Southwest. Nine criteria, aside from fossil evidence, are recognized for determining the

center of origin or the locality of dispersal: (1) Location of the greatest differentiation of a type; (2) location of dominance or great abundance of individuals; (3) location of synthetic or closely related forms; (4) location of maximum size of individuals; (5) location of greatest productiveness and its stability, in crops; (6) continuity and convergence of lines of dispersal; (7) location of least dependence upon a restricted habitat; (8) continuity and directness of individual variations or modifications radiating from the center of origin along the highways of dispersal; (9) direction indicated by biogeographical affinities and (10) annual migration routes in birds. There are three primary outlets of dispersal from the Southeast: (1) The Mississippi Valley and its tributaries; (2) the Coastal Plain, and (3) the Appalachian Mountains and adjacent plateaus. The first two have also functioned for tropical types and the third for boreal forms. Dispersal is both forward and backward along these highways. It is desirable to study individual variation of animals and plants along their lines of dispersal and divergence from the center of origin, in such characters as size, productiveness, continuity of variation, color variation, and changes of habit and habitats. Life areas should be studied as centers of dispersal and origin and hence dynamically and genetically.

*Description of Cephalogonimus vesicaudus, sp. nov.:* W. S. NICKERSON. (Read by title only.)

*Fresh Water Polychæta:* H. P. JOHNSON.

*The Lateral Line System of Polyodon spathula:* HENRY F. NACHTRIEB.

The paper considered only the general anatomical features of the lateral line of *Polyodon*. In general the systems of the

two sides are symmetrical, but in the position, distribution, number and minor details of the branches from the 'lateral' and 'main' canals there is considerable variation. In none of the several hundred specimens examined were these branches grouped as described and figured by Collinge. In all cases they were found all along the 'lateral' canal, the great majority being ventral to the canal. As a rule, one to three at the anterior end of each system begin on the ventral side of the canal and, after running a short distance in that direction, turn dorsalward and terminate in the usual branchlets and clusters of sense organs on the dorsal side of the canal.

The points made were demonstrated with dried skins of the fish upon which the systems had been painted over with white paint, and photographs of similarly prepared skins.

*Conditions of Fossilization:* J. CULVER HARTZELL. (Read by title only.)

Professor Hartzell's paper was a review of a series of investigations he has been making, the objective point of which is to find the laws (?) governing the conditions of fossilization for the various classes of Invertebrates in the same and in different formations.

Before the laws desired can be formulated, it is necessary (a) to know the mineral composition of the skeletal parts of living invertebrates; (b) to know the condition of the fossil, *i. e.*, whether it be the original, a mold or a cast; (c) to know the mineral composition of the fossil; (d) to know what mineral change has taken place during fossilization where the cast is one by molecular replacement; (e) to know the lithological composition of the formations in which fossils occur; (f) to know the relationship between the fossil and the formation.

The conversion of an organism into a fossil depends upon the character of its skeletal parts, the material in which it is buried and the material brought in in solution by infiltration. The material of which the skeletal parts is composed varies in different groups, being more durable in some than in others and therefore plays an important part in the preservation of the organism. The variation in the lithological character of the material in which the organism is buried also plays an important part. Certain organisms are preserved as originals, others as molds and casts, in the same formation, and locality. In this same formation, but in a locality of different lithological character, those groups which were lost under the former condition may be retained under the latter, and *vice versa*.

So far, twenty-five horizons and forty-four localities in the United States, Canada, England and Germany have been examined with special reference to the lithological character of each formation at the various localities and the conditions of preservation of the fossils. Tables have been prepared giving the general mineral character of the skeletal parts of living invertebrates, minerals replacing original minerals secreted by the organisms, and a comparative table showing the mineral composition of living and fossil invertebrates.

The paper was illustrated by means of photographs, drawings and models.

*Origin and Migration of the Germ-Cells in Squalus Acanthias:* FREDERICK ADAMS WOODS.

The germ-cells in this form are not derived from the germinal epithelium of the body cavity, as is commonly taught, but are traceable before the mesoderm has split to form the coelom, and in a region that may be called extra-embryonic.