

lating them they were placed in sterile glass chambers having close-fitting covers. In six days the apples showed signs of rotting and in ten days pycnidia had begun to form. On the fourteenth day after inoculation the entire epidermis was blackened and densely dotted with the protruding pycnidia. Here again no difference was observed either in the manner of growth or the decay produced by the two species of *Sphæroopsis*. An examination of the inoculated apples showed that they were entirely free from other fungi. Apples treated in the same way and put under similar conditions but stabbed with a sterile scalpel did not decay.

Finally, spores obtained from the inoculated apples were used to inoculate healthy branches of both the apple and the sumac. So far no difference can be observed in the growth of *Sphæroopsis rhoïna* and *Sphæroopsis malorum* on the apple tree, but the fact that growth has gone on from the points of inoculation is established. In the sumac, growth has not been so rapid.

The facts already established in these experiments go to show that *Sphæroopsis rhoïna* will cause black-rot in the fruit of the apple and will also produce the typical 'canker' on the branches and limbs just as readily as *Sphæroopsis malorum*. Although the evidence is not yet complete it is probable that the two species are identical.

P. J. O'GARA.

THE UNIVERSITY OF NEBRASKA,
August 27, 1902.

PALEONTOLOGICAL NOTES.

THE GENERIC NAME OMOSAURUS.

THE name *Omosaurus armatus* was applied by Owen in 1874 to a dinosaur from the Kimmeridge Clay described by him in 'A Monograph on the Fossil Reptilia of the Mesozoic Formations,' issued by the Paleontographical Society. The name first occurs on page 46 of the part printed in 1875.

The same generic name had, however, been used by Leidy in 1856 for a crocodilian described by him on page 256 of the *Proceedings of the Academy of Natural Sciences*, of Phila-

delphia, for that year, to which he gave the name *Omosaurus perplexus*.

Omosaurus Owen is thus preoccupied, and for the genus of Stegosauruses included under that name I propose the name *Dacentrurus* in allusion to the powerful spines with which the tail was armed.

A NEW GENERIC NAME FOR STEGOSAURUS MARSHI.

IN Vol. XXIII. of the *Proceedings of the U. S. National Museum*, pp. 591, 592, I described a new dinosaur from South Dakota under the name of *Stegosaurus marshi*, stating that it probably represented a distinct genus, although owing to lack of material generic characters could not be stated. Curiously enough, failure to give a new generic name has resulted in the creation of a synonym. Better acquaintance with dinosaurs in general and Stegosauruses in particular has shown that the species is not a Stegosaur, but is nearly related to the English *Polacanthus*. With the present material it is only possible to say that the main apparent differences between *Polacanthus* and *Stegosaurus marshi* are the greater size of the latter and the larger and more varied dermal spines with which it was clad. In the light of my past experience, I shall, however, take the liberty of giving a new generic name to the species, and for that purpose propose *Hoplitosaurus* in allusion to its heavy armature.

This genus and the English *Polacanthus* and *Acanthopholis* are characterized by the sudden and considerable expansion of the long bones at their articular faces, a feature particularly noticeable in the humerus at its distal extremity. It may be said that in the Stegosauridae not more than two pairs of spines appear to be present and these are near the end of the tail. The main dermal armor is in the form of very large and thin plates running from the head to near the end of the tail. In the three genera named above, placed by Mr. Lydekker in the Scelidosauridae, the dermal armor consists of numerous flattened scutes and many large variously shaped spines.

F. A. LUCAS.