

on a four years' course of instruction satisfactorily, will raise the requirements for admission to a proper point and then concentrate all their strength upon a thoroughly sound course of two years leading to the bachelor's degree. More depends upon the strict enforcement of proper standards of admission to college than is generally believed; that is at present the weakest point in college administration. The general standard of college education in the United States would be strengthened more if the weaker colleges would fix and rigidly enforce proper entrance requirements and concentrate all their money and energies upon two years of thorough college work than if they continue to spread a college course over four years with admission secured on nominal terms or on none at all.

The policy outlined would, I think, largely increase the number of students seeking a college education, and many who might enter one of the stronger colleges for the two years' course would remain for four years. The loss of income due to the dropping out of students after two years of residence would be more than made good very soon by the large increase in college attendance.

As the system of higher education in the United States has developed it has become apparent that we have substituted three institutions—secondary school, college and university—for the two—secondary school and university—which exist in France and Germany. The work done in the United States by the best colleges is done in France and Germany one half by the secondary school and one half by the university. The training given in Europe differs in many ways from that given here, but from an administrative point of view the comparison just made is substantially correct. The college, as we have it, is peculiar to our own national system of education, and is

perhaps its strongest, as it certainly is its most characteristic, feature. It breaks the sharp transition which is so noticeable in Europe between the close surveillance and prescribed order of the secondary school and the absolute freedom of the university. Its course of liberal study comes just at the time in the student's life to do him most good, to open and inform his intelligence and to refine and strengthen his character. Its student life, social opportunities, and athletic sports are all additional elements of usefulness and of strength. It has endeared itself to three or four generations of the flower of our American youth and it is more useful to-day than at any earlier time.

For all of these reasons I am anxious to have it preserved as part of our educational system and so adjusted to the social and educational conditions which surround us that a college training may be an essential part of the higher education of an American whether he is destined to a professional career or to a business occupation. It seems to me clear that if the college is not so adjusted it will, despite its recent rapid growth, lose its prestige and place of honor in our American life, and that it may eventually disappear entirely, to the great damage of our whole educational system.

NICHOLAS MURRAY BUTLER.

*ADDRESS OF THE PRESIDENT OF THE
BRITISH ASSOCIATION FOR THE
ADVANCEMENT OF SCIENCE.*

III.

THE UPPER AIR AND AURORAS.

THE present liquid ocean, neglecting everything for the moment but the water, was at a previous period of the earth's history part of the atmosphere, and its condensation has been brought about by the gradual cooling of the earth's surface. This resulting ocean is subjected to the pressure of the remaining uncondensed

gases, and as these are slightly soluble they dissolve to some extent in the fluid. The gases in solution can be taken out by distillation or by exhausting the water, and if we compare their volume with the volume of the water as steam, we should find about one volume of air in 60,000 volumes of steam. This would then be about the rough proportion of the relatively permanent gas to condensable gas which existed in the case of the vaporized ocean. Now let us assume the surface of the earth gradually cooled to some 200 degrees below the freezing-point; then, after all the present ocean was frozen, and the climate became three times more intense than any arctic frost, a new ocean of liquid air would appear, covering the entire surface of the frozen globe about thirty-five feet deep. We may now apply the same reasoning to the liquid air ocean that we formerly did to the water one, and this would lead us to anticipate that it might contain in solution some gases that may be far less condensable than the chief constituents of the fluid. In order to separate them we must imitate the method of taking the gases out of water. Assume a sample of liquid air cooled to the low temperature that can be reached by its own evaporation, connected by a pipe to a condenser cooled in liquid hydrogen; then any volatile gases present in solution will distil over with the first portions of the air, and can be pumped off, being uncondensable at the temperature of the condenser. In this way, a gas mixture, containing, of the known gases, free hydrogen, helium and neon, has been separated from liquid air. It is interesting to note in passing that the relative volatilities of water and oxygen are in the same ratio as those of liquid air and hydrogen, so that the analogy between the ocean of water and that of liquid air has another suggestive parallel. The total uncondensable gas separated in this way amounts to about one

fifty-thousandth of the volume of the air, which is about the same proportion as the air dissolved in water. That free hydrogen exists in air in small amount is conclusively proved, but the actual proportion found by the process is very much smaller than Gautier has estimated by the combustion method. The recent experiments of Lord Rayleigh show that Gautier, who estimated the hydrogen present as one five-thousandth, has in some way produced more hydrogen than he can manage to extract from pure air by a repetition of the same process. The spectroscopic examination of these gases throws new light upon the question of the aurora and the nature of the upper air. On passing electric discharges through the tubes containing the most volatile of the atmospheric gases, they glow with a bright orange light, which is especially marked at the negative pole. The spectroscope shows that this light consists, in the visible part of the spectrum, chiefly of a succession of strong rays in the red, orange and yellow, attributed to hydrogen, helium and neon. Besides these, a vast number of rays, generally less brilliant, are distributed through the whole length of the visible spectrum. The greater part of these rays are of, as yet, unknown origin. The violet and ultra-violet part of the spectrum rivals in strength that of the red and yellow rays. As these gases probably include some of the gases that pervade interplanetary space, search was made for the prominent nebular, coronal and auroral lines. No definite lines agreeing with the nebular spectrum could be found, but many lines occurred closely coincident with the coronal and auroral spectrum. But before discussing the spectroscopic problem it will be necessary to consider the nature and condition of the upper air.

According to the old law of Dalton, supported by the modern dynamical theory of

gases, each constituent of the atmosphere while acted upon by the force of gravity forms a separate atmosphere, completely independent, except as to temperature, of the others, and the relations between the common temperature and the pressure and altitude for each specific atmosphere can be definitely expressed. If we assume the altitude and temperature known, then the pressure can be ascertained for the same height in the case of each of the gaseous constituents, and in this way the percentage composition of the atmosphere at that place may be deduced. Suppose we start with a surface atmosphere having the composition of our air, only containing two ten-thousandths of hydrogen, then at thirty-seven miles, if a sample could be procured for analysis, we believe that it would be found to contain 12 per cent. of hydrogen and only 10 per cent. of oxygen. The carbonic acid practically disappears; and by the time we reach forty-seven miles, where the temperature is *minus* 132 degrees, assuming a gradient of 3.2 degrees per mile, the nitrogen and oxygen have so thinned out that the only constituent of the upper air which is left is hydrogen. If the gradient of temperature were doubled, the elimination of the nitrogen and oxygen would take place by the time thirty-seven miles was reached, with a temperature of *minus* 220 degrees. The permanence of the composition of the air at the highest altitudes, as deduced from the basis of the dynamical theory of gases, has been discussed by Stoney, Bryan and others. It would appear that there is a consensus of opinion that the rate at which gases like hydrogen and helium could escape from the earth's atmosphere would be excessively slow. Considering that to compensate any such loss the same gases are being supplied by actions taking place in the crust of the earth, we may safely regard them as necessarily permanent con-

stituents of the upper air. The temperature at the elevations we have been discussing would not be sufficient to cause any liquefaction of the nitrogen and oxygen, the pressure being so low. If we assume the mean temperature as about the boiling-point of oxygen at atmospheric pressure, then a considerable amount of the carbonic acid must solidify as a mist, if the air from a lower level be cooled to this temperature; and the same result might take place with other gases of relatively small volatility which occur in air. This would explain the clouds that have been seen at an elevation of fifty miles, without assuming the possibility of water vapor being carried up so high. The temperature of the upper air must be above that on the vapor pressure curve corresponding to the barometric pressure at the locality, otherwise liquid condensation must take place. In other words, the temperature must be above the dew-point of air at that place. At higher elevations, on any reasonable assumption of temperature distribution, we inevitably reach a temperature where the air would condense, just as Fourier and Poisson supposed it would, unless the temperature is arrested in some way from approaching the zero. Both ultra-violet absorption and the prevalence of electric storms may have something to do with the maintenance of a higher mean temperature. The whole mass of the air above forty miles is not more than one seven-hundredth part of the total mass of the atmosphere, so that any rain or snow of liquid or solid air, if it did occur, would necessarily be of a very tenuous description. In any case, the dense gases tend to accumulate in the lower strata, and the lighter ones to predominate at the higher altitudes, always assuming that a steady state of equilibrium has been reached. It must be observed, however, that a sample of air taken at an elevation of nine miles

has shown no difference in composition from that at the ground, whereas, according to our hypothesis, the oxygen ought to have been diminished to 17 per cent., and the carbonic acid should also have become much less. This can only be explained by assuming that a large intermixture of different layers of the atmosphere is still taking place at this elevation. This is confirmed by a study of the motions of clouds about six miles high, which reveals an average velocity of the air currents of some seventy miles an hour; such violent winds must be the means of causing the intermingling of different atmospheric strata. Some clouds, however, during hot and thundery weather, have been seen to reach an elevation of seventeen miles, so that we have direct proof that on occasion the lower layers of atmosphere are carried to a great elevation. The existence of an atmosphere at more than a hundred miles above the surface of the earth is revealed to us by the appearance of meteors and fireballs, and when we can take photographs of the spectrum of such apparitions we shall learn a great deal about the composition of the upper air. In the meantime Pickering's solitary spectrum of a meteor reveals an atmosphere of hydrogen and helium, and so far this is corroborative of the doctrine we have been discussing. It has long been recognized that the aurora is the result of electric discharges within the limits of the earth's atmosphere, but it was difficult to understand why its spectrum should be so entirely different from anything which could be produced artificially by electric discharges through rarefied air at the surface of the earth. Writing in 1879, Rand Capron, after collecting all the recorded observations, was able to enumerate no more than nine auroral rays, of which but one could with any probability be identified with rays emitted by atmospheric air under an electric discharge. Vogel attributed

this want of agreement between nature and experiment, in a vague way, to difference of temperature and pressure; and Zollner thought the auroral spectrum to be one of a different order, in the sense in which the line and band spectra of nitrogen are said to be of different orders. Such statements were merely confessions of ignorance. But since that time observations of the spectra of auroras have been greatly multiplied, chiefly through the Swedish and Danish Polar Expeditions, and the length of spectrum recorded on the ultra-violet side has been greatly extended by the use of photography, so that, in a recent discussion of the results, M. Henri Stassano is able to enumerate upwards of one hundred auroral rays, of which the wavelength is more or less approximately known, some of them far in the ultra-violet. Of this large number of rays he is able to identify, within the probable limits of errors of observation, about two thirds as rays, which Professor Liveing and myself have observed to be emitted by the most volatile gases of atmospheric air unliquefiable at the temperature of liquid hydrogen. Most of the remainder he ascribes to argon, and some he might, with more probability, have identified with krypton or xenon rays, if he had been aware of the publication of wave-lengths of the spectra of those gases, and the identification of one of the highest rays of krypton with that most characteristic of auroras. The rosy tint often seen in auroras, particularly in the streamers, appears to be due mainly to neon, of which the spectrum is remarkably rich in red and orange rays. One or two neon rays are amongst those most frequently observed, while the red ray of hydrogen and one red ray of krypton have been noticed only once. The predominance of neon is not surprising, seeing that from its relatively greater proportion in air and its low den-

sity it must tend to concentrate at higher elevations. So large a number of probable identifications warrants the belief that we may yet be able to reproduce in our laboratories the auroral spectrum in its entirety. It is true that we have still to account for the appearance of some, and the absence of other, rays of the newly discovered gases, which in the way in which we stimulate them appear to be equally brilliant, and for the absence, with one doubtful exception, of all the rays of nitrogen. If we cannot give the reason of this, it is because we do not know the mechanism of luminescence—nor even whether the particles which carry the electricity are themselves luminous, or whether they only produce stresses causing other particles which encounter them to vibrate; yet we are certain that an electric discharge in a highly rarefied mixture of gases lights one element and not another, in a way which, to our ignorance, seems capricious. The Swedish North Polar Expedition concluded from a great number of trigonometrical measurements that the average above the ground of the base of the aurora was fifty kilometers (thirty-four miles) at Cape Thorsden, Spitzbergen; at this height the pressure of the nitrogen of the atmosphere would be only about one tenth of a millimeter, and Moissan and Deslandres have found that in atmospheric air at pressures less than one millimeter the rays of nitrogen and oxygen fade and are replaced by those of argon and by five new rays which Stassano identifies with rays of the more volatile gases measured by us. Also Collie and Ramsay's observations on the distance to which electrical discharges of equal potential traverse different gases explosively throw much light on the question; for they find that, while for helium and neon this distance is from 250 to 300 mm., for argon it is $45\frac{1}{2}$ mm., for hydrogen it is 39 mm., and for air and oxygen still less. This in-

dicates that a good deal depends on the very constitution of the gases themselves, and certainly helps us to understand why neon and argon, which exist in the atmosphere in larger proportions than helium, krypton or xenon, should make their appearance in the spectrum of auroras almost to the exclusion of nitrogen and oxygen. How much depends not only on the constitution and it may be temperature of the gases, but also on the character of the electric discharge, is evident from the difference between the spectra at the cathode and anode in different gases, notably in nitrogen and argon, and not less remarkably in the more volatile compounds of the atmosphere. Paulsen thinks the auroral spectrum wholly due to cathodic rays. Without stopping to discuss that question, it is certain that changes in the character of the electric discharge produce definite changes in the spectra excited by them. It has long been known that in many spectra the rays which are inconspicuous with an uncondensed electric discharge become very pronounced when a Leyden jar is in the circuit. This used to be ascribed to a higher temperature in this condensed spark, though measurements of that temperature have not borne out the explanation. Schuster and Hemsalech have shown that these changes of spectra are in part due to the oscillatory character of the condenser discharge which may be enhanced by self-induction, and the corresponding change of spectrum thereby made more pronounced. Lightning we should expect to resemble condensed discharge much more than aurora, but this is not borne out by the spectrum. Pickering's recent analysis of the spectrum of a flash obtained by photography shows, out of nineteen lines measured by him, only two which can be assigned with probability to nitrogen and oxygen, while three hydrogen rays most likely due to water are very conspicu-

ous, and eleven may be reasonably ascribed to argon, krypton and xenon, one to more volatile gas of the neon class, and the brightest ray of all is but a very little less refrangible than the characteristic auroral ray, and coincides with a strong ray of calcium, but also lies between, and close to, an argon and a neon ray, neither of them weak rays. There may be some doubt about the identification of the spectral rays of auroras because of the wide limits of the probable errors in measuring wave-lengths so faint as most of them are, but there is no such doubt about the wave-lengths of the rays in solar protuberances measured by Deslandres and Hale. Stassano found that these rays, 44 in number, lying between the Fraunhofer line *F* and 3,148 in the ultra-violet agree very closely with rays which Professor Liveing and myself measured in the spectra of the most volatile atmospheric gases. It will be remembered that one of the earliest suggestions as to the nature of solar prominences was that they were solar auroras. This supposition helped to explain the marvellous rapidity of their changes, and the apparent suspension of brilliant self-luminous clouds at enormous heights above the sun's surface. Now the identification of the rays of their spectra with those of the most volatile gases, which also furnish many of the auroral rays, certainly supports that suggestion. A stronger support, however, seems to be given to it by the results obtained at the total eclipse of May, 1901, by the American expedition to Sumatra. In the *Astrophysical Journal* for June last is a list of 339 lines in the spectrum of the corona photographed by Humphreys, during totality, with a very large concave grating. Of these no fewer than 209 do not differ from lines we have measured in the most volatile gases of the atmosphere, or in krypton or xenon, by more than one unit of wave-length on Arm-

strong's scale, a quantity within the limit of probable error. Of the remainder, a good many agree to a like degree with argon lines, a very few with oxygen lines, and still fewer with nitrogen lines; the characteristic green auroral ray, which is not in the range of Humphreys' photographs, also agrees within a small fraction of a unit of wave-length with one of the rays emitted by the most volatile atmospheric gas. Taking into account the Fraunhofer lines *H*, *K* and *G*, usually ascribed to calcium, there remain only fifty-five lines of the 339 unaccounted for to the degree of probability indicated. Of these considerably more than half are very weak lines which have not depicted themselves on more than one of the six films exposed, and extend but a very short distance into the sun's atmosphere. There are, however, seven which are stronger lines, and reach to a considerable height above the sun's rim, and all have depicted themselves on at least four of the six films. If there be no considerable error in the wave-lengths assigned (and such is not likely to be the case), these lines may perhaps be due to some volatile element which may yet be discovered in our atmosphere. However that may be, the very great number of close coincidences between the auroral rays and those which are emitted under electric excitement by gases of our atmosphere almost constrains us to believe, what is indeed most probable on other grounds, that the sun's coronal atmosphere is composed of the same substances as the earth's and that it is rendered luminous in the same way—namely, by electric discharges. This conclusion has plainly an important bearing on the explanation which should be given of the outburst of new stars and of the extraordinary and rapid changes in their spectra. Moreover, leaving on one side the question whether gases ever become luminous by the direct action of

heat, 'apart' from such transfers of energy as occur in chemical change and electric disturbance, it demands a revision of the theories which attribute more permanent differences between the spectra of different stars to differences of temperature, and a fuller consideration of the question whether they cannot with better reason be explained by differences in the electric conditions which prevail in the stellar atmosphere.

If we turn to the question what is the cause of the electric discharges which are generally believed to occasion auroras, but of which little more has hitherto been known than that they are connected with sunspots and solar eruptions, recent studies of electric discharges in high vacua, with which the names of Crookes, Röntgen, Lenard and J. J. Thomson will always be associated, have opened the way for Arrhenius to suggest a definite and rational answer. He points out that the frequent disturbances which we know to occur in the sun must cause electric discharges in the sun's atmosphere far exceeding any that occur in that of the earth. These will be attended with an ionization of the gases, and the negative ions will stream away through the outer atmosphere of the sun into the interplanetary space, becoming, as Wilson has shown, nuclei of aggregation of condensable vapors and cosmic dust. The liquid and solid particles thus formed will be of various sizes; the larger will gravitate back to the sun, while those with diameters less than one and a half thousandths of a millimeter, but nevertheless greater than a wave-length of light, will, in accordance with Clerk-Maxwell's electromagnetic theory, be driven away from the sun by the incidence of the solar rays upon them, with velocities which may become enormous, until they meet other celestial bodies, or increase their dimensions by picking up more cosmic dust or diminish

them by evaporation.' The earth will catch its share of such particles on the side which is turned towards the sun, and its upper atmosphere will thereby become negatively electrified until the potential of the charge reaches such a point that a discharge occurs, which will be repeated as more charged particles reach the earth. This theory not only accounts for the auroral discharges, and the coincidence of their times of greatest frequency with those of the maxima of sunspots, but also for the minor maxima and minima. The vernal and autumnal maxima occur when the line through the earth and sun has its greatest inclination to the solar equator, so that the earth is more directly exposed to the region of maximum of sunspots, while the twenty-six days' period corresponds closely with the period of rotation of that part of the solar surface where faculæ are most abundant. J. J. Thomson has pointed out, as a consequence of the Richardson observations, that negative ions will be constantly streaming from the sun merely regarded as a hot body, but this is not inconsistent with the supposition that there will be an excess of this emission in eruptions, and from the regions of faculæ. Arrhenius' theory accounts also, in a way which seems the most satisfactory hitherto enunciated, for the appearances presented by comets. The solid parts of these objects absorb the sun's rays, and as they approach the sun become heated on the side turned towards him until the volatile substances frozen in or upon them are evaporated and diffused in the gaseous state in surrounding space, where they get cooled to the temperature of liquefaction and aggregated in drops about the negative ions. The larger of these drops gravitate towards the sun and form clouds of the coma about the head, while the smaller are driven by the incidence of the sun's light upon them away from the sun and form the tail. The

curvature of the tail depends, as Bredichin has shown, on the rate at which the particles are driven, which in turn depends on the size and specific gravity of the particles, and these will vary with the density of the vapor from which they are formed and the frequency of the negative ions which collect them. In any case Arrhenius' theory is a most suggestive one, not only with reference to auroras and comets, and the solar corona and chromosphere, but also as to the constitution of the photosphere itself.

VARIOUS LOW-TEMPERATURE RESEARCHES.

We may now summarize some of the results which have already been attained by low-temperature studies. In the first place, the great majority of chemical interactions are entirely suspended, but an element of such exceptional powers of combination as flourine is still active at the temperature of liquid air. Whether solid fluorine and liquid hydrogen would interact no one can at present say. Bodies naturally become denser, but even a highly expansive substance like ice does not appear to reach the density of water at the lowest temperature. This is confirmatory of the view that the particles of matter under such conditions are not packed in the closest possible way. The force of cohesion is greatly increased at low temperatures, as is shown by the additional stress required to rupture metallic wires. This fact is of interest in connection with two conflicting theories of matter. Lord Kelvin's view is that the forces that hold together the particles of bodies may be accounted for without assuming any other agency than gravitation or any other law than the Newtonian. An opposite view is that the phenomena of the aggregation of molecules depend upon the molecular vibration as a physical cause. Hence, at the zero of absolute temperature, this vibrating energy

being in complete abeyance, the phenomena of cohesion should cease to exist, and matter generally be reduced to an incoherent heap of cosmic dust. This second view receives no support from experiment.

The photographic action of light is diminished at the temperature of liquid air to about twenty per cent. of its ordinary efficiency, and at the still lower temperature of liquid hydrogen only about ten per cent. of the original sensitivity remains. At the temperature of liquid air or liquid hydrogen a large range of organic bodies and many inorganic ones acquire under exposure to violet light the property of phosphorescence. Such bodies glow faintly so long as they are kept cold, but become exceedingly brilliant during the period when the temperature is rising. Even solid air is a phosphorescent body. All the alkaline earth sulphides which phosphoresce brilliantly at the ordinary temperature lose this property when cooled, to be revived on heating; but such bodies in the first instance may be stimulated through the absorption of light at the lowest temperatures. Radio-active bodies, on the other hand, like radium, which are naturally self-luminous, maintain this luminosity unimpaired at the very lowest temperatures, and are still capable of inducing phosphorescence in bodies like the platino-cyanides. Some crystals become for a time self-luminous when cooled in liquid air or hydrogen, owing to the induced electric stimulation causing discharges between the crystal molecules. This phenomenon is very pronounced with nitrate of uranium and some platino-cyanides.

In conjunction with Professor Fleming a long series of experiments was made on the electric and magnetic properties of bodies at low temperatures. The subjects that have been under investigation may be

classified as follows: 'The Thermo-Electric Powers of Pure Metals'; 'The Magnetic Properties of Iron and Steel'; 'Dielectric Constants'; 'The Magnetic and Electric Constants of Liquid Oxygen'; 'Magnetic Susceptibility.'

The investigations have shown that electric conductivity in pure metals varies almost inversely as the absolute temperature down to *minus* 200 degrees, but that this law is greatly affected by the presence of the most minute amount of impurity. Hence the results amount to a proof that electric resistance in pure metals is closely dependent upon the molecular or atomic motion which gives rise to temperature, and that the process by which the energy constituting what is called an electric current is dissipated essentially depends upon non-homogeneity of structure and upon the absolute temperature of the material. It might be inferred that at the zero of absolute temperature resistance would vanish altogether, and all pure metals become perfect conductors of electricity. This conclusion, however, has been rendered very doubtful by subsequent observations made at still lower temperatures, which appear to point to an ultimate finite resistance. Thus the temperature at which copper was assumed to have no resistance was *minus* 223 degrees, but that metal has been cooled to *minus* 253 degrees without getting rid of all resistance. The reduction in resistance of some of the metals at the boiling-point of hydrogen is very remarkable. Thus copper has only one per cent., gold and platinum three per cent., and silver four per cent. of the resistance they possessed at zero C., but iron still retains twelve per cent. of its initial resistance. In the case of alloys and impure metals, cold brings about a much smaller decrease in resistivity, and in the case of carbon and insulators like guttapercha,

glass, ebonite, etc., their resistivity steadily increases. The enormous increase in resistance of bismuth when transversely magnetized and cooled was also discovered in the course of these experiments. The study of dielectric constants at low temperatures has resulted in the discovery of some interesting facts. A fundamental deduction from Maxwell's theory is that the square of the refractive index of a body should be the same number as its dielectric constant. So far, however, from this being the case generally, the exceptions are far more numerous than the coincidences. It has been shown in the case of many substances, such as ice and glass, that an increase in the frequency of the alternating electromotive force results in a reduction of the dielectric constant to a value more consistent with Maxwell's law. By experiments upon many substances it is shown that even a moderate increase of frequency brings the large dielectric constant to values quite near to that required by Maxwell's law. It was thus shown that low temperature has the same effect as high frequency in annulling the abnormal dielectric values. The exact measurement of the dielectric constant of liquid oxygen as well as its magnetic permeability, combined with the optical determination of the refractive index, showed that liquid oxygen strictly obeys Maxwell's electro-optic law even at very low electric frequencies. In magnetic work the result of greatest value is the proof that magnetic susceptibility varies inversely as the absolute temperature. This shows that the magnetization of paramagnetic bodies is an affair of orientation of molecules, and it suggests that at the absolute zero all the feebly paramagnetic bodies will be strongly magnetic. The diamagnetism of bismuth was found to be increased at low temperatures. The magnetic moment of a steel magnet is tempo-

rarily increased by cooling in liquid air, but the increase seems to have reached a limit, because on further cooling to the temperature of liquid hydrogen hardly any further change was observed. The study of the thermo-electric relations of the metals at low temperatures resulted in a great extension of the well-known Tait thermo-electric diagram. Tait found that the thermo-electric power of the metals could be expressed by a linear function of the absolute temperature, but at the extreme range of temperature now under consideration this law was found not to hold generally; and further, it appeared that many abrupt electric changes take place, which originate probably from specific molecular changes occurring in the metal. The thermo-electric neutral points of certain metals, such as lead and gold, which are located about or below the boiling-point of hydrogen, have been found to be a convenient means of defining specific temperatures in this exceptional part of the scale.

The effect of cold upon the life of living organisms is a matter of great intrinsic interest, as well as of wide theoretical importance. Experiment indicates that moderately high temperatures are much more fatal, at least to the lower forms of life, than are exceedingly low ones. Professor McKendrick froze for an hour at a temperature of 182° C. samples of meat, milk, etc., in sealed tubes; when these were opened after being kept at blood heat for a few days, their contents were found to be quite putrid. More recently some more elaborate tests were carried out at the Jenner Institute of Preventive Medicine on a series of typical bacteria. These were exposed to the temperature of liquid air for twenty hours, but their vitality was not affected, their functional activities remained unimpaired, and the cultures which they yielded were normal in every respect.

The same result was obtained when liquid hydrogen was substituted for air. A similar persistence of life in seeds has been demonstrated even at the lowest temperatures; they were frozen for over a hundred hours in liquid air, at the instance of Messrs. Brown and Escombe, with no other result than to affect their protoplasm with a certain inertness, from which it recovered with warmth. Subsequently commercial samples of barley, pea, vegetable marrow, and mustard seeds were literally steeped for six hours in liquid hydrogen at the Royal Institution, yet when they were sown by Sir W. T. Thiselton Dyer at Kew in the ordinary way, the proportion in which germination occurred was no less than in the other batches of the same seeds which had suffered no abnormal treatment. Bacteria are minute vegetable cells, the standard of measurement for which is the 'mikron.' Yet it has been found possible to completely triturate these microscopic cells, when the operation is carried out at the temperature of liquid air, the cells then being frozen into hard breakable masses. The typhoid organism has been treated in this way, and the cell plasma obtained for the purpose of studying its toxic and immunizing properties. It would hardly have been anticipated that liquid air should find such immediate application in biological research. A research by Professor Macfadyen, just concluded, has shown that many varieties of microorganisms can be exposed to the temperature of liquid air for a period of six months without any appreciable loss of vitality, although at such a temperature the ordinary chemical processes of the cell must cease. At such a temperature the cells cannot be said to be either alive or dead, in the ordinary acceptance of these words. It is a new and hitherto unobtained condition of living matter—a third state. A final in-

stance of the application of the above methods may be given. Certain species of bacteria during the course of their vital processes are capable of emitting light. If, however, the cells be broken up at the temperature of liquid air, and the crushed contents brought to the ordinary temperature, the luminosity function is found to have disappeared. This points to the luminosity not being due to the action of a ferment—a ‘Luciferase’—but as being essentially bound up with the vital processes of the cells, and dependent for its production on the intact organization of the cell. These attempts to study by frigorific methods the physiology of the cell have already yielded valuable and encouraging results, and it is to be hoped that this line of investigation will continue to be vigorously prosecuted at the Jenner Institute.

And now, to conclude an address which must have sorely taxed your patience, I may remind you that I commenced by referring to the plaint of Elizabethan science, that cold was not a natural available product. In the course of a long struggle with nature, man, by the application of intelligent and steady industry, has acquired a control over this agency which enables him to produce it at will, and with almost any degree of intensity, short of a limit defined by the very nature of things. But the success in working what appears, at first sight, to be a quarry of research that would soon suffer exhaustion, has only brought him to the threshold of new labyrinths, the entanglements of which frustrate, with a seemingly invulnerable complexity, the hopes of further progress. In a legitimate sense all genuine scientific workers feel that they are ‘the inheritors of unfulfilled renown.’ The battlefields of science are the centers of a perpetual warfare, in which there is no hope of final victory, although partial conquest is ever

triumphantly encouraging the continuance of the disciplined and strenuous attack on the seemingly impregnable fortress of Nature. To serve in the scientific army, to have shown some initiative, and to be rewarded by the consciousness that in the eyes of his comrades he bears the accredited accolade of successful endeavor, is enough to satisfy the legitimate ambition of every earnest student of Nature. The real warranty that the march of progress in the future will be as glorious as in the past lies in the perpetual reinforcement of the scientific ranks by recruits animated by such a spirit, and proud to obtain such a reward.

JAMES DEWAR.

SCIENTIFIC BOOKS.

Notes on Naval Progress. July, 1902. Office of Naval Intelligence. Washington, Government Printing Office. 1902. 8vo. Paper. Pp. 502; over 100 illustrations, plates, maps, tables, etc.

This very large and exceedingly valuable document constitutes No. XXI, General Information Series, of the Office of Naval Intelligence, a division of the Naval Organization which has now for many years been justifying its existence by great and increasing efficiency. Under the supervision of Captain Sigsbee, the present Chief Intelligence Officer, it is evidently fully maintaining its standing. The contributors to this bulky volume are usually young officers of the navy who exhibit that talent for exact, concise and comprehensive composition which is the distinguishing characteristic of a good official report, and that excellence in style which seems so common a talent with military and naval officers. The two probably necessarily go together and are the outcome of familiarity with, often a minute study of, the reports and writings of great commanders quite as much as of careful drill at the governmental technical schools.

The volume in hand contains notes on ships and torpedo-boats, on ordnance and armor, on engineering progress, electricity, wireless