SCIENCE

FRIDAY, JUNE 4, 1915

CONTENTS

The Electrical Photometry of Stars: Professor Joel Stebbins	809
Mr. Edison's Service for Science: PRESIDENT RICHARD C. MACLAURIN	813
The Proceedings of the National Academy as a Medium of Publication: Dr. GEORGE EL- LERY HALE	815
The Seattle Meeting of the American Chem- ical Society: Dr. CHAS. L. PARSONS	817
Scientific Notes and News	818
University and Educational News	822
Discussion and Correspondence:— Bird Collecting and Ornithology: DR. WIL- LARD G. VAN NAME. Fundamental Equa- tions of Mechanics: T. L. PORTER AND R. C. GOWDY. Another State Park Needed: R. C. BENEDICT	823
Scientific Books:— Stevens's Theory of Measurements: PRO- FESSOR A. DE FOREST PALMER. Child on Electric Arcs: PROFESSOR R. G. HUDSON	828
Scientific Journals and Articles	829
Scientific Results of the Terra Nova Expedi- tion: Dr. Edward W. BERRY	830
Special Articles:— A. Botanical Index of Cretaceous and Ter- tiary Climates: I. W. BALLEY AND E. W. SINNOTT. The Brown Grape Aphil: A. C. BAKER AND W. F. TURNER. The Relation of Mitochondria to Granules of the Vital Azo Dyes: KATHERINE J. SCOTT	831
The American Philosophical Society: PRO- FESSOR HORACE CLARK RICHARDS	835
The American Physical Society: Professor A. D. Cole	841
The Entomological Society of America: PRO- FESSOR ALEX. D. MACGILLIVRAY	842
The Indiana Academy of Science: Dr. A. J. BIGNEY	843
<i>(</i>)	

THE ELECTRICAL PHOTOMETRY OF STARS1

In measures of the light of stars there are some advantages and some drawbacks as compared with photometric work in the laboratory. First of all, we are not concerned with absolute measures of intensity. but what we want to know is how the light of a heavenly body varies. If the light is constant, there is not much to be learned, but if it changes, we may infer a great deal from the law of variation. In laboratory and commercial photometry, it is customary to measure what may be called the visual brightness of a source of light, but with the stars it is immaterial for many purposes whether we study the changes of the red, or the blue, or any other part of the spectrum, though in fact any complete stellar photometry should include measures in all regions, infra-red, visible and ultra-violet.

The chief disadvantage in stellar photometry is that the stars are so faint that it is usually not feasible to expand their images out into surfaces, and most forms of stellar photometer depend upon comparisons of two point images by the eye. Although the eye is a wonderful instrument, especially in the range of intensity over which it may be used, the limit of accuracy attained by looking first at one light and then at another is much the same as though instead of using a balance we should weigh objects by lifting them in our hands. It is safe to say that no observer has ever been able to get visual results accurate to 1 per cent., and in the best measures there are occasional errors of 10 per cent., 20 per cent. and even more. It was hoped that the introduc-

¹ Read at the meeting of the National Academy of Sciences, April 20, 1915.

MSS. intended for publication and books, etc., intended for review should be sent to Professor J McKeen Cattell, Garrigonon-Hudson, N. Y