SCIENCE

FRIDAY, OCTOBER 15, 1915

CONTENTS The Physiological Importance of Phase Boundaries: Professor W. M. Bayliss 509 The Publication of the Results of Investigations made in Experiment Stations in Technical Scientific Journals: Dr. RAYMOND PEARL 518 The Naval Consultation Board 522 Scientific Notes and News 523 University and Educational News 526 Discussion and Correspondence:-Potassium from the Soil: Professor E. W. Elementary Mechanics: Pro-FESSOR EDWIN BIDWELL WILSON. The End is not yet: Dr. WITMER STONE. The "Pan-American Scientific Congress': H. A. 527 Scientific Books:-Riley and Johannsen's Medical Entomology: W. H. HUNTER. Abderhalden on Abwehrfermente: Dr. John Auer. Olcott's Sun-lore of All Ages: Professor Chas. Hemoglobinophilic Bacteria: Dr. David John Davis 532 Special Articles:-Artificial Daylight for the Microscope: PROFESSOR SIMON H. GAGE. A New Alfalfa Leaf-spot in America: LEO E. MELCHERS. Differentiation of Wandering Mesenchymal Cells in the Living Yolk-sac: Dr. C. R. STOCKARD 534 Anthropology at the San Francisco Meeting:

MSS. intended for publication and books, etc., intended for review should be sent to Professor J. McKeen Cattell, Garrisonon-Hudson. N. Y.

PROFESSOR GEORGE GRANT MACCURDY 541

THE PHYSIOLOGICAL IMPORTANCE OF PHASE BOUNDARIES¹

Even a hasty consideration of the arrangements present in living cells is sufficient to bring conviction that the physical and chemical systems concerned operate under conditions very different from those of reactions taking place between substances in true solution. We become aware of the fact that there are numerous constituents of the cell which do not mix with one another. In other words, the cell system is one of many "phases," to use the expression introduced by Willard Gibbs.

Further, parts of this system which appear homogeneous under the ordinary microscope are shown by the ultra-microscope to be themselves heterogeneous. These are in what is known as the colloidal state. Some dispute has taken place as to whether this state is properly to be called a heterogeneous one, but it is sufficient for our purpose to note that investigation shows that the interfaces of contact between the components of such systems are the seat of the various forms of energy which we meet with in the case of systems obviously consisting of phases which can be separated mechanically, so that considerations applying to coarsely heterogeneous systems apply also to colloidal systems. Although the phases of a colloidal system can not be so obviously and easily separated as those of an ordinary heterogeneous one, this can be almost completely by filtration through membranes such as the gelatin in

¹ Address of the President of the Section of Physiology at the Manchester Meeting of the British Association for the Advancement of Science.