

London the collection is well forward. It should again be pointed out that every ton of nuts gathered means a saving of half a ton of grain. Present indications are that at least 25,000 tons of nuts will reach the Ministry of Munitions, but this is only about one eighth of the estimated crop for the country.

UNIVERSITY AND EDUCATIONAL NEWS

A BEQUEST of \$200,000 is left to Yale University by the terms of the will of the late Richard P. Sewell of Boston.

H. P. WOOD, head of the department of electrical engineering at the Georgia School of Technology, Atlanta, Ga., has been appointed president of the Academic Board of the United States Army School of Military Aeronautics, which has been established at the Georgia School of Technology.

PROFESSOR J. F. WILSON, who during the past year was professor of electrical engineering at Queen's University, Kingston, Ontario, has been appointed assistant professor of electrical engineering at the University of Southern California, Los Angeles.

DR. JOHN EDWARD MARR, F.R.S., fellow of St. John's College since 1881, university lecturer in geology at Cambridge University, has been elected to the Woodwardian professorship of geology in succession to the late Professor Hughes.

DISCUSSION AND CORRESPONDENCE METHODS FOR PREPARING ANIMAL MATERIAL TO BE DISSECTED

POSSIBLY the most common fixing and preserving fluid used for dissecting material is formalin. It is relatively inexpensive and especially convenient for collecting expeditions where a concentrated fluid is desirable. Animals preserved in it have rigid joints, however, and every one is familiar with the disagreeable characteristics of such material during dissection. Alcohol is much better from the standpoint of the dissector, but it has limitations when used alone.

Some of the "embalming fluid" mixtures used in preparing human cadavers for dissec-

tion are also splendid for smaller animals. Those containing phenol, alcohol and glycerine with no formalin give relatively flexible joints and pliable tissues. They also render the material resistant to a large amount of drying in the open air of a laboratory during dissection. Phenol is a relatively non-volatile antiseptic, and glycerine is very effective in preventing drying. Alcohol counteracts the action of the phenol in the solution, on the hands of the dissector. A good and much used solution consists of equal parts of phenol, alcohol and glycerine. Another less expensive fluid with arsenic and considerable water added to the above was described by Dr. W. C. Lusk some years ago¹ with an excellent discussion of principles involved in preparing cadavers for dissection.

As penetration by such fluids is slow, the mixture should be injected through some large artery, a femoral or carotid in the case of mammals. Small animals may be placed in solutions of about 80 per cent. alcohol in water when it is not practicable to inject them. In such cases, the usual practise of making a slit, at least in the ventral abdominal wall, should be followed. After all the tissues have been fixed, the material may be removed to a container which holds an "embalming fluid," such as I have mentioned, much diluted with water. Ten or more parts of water to one of the "embalming fluid" may be used. In fact, I have kept material which had already been thoroughly fixed in either formalin or alcohol, for several years in a solution consisting of water with 1 to 2 per cent. of phenol and 5 to 10 per cent. glycerine, with or without a little alcohol. Single specimens thus preserved have been used in dissection for many months without deterioration, so long as they were not kept out of the solution for more than a few hours or so at a time.

It is customary in human anatomy to leave cadavers on the dissecting tables for months without soaking. The glycerine in their tissues is wonderfully effective in checking drying. Nevertheless, unless the atmosphere of the room is very moist a good deal of drying

¹ *Anat. Record*, Vol. 3, No. 1.