

in the starchy central core of the typha rhizome was shown by A. P. Claassen, who estimated that one acre would yield a total dry weight of 10,792 pounds of cat-tail rhizomes, or more than two tons of flour, made from the central core.

Typha may be used as a substitute for high-priced corn. It would seem that the best time for feeding would be in the fall and winter, as the starchy content is likely to be highest then.

L. E. FREUDENTHAL

ROSALIE FARM,  
LAS CRUCES, NEW MEXICO

#### SOIL SHIFTING AND DEPOSITS

MR. PETERSON'S article on deposition of soil in the Palouse area of eastern Washington and Idaho, which appeared in *SCIENCE*, January 27, 1922, should prove of interest and value to foresters as well as agriculturists in this region. The questions naturally arise: How far is this soil carried into the Bitterroot mountains, and how does it influence the character of the soil and vegetation within the forest areas? The writer's observations in this respect may be of interest in this connection.

Dust storms, commonly referred to as "Palousers," are of comparatively frequent occurrence throughout northern Idaho and northwestern Montana. They accompany high winds from the west and southwest; they are well known and despised by housekeepers in Kalispell, Missoula, Thompson Falls, Libby and all surrounding towns. The dust penetrates into every house and office. When accompanied by rain the window panes and buildings are besmirched with streaks of red soil. One of these storms in March, 1917, laid down on the snow within the timbered region of northern Idaho about 600 pounds of dust per acre. The dust from that storm hung on the trees, even at 6,000 feet elevation, along the Kootenai-Priest Divide throughout the summer of 1917. Settlers say that dust storms are common along the Cœur d'Alene, St. Joe and Clearwater rivers.

The writer has noted the billowy soil surface, unmistakably due to surface shifting of the soil, as far east as Pierce, Idaho, about

eighty miles east of Moscow. The soil is unusually deep and fertile and the vegetation is more profuse, with better growth of timber, over the larger portion of the Clearwater Forest in Idaho than occurs on the forests farther north or on the forests of western Montana. It is of interest to note that the Clearwater forest lies directly in the path of the strong west winds from the arid parts along the Columbia River, and that Lewis and Clark, as early as 1806, called attention to the unusually deep and seemingly fertile soil in the Clearwater basin.

These observations lead to the supposition that the accumulation and shifting of soil on the Palouse area have been effective in preventing natural establishment of the forest here in the past, though climatic records indicate that the area should grow western yellow pine; and they strengthen the belief that the unusually good growth of timber, profuse vegetation, and deep soils on certain parts of the western slopes of the Bitterroot mountains in Idaho, are due partly to the fact that soil is carried in by the westerly winds from lava plateaus along the Snake and Columbia rivers.

J. A. LARSEN

MISSOULA, MONTANA

#### QUOTATIONS

##### AN INTERNATIONAL LANGUAGE

THERE is an increasing demand among scientific men for international agreement as to the choice of a universal auxiliary language. After a long struggle, many of the fundamental tools of thought have been unified. All nations now use the same system of numbers, Arabic numerals, measurements of latitude and longitude, mathematical symbols, chemical formulæ, and, at least in science, the metric system.

But language, the master-key to thought and the vehicle of communication, remains under the curse of Babel. Were it possible by acquiring a second language in addition to the natal language to convey ideas to fellow-workers in every part of the world and to receive their ideas, one of the greatest barriers to the progress of science would be broken down. Time and money would be saved, overlapping of