

even now it occupies an important position among the remedies that are available for the treatment of neuro-syphilis. Professor Leake's enthusiastic estimate of its value may be prophetic but can not be accepted as an accomplished fact.

Finally, there is some doubt as to whether one would be justified in referring to the results of the clinical investigations carried out at the University of Wisconsin as an "achievement of this cooperative research." All the work on which the application of tryparsamide to the treatment of neuro-syphilis was based was done by an independent group of workers; the study of tryparsamide by Lorenz, Loevenhart and their associates was not in any sense a chance study of "arsenical compounds which had been discarded for use in the general types of syphilitic infection," but this study was undertaken in response to a specific suggestion based on the results of laboratory and preliminary clinical investigations. The connection of the Wisconsin group of workers with the development of tryparsamide was, therefore, essentially the same as that of Alt and other clinicians with the development of salvarsan in that they were given the opportunity of carrying out one phase of the clinical investigations necessary to the determination of the probable therapeutic value of tryparsamide.

WADE H. BROWN

ROCKEFELLER INSTITUTE FOR
MEDICAL RESEARCH

IMMUNITY TO TUBERCULOSIS BY THE INJECTION OF EXTRACT OF HEART MUSCLE

THE article by Professor Elliott C. Prentiss in the issue of *SCIENCE* for July 31 prompts me to make a brief report of some experimental work undertaken a goodly number of years ago, but for various extraneous reasons never carried to completion. In my work as pathologist to the Philadelphia General Hospital and to the University Hospital, I was struck by the rarity, in fact, the virtual non-occurrence, of tuberculosis of the heart muscle. This seemed to me to have but one of two explanations: (a) that the activity of the intracardiac circulation was such that the tubercle bacillus had no chance to lodge; (b) that the immunity of the myocardium was due to a biologic property, to some antituberculous substance in the muscle. Assuming that the latter was the more probable reason, I undertook a series of experiments in which I made extracts of beef's heart and mixed the filtrate obtained with virulent tubercle bacilli and injected the suspension into guinea pigs. In this work I had the help of Dr. Karl F. Meyer, now of California. It seemed to me that the material had a dis-

tinct inhibitory influence. Unfortunately, Dr. Meyer left Philadelphia for his new field, while my own work in clinical medicine and teaching came to absorb nearly all my time. I have proposed a repetition of the experiments to several men whose facilities and opportunities for research of this character were greater than my own, but so far no one has begun to work actively upon the problem.

If the immunity of the heart muscle and of the skeletal muscles is, as I believe, biological rather than mechanical, then it may be possible to extract an active agent by suitable means. In my own experiments pieces of heart muscle were cut up into small fragments under aseptic conditions and the fragments put into a grinding machine with china balls. The juice was then pressed out under great pressure and finally filtered through a Berkefeld filter. The filtrate, a golden colored liquid, was then used for the experiments. It is highly probable that the procedure could now be greatly simplified.

DAVID RIESMAN

UNIVERSITY OF PENNSYLVANIA

A COMBINED CULTURE MEDIUM AND INDICATOR FOR PARAMOECIUM

IN the April 10, 1925, issue of *SCIENCE*, Bragg and Hulpieu describe the effect of a stain obtained from red cabbage leaves as an indicator of the acidity of the food vacuoles of *Paramecium*. I have been unable to secure similar satisfactory results with the races I am using, but have found that a dilute infusion of red cabbage leaves (about 30 grams to one liter of water) is an excellent medium in which the animals not only reproduce rapidly but at the same time the color of the infusion indicates the chemical condition of the culture. When fresh, the cabbage leaf culture medium is light reddish purple in color, but about twenty-four hours after being seeded with *Paramecium*, it turns red, indicating the formation of acid. In four or five days to two weeks, as the paramecia increase in number, the medium gradually becomes alkaline, as is shown by its change of color to green.

The culture, as far as quantity of paramecia is concerned, is at its height when it becomes a brilliant green and has lost its early turbidity. The behavior of the cultures can be varied considerably by adding a trace of sodium bicarbonate or a weak acid. In from one to two months, the culture becomes the color of an old hay infusion, fails to react to acids or alkalies and the paramecia have either wholly or almost wholly died off.

ROBERT T. HANCE

ROCKEFELLER INSTITUTE
FOR MEDICAL RESEARCH