

eral government, except in the case of commitments already made to bring new areas under cultivation.

He will have charge of the courses on weather and climate and geomorphology.

UNIVERSITY AND EDUCATIONAL NOTES

HARVARD UNIVERSITY has received an anonymous gift of \$3,000,000 to build and endow a residence college of the type of the colleges of the Universities of Oxford and Cambridge.

Two bequests amounting to \$700,000 contained in the will of Charles Lennig, who died thirty-seven years ago, have become available to the University of Pennsylvania at the final distribution of the estate. One bequest of \$500,000 creates the "Charles Lennig Fund in Aid of Instruction in Theoretical and Practical Mechanics." Its income will be used for the acquisition of scientific works, structures, instruments, machines and material for the Towne Scientific School of the university. The other bequest of \$200,000 establishes the "Charles Lennig Beneficiary Fund," the income from which will be devoted to providing free scholarships.

GROUND has been broken on the Columbia University campus for a new building to house the natural science department, which will be erected at a cost of \$1,000,000. It will be situated in the southeast corner of the Grove on Amsterdam Avenue, facing 119th Street, and will be ten stories in height. Architecturally it will be almost identical with the Chandler Chemical Laboratories.

DR. ALEXANDER RUTHVEN has been appointed dean of administration with the duties of vice-president of the University of Michigan.

DR. LEON E. SMITH, of the Randal Morgan Laboratory of Physics of the University of Pennsylvania, has accepted the position of professor of physics and head of the department of physics at Denison University, Granville, Ohio.

DR. AMOS M. SHOWALTER, National Research Fellow, 1924-27, has been appointed assistant professor of botany in Washington University.

DR. HARRY HELSON, of the department of psychology of Cornell University, is now associate professor of experimental psychology in Bryn Mawr College, taking the place of Dr. Clarence E. Ferree, who is now at the Johns Hopkins University.

DR. GUY HAROLD SMITH, who was a member of the department of geography of the Ohio State University last year during the absence of Dr. Roderick Peattie, has joined the faculty of the department of geology and geography of the University of Illinois.

DISCUSSION AND CORRESPONDENCE

ON NUCLEAR DERIVATIVES AND THE LETHAL ACTION OF ULTRA-VIOLET LIGHT

THE bactericidal action of ultra-violet light has been known for fifty years, and has been repeatedly investigated. But few investigators have sought the mechanism of the reaction or the chemical units of the bacterial protoplasm so affected by the ultra-violet energy as to prevent the subsequent multiplication of the cells.

If measured monochromatic ultra-violet energy is used to kill bacteria such as *S. aureus*, lying in a single plane, and its effect is recorded statistically, characteristic and similar curves are produced at each wave-length studied. These curves show that an appreciable amount of energy must be incident on the bacteria before any of them succumb. With longer exposures they succumb along a gradient that is for the most part apparently exponential, but experimental evidence indicates that its course is determined by differences in the resistance of single bacteria, and that the curve is therefore one of probability. Wide differences are found in the incident energies required to produce these curves at different wave-lengths, and if the same points on each gradient (say 50 per cent. destruction) are joined by a smooth curve, its shape is such as to suggest immediately that it is reciprocally related to the absorption of ultra-violet energy by some sensitive element in the bacterial protoplasm.

In 1917 Harris and Hoyt¹ suggested that "the susceptibility of protoplasm to ultra-violet light is conditioned by the selective absorption of the toxic rays by the aromatic amino acid radicals of the proteins." Their conclusion was based on the observation that a screen or filter of an aromatic amino acid solution—tyrosine or aminobenzoic acid, for example—greatly prolonged the exposure to the quartz mercury arc necessary to kill *Paramecia*, and that therefore these substances must be absorbing the very wave-lengths responsible for the lethal changes in these organisms.

Such a conclusion, however, does not exclude the possibility that other biochemical entities essential to life may also show a selective absorption over the toxic range, and it may be that the lethal reaction is due to some, or some one, of these other substances.

Recently a further search has been made for the substance most probably involved. Since the nucleus

¹ F. I. Harris and H. S. Hoyt, SCIENCE, 1917, N.S. 46: 318.