

Over a range of several different temperatures marked differences in varieties could be noted.

BOTANY DEPARTMENT, STUART DUNN
NEW HAMPSHIRE AGRICULTURAL
EXPERIMENT STATION

THE USE OF SOLID CARBON DIOXIDE IN MAKING FREEZING-POINT DETERMINATIONS WITH PLANT JUICES

DUNN¹ has recently employed "Dry Ice" as a refrigerant in subjecting tissues of apple stems to freezing temperatures.

The ease with which this material can be handled and the ease with which low temperatures can be obtained and maintained caused the writer to test out its use as a refrigerant in determining the freezing-point depression of plant juices.

Although others may possibly have employed Dry Ice for this purpose, it seems to the writer that knowledge of this method should receive more publicity. Because of the ease of manipulation and the cleanliness and rapidity of the method, it seems as though its use would be of value to workers in the plant sciences.

In this method an ether bath surrounds the air jacket of the usual Beckman freezing-point apparatus. The temperature of the bath is regulated by adding to it pieces of Dry Ice (solid CO₂) until the desired temperature is reached. When the pieces of Dry Ice drop into the ether they sink, causing at the same time violent bubbling of the liquid, while the temperature of the bath becomes lower. The bubbling causes the temperature of the bath to become uniform throughout, dispensing thus with stirring.

The air-jacket may be filled with ether or alcohol if desired. The ether bath should be in a cylindrical container, tall enough to accommodate the air-jacket and its contents and the auxiliary thermometer. The container may be of metal, enamelware or glass, and should have a capacity preferably of two to two and one half liters, and tall enough to accommodate the freezing tube.

The desired temperature of the bath may be maintained during an experiment by adding small pieces of solid CO₂ whenever there is a tendency for the temperature of the bath to rise. The apparatus should be placed in sawdust or some other insulating material.

The advantages of using ether for a bath lie in the following points: (1) Ordinary ether is cheap. (2) The freezing-point of ether is very low. (3) Ether is volatile, leaving the apparatus clean after its use. (4) In other liquids that are volatile and

do not adhere to the parts of the apparatus in contact with them, difficulty is obtained in lowering the temperature of the bath if water is present. For this reason alcohol has been discarded as a bath. In alcohol the pieces of CO₂ become coated with ice which retards or stops the volatilization of the CO₂.

It is apparent that this method excels as cooling is produced by drawing air through the ether bath causing rapid evaporation therein because of the more rapid lowering of the temperature.

The advantages of this method over methods where a salt-ice mixture is used are the following: (1) The temperature of the cooling mixture is more easily controlled. (2) The material is cleaner and easier to handle. (3) The ether in the bath may be used over and over again. (4) The temperature of the ether bath can be lowered more rapidly and accurately.

L. P. LATIMER

DEPARTMENT OF HORTICULTURE,
NEW HAMPSHIRE AGRICULTURAL
EXPERIMENT STATION

SPECIAL ARTICLES

SERIES IN THE ARC SPECTRUM OF BROMINE¹

RECENTLY we have photographed the spectrum of bromine as emitted by a Geissler tube, from the ultra-violet to beyond 9300A in the near infra-red. The type of spectrum obtained depends both on the pressure of the gas within the tube and the character of the exciting discharge. When the gas at low pressure is activated by an uncondensed discharge from a high-voltage transformer the spectrum observed is predominantly that of the neutral atom, the arc spectrum.

With the new wave-length data and the wave-lengths observed by Turner² in the Schumann region, we have succeeded in working out the structure of the arc spectrum, Br I. The theoretical structure of the spectrum is similar to that of Cl I which we described in our note to SCIENCE for October 12, 1928. Turner's lines represent the combination of the lowest term $s^2p^4 \cdot 4p \ ^2P$ with the higher 2P and 4P terms coming from the electron configuration $s^2p^4 \cdot 5s$. These terms, in turn, combine with still higher terms coming from $s^2p^4 \cdot 5p$ and $s^2p^4 \cdot 6p$ to give the prominent arc lines observed in the infra-red and in the green and blue.

For the lines observed by Turner, we give the following classification.

The lines from the $5p$ and $6p$ electrons are in Rydberg sequence. We therefore use them in calculating

¹ Stuart Dunn, "The Use of 'Dry-Ice' or Solid Carbon Dioxide as a Laboratory Refrigerant," SCIENCE, March 29, 1929.

² Publication approved by the director of the Bureau of Standards, of the U. S. Department of Commerce.

² *Physical Review*, 27: 400. 1926.

λ	Int.	Term combination
1576.5	6	$^2P_2 - ^4P_3$
1540.8	6	$^2P_2 - ^4P_2$
1488.6	8	$^2P_2 - ^4P_1$
1633.6	10	$^2P_1 - ^4P_2$
1575.0	9	$^2P_1 - ^4P_1$
1449.9	3	$^2P_2 - ^2P_1$
1582.4	8	$^2P_1 - ^2P_2$
1531.9	7	$^2P_1 - ^2P_1$

the ionization potential of approximately 12.2 volts for the neutral atom. This value is checked by interpolation from the known ionization potentials of Ga, Ge and Kr.

Both Kimura³ and Hori⁴ have published observations of the complex structure of Br lines. We have used the Fabry-Perot interferometer in observing the Br spectrum and we confirm their findings as to the fine structure of numerous lines.

The details of this investigation will appear in an early number of the Bureau of Standards *Journal of Research*.

T. L. DEBRUIN
C. C. KIESS

BUREAU OF STANDARDS

ON THE BIOLOGICAL EFFECTS OF X-RAYS

THE object of this note is to record briefly the results of experiments upon *Saprolegnia ferax* carried out during the summers of 1925 and 1926 at Woods Hole. It was to be expected that with modern radiation apparatus and a knowledge of the technique as applied to *Drosophila*, *Saprolegnia* with its extremely sensitive behavior toward chemical changes and its various modes of reproduction would provide changes under the action of X-rays which could be treated statistically. Except for a possible stimulus to nuclear division in the mycelium under the influence of X-rays, extensive experimentation on the rate of growth in culture media, on the formation and liberation of zoospores, the formation of oogonia and oospores, and the movement of protoplasm, failed to produce any results which could be attributed to the action of X-rays. The amount of radiation was enormously greater than had been employed with *Drosophila*. A dosage of 50,000 volts at 2.5 m.a. for twenty minutes with a standard Coolidge tube, and the material at 12 cm from the tungsten target (in our method of recording represented by 32D),¹ had been sufficient to cause complete sterility in *Drosophila* for two days and partial sterility for ten days. Ap-

plications as high as 75,000 volts at 10 m.a. for forty-five minutes at a similar distance from the tube failed to produce any results in *Saprolegnia*. The material radiated was exposed in water in small petrie dishes, duplicate material being kept as a control. The entire life cycle of *Saprolegnia* may be carried out in from two to three days. These experiments were so conducted that samples of the radiated material could be removed for study or staining without disturbing the remaining material. It was desirable to keep a large amount of material permanently, and after experimentation with many stains it was found that gentian violet for eighteen to twenty-four hours, destained by xylol, would show chromosomes in the mycelium and the oogonia without recourse to sectioning. The chromosomes, however, are minute.

As regards *Drosophila*, the writer does not presume to say that mutations can not possibly be induced by X-rays, but a series of carefully planned and extensive experiments carried out in 1922-1924 by the writer with the cooperation of Dr. Mavor, having as one of the primary ideas the possible production of mutations by X-rays, failed entirely to produce any physical mutations which could be detected. In these experiments, with special reference to three characters of the second chromosome, black, purple and curved, under various dosages of X-rays, some 120,000 individuals were individually examined in four repeated experiments, one half the number being controls. In no case could mutations be detected in greater numbers than appeared normally in the stock. Crossovers from the "recovery period," i.e., the period of greatest percentage of crossover after X-ray treatment, were bred in numerous auxiliary experiments; in one of these the increased crossover percentage was clearly demonstrated as not inherited,² and there was no sign of mutations.

From these experiments the writer believes that X-rays tend to show the extraordinary resistance of the germ-plasm to change by experimental means. Either the organism is killed, i.e., by lethal rays during maturation divisions in the egg, resulting in complete or partial sterility; or the changes in percentage of crossing-over which accompany the recovery period disappear in the succeeding generation.

H. K. SVENSON

CAMBRIDGE, MASSACHUSETTS

CONTROL OF THE COTTON BOLL WEEVIL BY INSECT ENEMIES

PRIOR to the use of poisons for the control of the cotton boll weevil (*Anthonomus grandis* Boh.), the attention of many entomologists interested in the

² *Am. Nat.*, 58: 311-315. 1924.

³ Memoirs, College of Science, Kyoto, 4: 139. 1920.

⁴ Memoirs, College of Science, Kyoto, 9: 307. 1926.

¹ J. W. Mavor and H. K. Svenson, *Genetics*, 9: 588-608 (1924), and previous papers.