

nated eggs was given and after three weeks the experiment was terminated. Examination of the flushed intestinal contents of the fifty-seven chickens gave 133 nematodes, whereas in the scrapings of the intestinal mucosa no worms were found in any case. The technique worked admirably with both small and larger worms, the range in length being from 3.2 mm to 95.2 mm. Numbers of worms likewise caused no difficulty, for as few as one and as many as twenty-five worms were present in a chicken. Thirty-three per cent. of the birds were infested at autopsy.

In the second experiment, thirty chickens were parasitized at the age of nine weeks by giving to each bird fifty embryonated eggs of the nematode. Two weeks later the experiment was terminated with results similar to those of the first experiment, *viz.*, that from the flushed intestinal contents 186 worms were isolated, while in the scrapings of the mucosa of the same intestines not a worm was found. The percentage of infested birds in this experiment was 92; the range of individual infestations was from one to thirty-three worms, and the lengths of the worms varied from 2.1 mm to 11.5 mm. The results of these experiments give evidence that the technique is highly efficient in the removal of roundworms from the intestines of chickens.

The temperature of the water for flushing the intestine may vary many degrees and still be effective. Temperatures above 60° C. and below 35° C. caused contractions of the muscles of the intestine and thus interfered with distention and free flushing.

While the technique is especially valuable for small worms, it works equally well with larger ones, and should be readily adapted to studies on the various larval and adult nematodes, living free in the small intestine of birds and reptiles and of small and medium-sized mammals. Apparatus such as shown in Fig. 1, while desirable, is not necessary for the application of this technique. The flushing cone (Fig. 2) can be used on any hot-water faucet to which a hose

couple can be attached. It was made by threading a small brass cone on a hose couple.

JAMES E. ACKERT

L. O. NOLF

KANSAS STATE AGRICULTURAL COLLEGE

PARAMECIUM BURSARIA AS A LABORATORY DEMONSTRATION OF CYCLOSIS

THE use of *Paramecium bursaria* for demonstration of cyclosis in laboratory classes in zoology has several possible advantages, as compared with the customary use of *Nitella*. In the first place, it obviates the necessity for drawing on the plant kingdom for illustrative material. Furthermore, this ciliate may be maintained easily in laboratory cultures at any season of the year, and in addition it furnishes, with its contained *Chlorella*, an excellent example of symbiosis.

Cyclosis is unusually rapid in this species of *Paramecium*, and is readily followed under a 4 mm objective. The writers have found that especially interesting preparations may be made by staining vitally with neutral red. Clean slides, after being warmed slightly over a flame to eliminate excess moisture, are filmed with a solution of neutral red (1:1500, or more dilute) in absolute alcohol. After the film has dried a drop of culture material is added, and a cover-slip sealed in place with melted vaseline. Numerous small scattered globules are stained with neutral red, and these add to the clearness with which cyclosis may be observed. In addition, this method affords a good laboratory demonstration of the effects of vital dyes on a protozoon, while the neutral red also serves as an indicator of the pH of the inclusions. If the dye solution is dilute enough, the organisms should live for twenty-four hours or more; hence, if several laboratory sections are to be supplied, the same preparations may be used in successive laboratory periods.

R. F. NIGRELLI,

R. P. HALL

UNIVERSITY COLLEGE,

NEW YORK UNIVERSITY

SPECIAL ARTICLES

MICHIGAN PAPYRUS 620; THE INTRODUCTION OF ALGEBRAIC EQUATIONS IN GREECE

THE Egyptians some two thousand years before the Christian era set up equations of a purely algebraic type. In one type of these problems, as given in the Ahmes Papyrus,¹ an unknown number with some fractional part of it is set equal to a known number and the solution is effected by the so-called method of false position. In other problems² two

¹ T. Eric Peet, "The Rhind Mathematical Papyrus," Liverpool, 1923.

² H. Schaack-Schackenburg, "Der Berliner Papyrus 6619," *Zeitschrift für Ägyptische Sprache und Altertumskunde*, Vol. XXXVIII (1900) and Vol. XL (1902).

unknowns appear as the sides of a rectangle with known area, and their ratio being given, the two unknowns are determined. The brief translation of portions of the Moscow papyrus given by Touraëff³ also indicate that the analogous problems which appeared in Euclid's *Data* had their beginnings in ancient Egypt. The eagerly awaited complete trans-

³ B. Touraëff, "The Volume of the Truncated Pyramid in Egyptian Mathematics," "Ancient Egypt," 1917, pp. 100-102; L. C. Karpinski, "An Egyptian Mathematical Papyrus in Moscow," *SCIENCE*, 57 (1923): 528-529. In the article in *SCIENCE* I pointed out the importance of the other problems to which Touraëff refers. L. C. K.