

Much of the sand of the vicinity of Fifteen Miles Falls is so rich in mica and slate that it was considered unsuitable for the concrete of the dams. Large masses of trap and diabase were located at the northern end of the lower dam in Barnet and are being utilized in large quantities in the concrete construction. During the afternoon the geologists spent an interesting period deciphering the relations of the igneous rocks at this quarry. Four types are present, a coarse diabase, two traps of intermediate and fine grain and an aplitic granite. The pyroxenes of these rocks have been largely altered to hornblende. The coarse diabase has been further metamorphosed by the alteration of its plagioclase feldspars to albite and zoisite and the fine traps have developed a schistosity accompanied by the production of much chlorite. There was a vigorous discussion concerning the relative ages of the coarse diabase and the traps, but it was generally agreed that the alteration of these rocks had been largely influenced by the later intrusion of the aplitic granite.

Saturday evening R. J. Lougee discussed his recent study of the Connecticut valley lake stretching from Middletown, Conn., to Beecher Falls, near the Canadian boundary. The deltaic terraces bordering this

lake have been tilted since their deposition at the close of the Pleistocene from an altitude of 160 feet near Middletown, Conn., to an altitude of 657 feet at Hanover, N. H. The dam holding back this first lake broke, perhaps at the White Rocks Narrows near Middletown, lowering the lake approximately one hundred feet. A second lake surface can be traced northward from Hanover at an elevation of 565 feet to Beecher Falls at an altitude of 1,080 feet. Comparing the isobases of the Atlantic beaches along the New England coast with the lake terraces it would appear that the lake level was approximately 180 feet above the surface of the ocean at the time of its formation.

Sunday morning a number of the party visited the site of the future power plant which is being erected below the lower dam. Throughout the excursion the New England Power Association extended every courtesy to the geologists. Mr. A. C. Eaton, chief hydraulic engineer for the association, was specially helpful. All who took part in the excursion were much impressed by the magnitude of the undertaking and by the interesting geologic facts revealed in the large excavations.

W. G. FOYE,
Secretary

SCIENTIFIC APPARATUS AND LABORATORY METHODS

A USEFUL MODIFICATION OF AMANN'S MEDIUM

IN the belief that any promising method develops into more certain usefulness when generally known, I wish to add an additional point to Mr. Linder's timely note on Amann's lacto-phenol-cotton blue medium. During the past two years, since he brought it to our attention, this medium has been used with success in this laboratory by the students and myself for making temporary and permanent mounts of a great variety of materials. Its combined killing, clearing and staining action, rapid restoration of normal turgor and fixation *in situ* of easily disattached spores, are marked advantages over glycerine or glycerine jelly media. Cotton blue, however—although bringing out more structures than eosine—does not usually differentiate nuclei. To meet this difficulty I tried various nuclear stains that would mix with Amann's formula. The addition of a very little Nigrosin (water soluble), either aqueous, or in saturated picric acid as described by Curtis and Colley in the *American Journal of Botany*, for February, 1915 (with subsequent filtration or settling if necessary), was found to differentiate nuclei as well as other cell structures in a considerable number of representative fungi and algae.

Moreover, with sections of host plants invaded by mycelium of such fungi as the downy mildews, a pro-

longed stay in the medium yields a differentiation of the host and parasite like that resulting from the more complicated treatment with lacto-phenol-cotton blue and safranin described by Lepik in *Phytopathology* for October, 1928.


The medium lends itself well to making permanent mounts either by the double cover-glass method of Diehl (*SCIENCE*, p. 276, 1929), or by the older but quite reliable method described by Bullard in the *Transactions* of the American Microscopical Society in April, 1921, provided one exercises the same scrupulous care to insure perfect sealing (preferably with King's cement) that is necessary when mounting in glycerine. When making permanent mounts in very damp weather, as during a tropical rainy season, it is advisable, before sealing, to concentrate the medium on the slides in a desiccator.

WM. H. WESTON, JR.
LABORATORIES OF CRYPTOGAMIC BOTANY,
HARVARD UNIVERSITY

A CIRCUIT BREAKER FOR WATER-COOLED X-RAY TUBES

A RECENT issue of the *Journal of the Optical Society*¹ contains the description of a device for pro-

¹ L. W. McKeehan and L. M. Kirkpatrick: *Journal Opt. Soc.*, April, 1929.



protecting water-cooled X-ray tubes from failures of the water line, the operation of which depends on the outflow from the tube. A device actuated by the water flowing out of the X-ray tube was constructed by the author and used at Cornell University in 1926-1927. Inasmuch as the arrangement used is much simpler in construction than the one described in the journal, and quite efficient, it seemed worth while to give its description here.

A beaker having three tubes sealed to it is placed in the path of the water from the X-ray tube, the water entering the beaker through tube A (see

figure). The diameter of tube B is slightly smaller than that of A, and can not carry the outflow. The water in the beaker rises to the level of the large tube C raising a cork plug with a mercury cup in the center. In this position two copper electrodes passing through the cover of the beaker dip into the mercury and close the primary circuit of the transformer. It is, therefore, impossible to operate the tube without turning the water on. In case of failure of the water system the level of the water in the beaker drops and with it the plug with the mercury cup, thereby breaking the primary circuit. Temporary fluctuations of the water pressure will not affect the circuit, as it takes a few seconds for the water level to drop so low as to break the circuit owing to the small diameter of tube B. It is necessary to make the wooden plug waterproof and to prevent its swelling. This is accomplished by keeping the plug in boiling paraffin for about an hour. The copper electrodes are sealed into glass tubes passing through a rubber stopper in the cover. The level of the electrodes is easily adjusted by pushing the glass tubes in or out of the stopper.

A. A. BLESS

UNIVERSITY OF FLORIDA

SPECIAL ARTICLES

VITAMIN B DEFICIENCY IN NURSING YOUNG RATS AND LEARNING ABILITY¹

INNUMERABLE experimental studies have established the causal relationship between deficient diets and various forms of physical deformities. Howe² has produced experimentally in monkeys the most striking cases of tooth and nasal deformities resembling those often seen in human beings simply by placing the young and rapidly growing animals on a diet partially deficient in vitamins.

Similarly, diets deficient in vitamin B experimentally produce polyneuritis in young adult animals and play an important rôle in the etiology of pellagra and beriberi of human beings. These experiments show that the relation also exists between diet and the nervous system. However, thus far there has been practically no adequate experimental study on the relation of diet to learning ability.

It was the purpose of our investigation to determine the effect of various deficient diets upon the learning ability of the first, second and succeeding generations

of rats of known heritage, and to correlate these findings with the anatomical and chemical changes in their nervous system.

The present report, however, deals only with the number of trials required to learn a standard maze by seventy-one rats of the first generation, some of which were depleted of vitamin B during their nursing period. Other data will be reported in forthcoming articles.

As it was our purpose to deplete the animals during their nursing period, we could succeed only by depleting the nursing mothers. The technique we used in giving the new-born pups a diet at first deficient and later free of vitamin B is that described by Sure,³ who has demonstrated that the quantity of vitamin B in the milk of the nursing mother rat is proportional to the vitamin B content of the mother's diet.

Forty-three animals were thus depleted on this diet until they were strong enough to wean. The degree of depletion was, of course, subject to variation; some of the pups were depleted to a point so near death that their lives were saved with great difficulty, while others appeared quite normal all the way through except for retardation of the rate of growth. Twenty-eight other animals were used as controls, their

³ Barnett Sure, "Vitamin Requirements for Nursing Young, III," *The Journal of Nutrition*, 1: 139.

¹ From the Otho S. A. Sprague Memorial Institute and the departments of pathology and psychology, University of Chicago.

² Percy Howe, "Percy Howe's Letters," *The Dental Digest*.