

istic mechanism. It illustrates beautifully the dependence of space on time in our estimations of visual, tactile, kinesthetic and auditory space. In order to produce the *Tau* effect we may say, in general, vary the time interval in the opposite direction to the space interval and the latter will be distorted accordingly. So easy is it to demonstrate the *Tau* effect that it can be used as a parlor trick or game.

It may be thought that we are here dealing with a lightly dispelled illusion or error in judgment in which the subject unwittingly is judging the time intervals instead of the spatial distance between the spots touched. Nothing could be farther from the truth, for even when the subject knows what the effect consists in and is due to, if we vary our conditions by reversing the spatio-temporal relations, the subject will be wholly lost as to whether or not the spatial intervals are really equal or different and in what sense they differ. We have here, I believe, a *bona fide* example of the interdependence of time and space. They are so intimately related psychologically, as well as physically, that by varying them in opposite sense it is possible to demonstrate directly to an observer the distortions in space which relativists have told us about. It is interesting to note that whereas it is doubtful if the physicist can ever hope to do more than make relatively an intelligible abstraction to the layman, the psychologist by this simple experiment can directly demonstrate what the interdependence of time and space means in direct experience.

Several factors influencing the *Tau* effect which should be noted if one is to get it at its best are the following: (1) care should be taken to touch all the spots equally so that no one stands out more than another; (2) the greater (or less) the spatial distance between the second and third stimuli as compared with the first and second, the less (or greater) must the time interval be between the latter as compared with the former, if the effect is to appear; (3) the optimal effect is limited by the actual spatio-temporal intervals used: we have found that distances as great as 80 mm on the back of the arm and times as long as 1 second may be used. There is practically no lower time limit, although the second temporal interval should not bear a greater ratio to the first than 3 or 4 to 1.

A fuller, quantitative account will appear in one of the psychological journals.

HARRY HELSON

BRYN MAWR COLLEGE

ON THE AGE OF THE NEW ALBANY SHALE

IN an interesting article on "Petrified Wood in the New Albany Shale," published in *SCIENCE* for De-

cember 13, 1929, Chester A. Arnold described an occurrence of fossil wood in the upper part of the New Albany shale in Scott County, Indiana, which he referred to the genus *Callixylon*. The concluding paragraph of his contribution is as follows:

Although the wood is widely scattered, it appears to occur mostly near the top of the New Albany formation. While formerly considered as belonging to the upper Devonian and of the same age as the Genesee shale of New York, the New Albany shale is now viewed by some competent authorities as being, at least in part, of lower Mississippian age. This would place the Indiana wood in the Mississippian, and thus extend the range of *Callixylon* from the Devonian up into the Carboniferous. However, there is no record of its occurrence any higher than this basal member.

It seems to the present writer that Mr. Arnold is not justified in his conclusion quoted above. The New Albany shale as it occurs in the type locality in the vicinity of New Albany, Indiana, is a definite formation. Other black shales present in eastern Ohio, Kentucky and Tennessee, east of the Cincinnati anticline, were until a few years ago thought to be the equivalent of the New Albany shale at New Albany, Indiana. In recent years a part of the black shale in eastern Ohio, Kentucky and Tennessee has been shown to be of early Mississippian age. However, this does not prove that any part of the New Albany shale at New Albany, Indiana, is younger than the Upper Devonian, but rather that such part of the black shale of eastern Ohio and Kentucky as is now known to be of early Mississippian age is younger than any part of the typical New Albany shale at New Albany, Indiana. So far as known to the present writer, no one has ever shown that any part of the New Albany shale, as it is developed in the type locality near New Albany or farther north in Indiana or south in Kentucky, west of the Cincinnati anticline, is younger than upper Devonian age. Therefore, it seems that the more logical conclusion to be drawn from the occurrence of *Callixylon* in the upper part of the New Albany shale in Scott County, Indiana, would be that this shale is of upper Devonian age, because the genus of fossil wood that occurs in it has never been found in strata younger than the Devonian.

T. E. SAVAGE

UNIVERSITY OF ILLINOIS

FIREFLIES FLASHING IN UNISON

SEVERAL times in recent years correspondents of *SCIENCE* have directed attention to the synchronous flashing of a swarm of fireflies or other insects, as at page 132 in the issue for January 31, 1930.