

ciates it would seem not only logical but inevitable that bacteria in all stages of their developmental cycles should be found in nature. It is of value to demonstrate this fact because of its many important implications, biological and practical. It is important, also, in order to remove the many significant observations of Hadley and previous investigators from the shallow objection of being artificially induced laboratory "involution."s.

Following the methods of Hadley, and also with slight variations therefrom, we have demonstrated the presence of filterable microorganisms in soils, decomposing manure, hay infusions, fresh human feces and milk. We have followed the transformation of some of the "g" types (Hadley) to the point where they would be recognized as "bacteria" in the usual sense. It should hardly be necessary to state that all filtrates have been controlled and tested for "sterility" by the conventional methods. Berkefeld v and n filters have been used with the same positive results. Growths obtained from the filtrates have been refiltered and the "g" types again developed.

JAMES M. SHERMAN
CLAIR E. SAFFORD

CORNELL UNIVERSITY

POSITIVE GAS AND WATER PRESSURE IN OAKS

EVIDENCE of positive gas and water pressure in forest trees was observed by the writers during the late summer and early fall of 1930, in western North Carolina and northern Georgia hardwood forests.

In connection with a growth study carried on by the Appalachian Forest Experiment Station, numerous forest trees were drilled with the increment borer. This tool consists of a hollow steel bit, turned like an auger into the bole of a tree (usually at breast height). The hollow bit will then extract a solid core of wood somewhat smaller in diameter than a pencil, extending from bark to pith. A sudden reverse twist of the handle frees the core from solid wood at the inner end of the bit. Annual rings on the core can be counted for determining age or measuring growth along that particular radius.

It was the writers' observation that frequently after the bit had been inserted two inches or more, scarlet oaks (*Quercus coccinea*) would emit enough liquid to cause a dripping from the outer end of the borer. Often there was an accompanying hissing sound as though gas were escaping, though not with any great pressure.

One 14-inch apparently sound scarlet oak was encountered which had sap pressure enough to eject the 3-inch core with considerable force and follow it with a stream of liquid which was thrown 3 to 4 feet

from the base of the tree. This liquid had the characteristic ill-smelling odor of scarlet oak sap. The occurrence seems more remarkable considering the very dry season preceding it. At that time, late summer, the rainfall deficiency for the calendar year was 12.73 inches related to a normal of 40.28 inches.

The only two species in which inflammable gas was found were the chestnut oak (*Quercus montana*) and the white oak (*Quercus alba*). In some of these trees, which ranged from 8 to 16 inches, diameter breast high, the positive gas pressure was sufficient to blow the core out of the hollow bit with considerable force. The gas was frequently lighted and would shoot a blue flame, sometimes 1½ to 2½ feet long, extending horizontally near the source but gradually curving upward to its tip. The flame would burn steadily for thirty seconds or so, then gradually lose force and become smaller. It was usually snuffed out soon after lighting to preserve the temper of the borer. In all trees issuing inflammable gas the heartwood was unsound, apparently affected by a dry rot. Curiously enough failing pressure could occasionally be revived by turning the borer into the tree a little farther; sometimes the issuance of gas was stopped completely by turning it too far. Because of the fact that the flame was blue it suggested a gas similar to, if not, methane, which is known to be a product in the decomposition of cellulose.

The writers have not observed positive gas or sap pressure in any other species or at any other season of the year than indicated above. Wood,¹ however, observed inflammable gas in white oak and red oak (*Quercus borealis*) in West Virginia during the fall of 1927. Such phenomena as have been observed may be of interest to investigators of the water and gas systems of trees.

C. A. ABELL
C. R. HURSH

APPALACHIAN FOREST EXPERIMENT STATION,
ASHEVILLE, NORTH CAROLINA

ANALYSES OF THE BLOOD OF IDIOTS

IN the issue of SCIENCE for March 20, page 316, there appears a note headed "Biochemistry in Relation to Intelligence." In this note the writer, H. D. Powers, claims to have found an abnormal amount of inorganic phosphate in the blood plasma of idiots. In results which we have obtained from analyses of blood samples taken from idiots of classified mentality we have been unable to obtain such results. Inorganic phosphorus in blood plasma samples from twenty-five idiots has been found to be within the normal limits. Our results range from 3.1 mgs to 5.0 mgs per 100 cc of blood, with an average of 4.0

¹ L. M. Wood, "Gas from Trees," Service Bulletin, U. S. Forest Service, Washington, D. C. December 1, 1930.