

colloidal suspension in the soil solution. In addition, iron and aluminum are found adsorbed by this organic material to the extent of about 40 p.p.m. and 20 p.p.m. of the soil, respectively. These soils and soil solutions with this treatment remain acid between values of pH 6.1 and 6.7.

Iron toxicity, due to increase in soluble iron after liming, has been claimed for the Louisiana soils, but the evidence that we possess indicates that the presence of the soluble sesquioxides in small amounts after sodium alkali treatment does not inhibit production from these Quebec soils but permits positive increases.

G. T. SHAW
R. R. McKIBBIN

MACDONALD COLLEGE

THE ATTRACTION OF SPHERES

It is appropriate that some warning should be given the inexperienced reader of Mr. Thaddeus Merriman's article in *SCIENCE* for April 14 (p. 371). His method of calculating the attraction of a homogeneous sphere upon a neighboring body involves assumptions inconsistent with the generally accepted methods of the integral calculus, as well as with the Newtonian law

of gravitation. Newton's beautiful geometrical proof that the net attraction of such a sphere is exactly the same as if all the mass was concentrated at its center is, of course, fully confirmed by the rigorous analytical discussion which may be found in any standard treatise on mechanics, and so need not be reproduced here.

Mr. Merriman's proposed substitute law is inconsistent not only with theory but with the fact. For a small sphere near the earth's surface, it reduces to $F = G \frac{M \cos \alpha}{d^2}$, when M is the earth's mass, d its radius, and α the angle between a line drawn to the earth's center and one tangent to a circle of radius 0.424 times the earth's. The factor $\cos \alpha$, which is omitted in Newton's equation, has the value 0.911 for a point on the earth's surface. Without it the attraction at the surface is known to be in complete agreement with that inferred from the orbital motion of the moon. To introduce it destroys the agreement, which provided Newton with the first conclusive test of his theory.

HENRY NORRIS RUSSELL

PRINCETON
APRIL 21, 1933

SCIENTIFIC APPARATUS AND LABORATORY METHODS

A TWO-FIELD STROBOSCOPE


IN stroboscopic observation of vibrating strings for measurement of frequency, no use seems to have been made of a two-field illumination. The two fields are contiguous and are alternately illuminated. The two-field method removes some ambiguities of the one-field method. But more important is the fact that a feeble resonance of the string to a tuning-fork or other source is more easily detected.

For the laboratory, the simplest device is the type of Neon bulb built for a. c. lighting circuits and having semi-cylindrical electrodes. Such a bulb can be held behind a string in such a manner that half of each electrode is visible. The two electrodes light up in alternation, the flash frequency of either side being equal to the frequency of the current.

For lecture demonstration, one can take a siren disk with an even number of holes. Alternate holes are half covered on the half nearest the center of the disk; the remaining holes are half covered on the side next the edge of the disk. The disk is set as closely as possible to the string, with the center of the disk level with the string. When a hole is level with the string, a strong beam of light is made convergent by means of a condenser, and is arranged to just cover either type of hole. On the other side, an objective focusses the string on a screen. When the disk is rotated and the string made to vibrate, a

two-field stroboscopic view of the string is obtained. (Many other methods may be used.)

To understand the patterns seen, the procedure is to construct in advance the patterns to be expected. To do this, imagine a stroboscope disk with a single spot; rotate this at S revolutions per second, S being the string frequency; and illuminate the disk by F flashes per second. When the pattern has been worked out, project the spots of the pattern on a vertical line in the plane of the disk. Then repeat the process for a series of flashes beginning one half flash-period later. The projections of this second pattern gives the string pattern in the other field. The figure herewith illustrates the case when the flash frequency is $3/5$ of the string frequency. Small circles give the first disk pattern, and crosses give the second.



Disk Pattern

Two-field Pattern

Figure illustrating the construction of the two-field pattern for $F = S.3/5$.

A few simple rules may be given here. Assume that $F = S.f/s$.

For the disk patterns:

(1) Sf/s flashes per second produce a pattern of f spots, independently of the value of s .

(2) If s is even, the crosses coincide with the circles.

(3) If s is odd, the crosses alternate with the circles.

For the two-field string patterns:

(1) The number of lines seen in either field is equal to f ; when two lines merge into one, that line must be counted twice.

(2) If s is even, the two fields are identical, and appear as a single field.

(3) If s is odd, the two fields are similar but not identical.

It is rarely the case that f/s is a simple ratio. Then, as in Lissajou curves, one sees changing patterns. These can be worked out by shifting the initial position of the disk in the construction shown above. It is just this fact that makes the two-field method so convenient. Assume that a string has been tuned to the frequency F , for which we obtain just one line in either field. If the string starts to resonate to some source, one can usually observe two lines moving in opposite directions. The writer has used a $\times 25$ microscope to observe such resonances.

The changing patterns just mentioned can be worked out by a slow rotation of the disk patterns, as stated. From this it will be understood that a complete cycle of change means that the disk has moved by an angle equal to the angle between two spots of the same set. Or in the two-field patterns, we have made a complete cycle of changes when we return to some pattern in the same way. Then, if the pattern corresponds to a ratio f/s , and there are R complete repetitions per second,

$$\frac{F}{f} - \frac{S}{s} = \frac{\pm R}{fs}$$

Although, for simplicity, we have spoken of strings,

the two-field method is adaptable in many ways to the observation of other bodies, *e.g.*, the writer used a Neon bulb in the selection of steel springs which were to vibrate at a frequency of 240.

PAUL F. GAHR

WELLS COLLEGE

REGULATING THE AIR SUPPLY OF MICRO BURNERS

Most of the micro burners obtainable from supply houses have an inflexible and very often an inefficient adjustment of the gas-air mixture. The usual practise with laboratory workers has been to test a number of burners and then select the best of the lot, without attempting adjustment. Obviously, this is an awkward and somewhat unsatisfactory solution of the problem.

A method, almost as convenient as that used for the Bunsen burner, has been found for regulating the air supply of micro burners. The small burner is connected to the gas line and lit. While the flame is burning, twist with a pair of pliers the metal gas connection which projects from the base, loosening or tightening until the proper type of flame is obtained.

Occasionally a burner will be found which requires further treatment. In these cases the gas connection is unscrewed and, using a small three-cornered file, two flat channels about 5 mm wide and 6 or 8 mm long are filed on each side of the tube. These shallow channels start from the middle of the tube and slant upward, meeting at the pin hole gas outlet on top. The connection is replaced frequently while filing and the gas-air mixture tested as described above, in order to prevent spoiling the burner by too much filing. If necessary, the small gas outlet can be made larger by reaming with a pin. However, it is usually found that the opening is already too large. This can be remedied by filing the sides of the hole rather thin and then pressing the edges inward until the desired opening is obtained.

CORNELIUS A. DALY

MEDICAL DEPARTMENT, UNION UNIVERSITY
ALBANY MEDICAL COLLEGE

SPECIAL ARTICLES

ACCELERATED, EXPERIMENTAL POLIOMYELITIS IN NASALLY INSTILLED MONKEYS

IN a previous communication,¹ attention was directed to the accelerating effect of a second virus injection, after an interval of from 7 to 10 days, in provoking experimental poliomyelitis in *Macacus rhesus*. The earlier report dealt with the phenomenon from

¹ S. Flexner, "Accelerated Infection in Experimental Poliomyelitis," *SCIENCE*, 1931: 74, 520.

the point of view of establishing human strains of virus in monkeys. Since that report was made, additional studies of the subject of accelerated infection have been undertaken.

The first test, made by injecting the human virus intracerebrally and intraperitoneally, were soon followed by the injection of sub-infective doses of monkey virus intracerebrally alone. The results showed that acceleration does not depend on the