

curate title would have been "Sahuaro Susceptible to Crown Gall."

J. G. BROWN
M. M. EVANS

UNIVERSITY OF ARIZONA

THE SPECIFICITY OF FROG FLUKES

THERE are few animals more subject to parasitism than are frogs, and the ease with which these animals are collected and kept in captivity makes them ideal for the study of parasites of many kinds. In recent years the life-histories of a few trematodes which infect the frog have been worked out, but unfortunately very little attention has been given to the interesting problem of the specificity of these flukes for their hosts. The experimental approach to this problem is attended with some difficulty, since the mollusk, insect and frog which are involved as hosts must be laboratory-raised, and the investigator has usually been content if he has been fortunate enough to have the necessary hosts to demonstrate the life-history, without trying to find out just how specific the parasite may be for its hosts.

Considerable light can be thrown on this problem if frogs are carefully examined both within their natural range and in regions where they have been successfully introduced. A good example of the latter is to be found in the Hawaiian Islands and in the Sacramento Valley in California. In the former there are no native frogs, but such species as *Rana rugosa* from the Orient and *Rana catesbeiana* from the eastern part of the United States are well established. In a correspondence with Dr. C. H. Edmondson, of the University of Hawaii, I am informed that several hundreds of the American species have been used in his laboratories during the past few years, and not a single fluke has been found in them. Apparently the introduced bullfrog lost its flukes because of the lack of suitable mollusks or insects which serve as hosts. However, trematodes from domestic mammals which were reared on the islands have been found.

The frog "plant" in the vicinity of Gridley, California, was made several years ago, and at present the bullfrog has spread over most of the rice fields in that part of the state. They are very common and are even penetrating the near-by mountain streams, where they are living side by side with two California species, *Rana aurora* and *Rana boylei*. Although the two latter frogs have been found to harbor eight species of trematodes, none has ever been found in the introduced *Rana catesbeiana*. We use this frog in the elementary classes in biology and physiology, and I have personally examined the intestine, bladder and lungs of a large number but without the trace of a trematode. This is of especial interest, because two species of snails of the genus *Gyraulus* and at least

seven species of dragonflies belonging to the genus *Sympetrum* are found in California, and species of these genera serve as hosts for certain frog lung flukes in Michigan and for other frog lung flukes in California. Both of the above-mentioned genera are represented in this locality.

Young bullfrogs and a single eastern leopard frog, *Rana pipiens*, were fed the metacercaria of *Ostiolium oxyorchis*, a common lung fluke in *Rana aurora*, but without results.

These observations indicate that there is a rigid specificity for the hosts by frog flukes and that the introduction of frogs into new regions may result in the loss of flukes because of the lack of suitable hosts. If this specificity should prove experimentally to be as rigid as is indicated, it might be used as a means of determining the identity or assisting in the differentiation of species of mollusks concerning which there is at present a difference of opinion. Whether this freedom from these parasites plays any appreciable part in the success of the adventitious bullfrogs over the indigenous frogs can not be stated, because it is doubtful whether the frog flukes actually harm their hosts seriously, but in the Gridley rice fields the bullfrogs outnumber all other native species ten to one. However, many other factors might cause this difference in numbers.

LLOYD G. INGLES

STATE TEACHERS COLLEGE,
CHICO, CALIFORNIA

POST-BOLSON FAULTING IN NEW MEXICO

DURING a recent trip into the San Andreas Mountains, the writers noted two low but interesting scarps in an area of bolson deposits. These were located at 25.2 miles and at 29.1 miles, respectively, west of Tularosa, on New Mexico state highway number 52. Unfortunately, time did not permit tracing these features either way from the highway, but a few facts of some importance were observed.

Both escarpments are moderately straight, and about the same height, approximately fifteen feet. Both face east, and neither one, at least within sight of the highway, shows any indication of an opposite-facing cliff, as would be the case if they were the banks of arroyos.

Even more interesting is the fact that each scarp shows a gentle but unmistakable back slope to the west, that of the easternmost being about fourteen feet per mile and extending for 3.9 miles; that of the westernmost thirty feet per mile for 1.0 mile. Both are considerably west of the center of the Tularosa basin. As a result of the back slopes, the surface of the basin floor actually declines westward from the scarps towards the adjacent mountains, just the reverse of what would be expected, were the intervening