

cal time they are half way from their ancestors to mammals.

HOWARD AYERS

UNIVERSITY OF CINCINNATI

A NEMATODE PARASITE IN MYXINIDS

IN studying *Bdellostoma*, the nematode parasite was found in the large subdermal blood sinus. It was usually located in the dorsal part of the sinus in the head region, and in the preserved specimens the worms were found closely coiled together and interlaced with the strong tendons connecting the skin to the trunk in the median dorsal line, also sometimes intertwined among the connective tissue strands which fasten the skin of the ventral and ventro-lateral part of the head to the muscles. The myxinoids appear not to harbor many parasites, which fact may be related to their long-continued existence. Specimens of this nematode were sent to G. Steiner, senior nematologist, of the U. S. Department of Agriculture in 1928. Steiner named it *Tetanonema strongylurus*, new genus, new species. At that time nothing was known of the method of propagation. I have recently found, however, that the microscopic embryos of the parasite leave the body of the parent *Bdellostoma* in the eggs and are to be found in the growing embryo before hatching as well as after hatching, coiled in the tissues of the gills. Whether they infest other parts of the embryo *Bdellostoma* is yet unknown. So far as I know this is the sole animal parasite occurring in the myxinoids, and no plant parasite is known to attach itself to or live within the body of these animals. How the filaria of this nematode enter the *Bdellostoma* egg and at what stage of the growth of the egg is not known. They enter before the micropyle is fully formed. None of the adult nematodes have been found in the gills of the adult *Bdellostoma*, but colonies of the worms numbering up to one hundred or more occur in the subdermal blood sinus.

HOWARD AYERS

UNIVERSITY OF CINCINNATI

THE THICKNESS OF THE GLACIAL DEPOSITS IN OHIO

AN examination of nearly 2,800 well records, obtained from the files of four oil companies which operate in Ohio, has revealed some interesting facts concerning the topography of Ohio in preglacial and interglacial time. It was possible to obtain figures for the thickness of the glacial drift over an area of approximately 4,637 square miles in north central Ohio, in Cuyahoga, Summit, Lorain, Medina, Ashland, Wayne, Richland, Holmes, Tuscarawas, Portage and Coshocton counties. As expected, it was found that the thickness of the glacial drift is much greater in the preglacial and interglacial valleys, the majority of which coincide, in trend, with the present valleys.

On the buried uplands between the valleys, the thickness of the drift varies, and appears to be thinnest along Lake Erie, where in places it is not much above 20 feet in thickness. Fourteen quadrangles show average thicknesses over the buried uplands varying from 37 to 67 feet. For the entire area, the thickness over the buried uplands averages 51 feet.

The depth of the glacial drift in the buried valleys averages from 155 feet in the Wellington quadrangle to 380 feet in the Cleveland quadrangle. The average thickness of the drift for all the buried valleys in the entire area is 205 feet. The maximum thicknesses in feet for the fourteen quadrangles range from 192 feet for the least, to 763 feet for the greatest. Thicknesses of 300 feet and over are common throughout the area. The thickness of the drift, over all, taking the average for buried valleys and uplands in the entire area, is 95.7 feet. This figure was derived by taking the average of all the well records, a total of 2,782. The Oberlin and Berea quadrangles, located farthest north, along Lake Erie, have the least thickness, 54 and 43 feet, respectively, whereas the area farther south represented by the Medina, Akron, Ashland, West Salem, Wooster, Massillon, Canton, Perrysville, Loudonville and Millersburg quadrangles have thicknesses of 85, 111, 90, 81, 80, 154, 97, 88, 90 and 94 feet, respectively. The greater thicknesses are explained by the facts that, where they occur, there are either deep buried valleys or morainic belts.

The greatest thickness in the entire region and possibly in the State of Ohio, 763 feet, is located south of Cleveland, in Newburg township, about two miles directly east of the village of Independence. Here a north-south buried valley marks the course of the preglacial or interglacial Cuyahoga River. Other thicknesses in this valley are 605, 560, 519 and 500 feet and many more greater than 400 feet. The elevation of the bed-rock which marks the buried surface, at the point where the thickness of the drift is 763 feet, is 13 feet below sea-level and 586 feet below the present level of Lake Erie. The bottom of this valley stands 800 to 900 feet below the buried upland, as indicated by well records in Newburg and Independence townships. It appears to have been a narrow gorge in the youthful or early mature stage in the cycle of erosion and may have stood much higher than at present. The buried valleys throughout the area are, with few exceptions, narrow at the bottom, but their upper portions are broad, with rather gently sloping uplands. It is probable that the valley of the old Cuyahoga, and other buried valleys in the region, are not the result of stream erosion alone, but have been deepened and otherwise changed by glacial erosion.

KARL VER STEEG

COLLEGE OF WOOSTER