elements are more easily disturbed and broken apart than is the case among more complex organisms. The study of fossils shows the extraordinary permanency of structures in the higher animals and plants. It may well be that the cytoplasm, in addition to its other functions, serves to protect the chromosomes from such disintegrating influences as are described by bacteriologists. The phenomena of crossing over show how readily a part of a chromosome becomes attached to another, and by analogy it is easy to understand why bacteria so readily "take on" free genes. Dr. Manwaring says: "Transmissible bacterial genes are apparently widely distributed in nature, being found, for example, in almost any contaminated surface water." The "polyvalent" genes may really be aggregates of two or more. It is conceivable that we may return to a sort of Darwinian pangenesis, and postulate the existence of many kinds of "free genes," which are ready to unite with the organized systems of genes when they have a chance. Some bold experimenter, perhaps using sperm cells on account of the absence of the thick cytoplasmic covering, may one of these days succeed in adding genes to the germinal elements of the higher organisms.

What we now want to know is whether the "dissociated genes" arise *de novo* from inorganic or non-living sources or whether they are always the result of the breaking up of living systems. Just as the inorganic letters of the alphabet c, a, t, when united give us the organic cat, so it is conceivable that the genes owe their significance as vital units to their being parts of a system, and not to any special "vital" properties of their own.²

In any case, we have plenty of evidence to show the extraordinary stability of genes in nature, their persistence during many millions of years, under all sorts of diverse conditions. This stability may in a sense be a product of natural selection, since it is essential for the processes of evolution and adaptation. Nature can not build on a quicksand. It does not seem probable that the phenomena described by Dr. Manwaring can be ascribed to perpetual or very frequent gene mutations, or to specific changes in genes induced by particular environmental factors. According to this view, bacterial genes may be about as stable as others, and there is no "Lamarckian world of bacteriology."

UNIVERSITY OF COLORADO

T. D. A. COCKERELL

DARWIN'S VIEW OF HEREDITY

IT seems that in the interest of modernity we ought to demonstrate the fallacies of our predecessors. One

² For a discussion of the gene as the unit of life, see Hurst, "The Mechanism of Creative Evolution," Chap. XVII, 1932. of the favorite methods adopted for this end is to schematize the theories of earlier workers and then show how modern advances have shown these schemes to be untenable. It seems to me that the time has come, however, when text-book writers ought to check over more thoroughly the written works of the author whose theory is being criticized. The particular instance of this which is rapidly becoming my private grouch is the apparent wide-spread belief that Darwin believed all variations to be inheritable, and thus grist for the natural selection mill. In a fairly recent textbook, for instance, there occurs the statement, "Darwin believed that all differences among individuals were hereditary."

I would like to call attention to some quotations from Volume 2 of Darwin's "Variation of Animals and Plants under Domestication." In the first paragraph of chapter 12: "It is obvious that a variation which is not inherited throws no light on the derivation of species, nor is of service to man, except in the case of perennial plants, which can be propagated by buds." Again, about two thirds of the way through the same chapter: "When a new peculiarity first appears, we can never predict whether it will be inherited."

It is true that he also stated, in Chapter 28, that "we are led to conclude that species have generally originated by the natural selection, not of abrupt modifications, but of extremely slight differences." This has frequently been stressed as a difference between his and more recent theories which stress mutations. Since, however, mutations no longer signify large abrupt changes alone, but simply heritable variations, however slight, and since we have found that the larger share of these are very slight alterations, it seems something of a quibble to say that, since Darwin did not believe that "sports" were especially significant in evolution—a view which modern geneticists would subscribe to if "sports" mean such large modifications as they did in Darwin's day, e.g., moss roses, hornless cattle, etc.—his view differs so radically from such a view as is, for instance, incorporated in Morgan's "Scientific Basis of Evolution."

With the hope that this protest will lead an occasional biologist either to glance through for the first time or to review once more one of Darwin's most significant contributions to scientific literature, I submit it.

GEORGE M. ROBERTSON

DARTMOUTH COLLEGE

TERRACES IN THE SUSQUEHANNA VALLEY BELOW HARRISBURG, PENNSYLVANIA

DURING the summer of 1931 the writer made a study of river terraces in the Susquehanna Valley