

tween the coarse clastic formations and the fine sea deposits are found most of the great coal formations which supply the markets of the world with this type of fuel.

It happens that there is abundant evidence of the existence of huge glaciers in the southern hemisphere during the very times when these curious alternations of deposits were being formed.¹ A relation between these continental glaciers and the sedimentary cycles has been proposed recently by the writers.² It is well known that the growth of these various great glaciers of the Pleistocene ice age was accompanied by raising and lowering of the sea level hundreds of feet and by world-wide climatic alternations. Many geologists believe the Permo-Carboniferous glaciers were more extensive than those of the recent ice age, and certainly the glacial conditions persisted over a much greater span of time than the recent ice age. They were probably accompanied by the same waxing and waning of the ice masses. The resulting changes of sea level and of world climate may have caused the alternating types of sediment which go to make up the coal series.

To see how these conditions would have operated, let us start with the formation of the first great glaciers. As the glaciers grew, more and more water would have been withdrawn from the shallow seas which were previously spread over much of the continental surfaces. As the seas withdrew the climate would have grown colder as an accompaniment of the advancing glacial conditions, and greater aridity may have resulted from the decrease of evaporating surfaces from which the atmosphere derived its moisture. The net result would have been the killing of vegetation on the slopes of the mountainous lands which were supplying the inland basins with sediment. Since most semi-arid regions now have periodic heavy rains, it is likely that these barren mountain slopes would have been subjected to violent storms with accelerated erosion, and coarse debris would have been spread as great fans over the emergent lowlands. At this time and in subsequent glacial epochs the coarse continental sediments of the cycles would have accumulated.

With the warming of the climate which led to the melting of the glaciers and the rise in sea level, the slopes would have again become cloaked with vegetation and the streams would have ceased contributing the coarse sediment which had been due to rapid slope wash. As the ground water was raised on the plains and profuse vegetation began to grow, swampy conditions would have developed and in these swamps the

peat which later formed coal could have accumulated. The upward growth of the great tangles of vegetation may have held back the advancing seas till large accumulations had been formed in many places.

When the rising seas finally overwhelmed the swamps the marine phases of the cycle set in and an accumulation of muddy sediments formed a cover over the peat beds causing the peat to turn gradually into coal. After the sea level had risen sufficiently to drown the lower courses of the land valleys the muds which had been washed into the open seas would be deposited in the resulting bays and the seas would have cleared and allowed the deposition of limestone. Return of the glacial climates would have led to a repetition of the cycle and of the special sequence of formations already described.

The explanation which has been outlined does not attempt to account for all the phenomena observed in connection with these sedimentary formations. The sequence of formations has been generalized and is actually more complex due to varying local conditions, to the distance of different areas from the sources of sediment, and perhaps to the oscillatory character of the advancing and retreating seas. The explanation has been proposed as a substitute to the hypothesis that the cycles were due to alternate uplift and sinking of the basins of sedimentation and of a much greater, but contemporaneous, uplift and sinking of the source areas. There are mechanical difficulties in such an explanation, especially in view of the wide-spread recognition of the cyclic phenomena during these periods. On the other hand, earth movements probably had an important effect on the sedimentation, but it seems likely that these movements were largely of the order of slow progressive sinking of the basins in which the sediments were accumulating and slow rising of the mountainous tracts which were the source areas of the sediment.

FRANCIS P. SHEPARD
HAROLD R. WANLESS

UNIVERSITY OF ILLINOIS

BOOKS RECEIVED

- Annales de L'Acfas. Vol. 1, 1935.* Pp. 177. Association Canadienne-Française Pour L'Avancement des Sciences, Montreal.
- CAMERON, GLADYS. *Essentials of Tissue Culture Technique.* Pp. xvi + 134. Illustrated by C. G. Grand. Farrar and Rinehart. \$3.00.
- MAGIE, WILLIAM F. *A Source Book in Physics.* Pp. xiv + 620. 111 figures. McGraw-Hill. \$5.00.
- ROULE, LOUIS. *Fishes and Their Ways of Life.* Pp. viii + 312. Translated by Conrad Elphinstone. Norton. \$3.75.
- STORMZAND, M. J. and ROBERT H. LEWIS. *New Methods in the Social Studies.* Pp. ix + 223. Farrar and Rinehart. \$1.75.
- WILSON, CARL and JULIA M. HABER. *An Introduction to Plant Life.* Pp. xiv + 493, 315 figures. Holt. \$3.00.

¹ A. L. Du Toit, Abst. XVI, International Geol. Cong., p. 27.

² Harold R. Wanless and Francis P. Shepard, "Sea Level and Climatic Changes Related to Late Paleozoic Cycles of Sedimentation." Presented before the Geological Society of America, December 29, 1934.