

and that cessation occurs with the appearance of morphological symptoms of impairment.

(2) *Amoeba proteus* shows negative phototropism toward ultra-violet light.

(3) The cell membrane of irradiated amebas is relatively inextensible.

W. A. BLACK

UNIVERSITY OF CALIFORNIA

### A CONTRIBUTION TO THE PHARMACOLOGY OF PHYSOSTIGMINE<sup>1</sup>

WHILE investigating the peripheral action of barbiturates, we observed that in all experimental animals where the cardiac vagus response to weak faradic stimulation had been abolished by barbiturates doses of physostigmine salicylate ranging from 0.2 to 0.35 mgm per kgm showed no detectable spontaneous effect on the heart rate. However, if two to three minutes were allowed to elapse after intravenous administration of this drug and then the peripheral vagus was stimulated with the same weak, or even weaker, faradic current as used above profound cardiac inhibition was produced. Occasionally the slowing of the heart which was produced by stimulation of the peripheral vagus persisted for several minutes after stimulation

was discontinued. A similar cardiac slowing was noted following injection of acetyl choline, but it was not as marked and consistent.

This physostigmine sensitization of the vagus to stimulation lasted for about 30 minutes and was promptly antagonized by intravenous injections of further doses of barbiturates.

On the assumption that barbiturates produced their vagus-impairing effects by ganglionic depression, we employed in another series of experiments nicotine salicylate in doses varying from 2 to 4 mgm per kgm to produce ganglionic paralysis in dogs and rabbits. After sufficient nicotine salicylate had been administered intravenously to render the peripheral vagus non-responsive to even strong faradic stimulation, physostigmine (0.2 to 0.3 mgm per kgm) was injected intravenously. In every case within three minutes following the injection of physostigmine, faradic stimulation of the peripheral end of the vagus nerve produced marked cardiac inhibition. In other words, physostigmine antagonized the synaptic paralysis produced by nicotine.

CHARLES R. LINEGAR

JAMES M. DILLE

THEODORE KOPFANYI

## SCIENTIFIC APPARATUS AND LABORATORY METHODS

### A PROPOSED METHOD FOR THE DIRECT MEASUREMENT OF CORRELATION

ATTENTION is called to the possibility of applying certain relations given by Yule to the development of a device for the physical determination of the correlation coefficient. The relations are:

$$\Sigma_1^2 + \Sigma_2^2 = \sigma_1^2 + \sigma_2^2$$

$$r^2 = 1 - \frac{\Sigma_1^2 \Sigma_2^2}{\sigma_1^2 \sigma_2^2}$$

where  $\Sigma_1$  and  $\Sigma_2$  are the maximum and minimum standard deviations of the scatter about the intersection of  $\text{mean}_1$  and  $\text{mean}_2$ , and  $\sigma_1$  and  $\sigma_2$  have the usual connotations.<sup>1</sup>

Given the correlation surface actually constructed, the measurement of all these standard deviations in terms of moments should be feasible. The equivalence of  $\sigma^2$  to the moment of inertia of the distribution about its mean is well known. It is possible that a physical method has not yet been applied to correlation because of the difficulty of obtaining cross-products. If Yule's  $\Sigma_1^2$  and  $\Sigma_2^2$  can be physically measured, this difficulty is eliminated by the proposed method.

<sup>1</sup> From the Department of Pharmacology and Materia Medica, Georgetown University, School of Medicine, Washington, D. C.

<sup>1</sup> "Introduction to the Theory of Statistics," p. 322, 1919.

A moment of inertia is measurable as  $\frac{T t^2}{2\theta}$ , where  $T$  is torque applied (constant), and  $t$  is time to reach  $\theta$ , which is resultant angular displacement. What would seem necessary then would be a device utilizing unit rotational force acting through unit time, whence a moment would be found in terms of distance rotated. A spark marker could be used to measure this distance. Having found the means of the constructed correlation surface by balancing,  $\sigma_1^2$  would be obtained by rotation of the surface about the line through  $\text{mean}_1$  parallel to the axis of the other variable, and correspondingly for  $\sigma_2^2$ . The axes of rotation for finding  $\Sigma_1^2$  and  $\Sigma_2^2$  would be passed diagonally through the intersection of  $\text{mean}_1$  and  $\text{mean}_2$  (center of gravity). The angles of these axes would be varied until the minimum or maximum rotation had been noted. Inspection of the surface would reveal the angles for these axes closely for correlations higher than .30.

For plotting the scatters there could be provided light, rigid trays having twenty cells each way and a small post in each cell. With thin but heavy metal coins having holes in the centers, the plotter could construct the surface as rapidly as one plots the usual scatter on paper. The coins should fit snugly and be thin enough to give the surface a relatively small vertical dimension. The physical measurements of the