certain crops grown in the same field, particularly corn and timothy sod. It rarely occurred in continuous culture of tobacco, and fields strongly infested with brown root-rot usually recovered after one or two years of tobacco culture. It is thought that the rate of decomposition and nitrification of crop residues is an important factor governing the occurrence of the disease. Its ephemeral character under certain conditions is well established.

Because of the facts presented under 1, 2, 3, 4 and 5, it seems that the unoxidized forms of nitrogen resulting from the decomposition of organic matter are indirectly the cause of brown root-rot. Secondly, because of its ephemeral and transitory character and its relation to the cropping system, as shown under 6, 7, and 8, it appears that the unoxidized form or forms of nitrogen concerned originate from plant residues and possibly, also, from organic nitrogenous fertilizers supplied in large quantities in tobacco culture. Finally, it appears that the rotting of the roots is due to the comparatively high concentration of nitrogen in the roots, or a narrowed carbon-nitrogen ratio, caused by rapid absorption of basic nitrogen. The basic forms of nitrogen are not in themselves toxic in the concentrations usually found in the soil, but produce a condition in the roots which makes them easy prey to decay organisms commonly present in the soil. This explanation of the cause and nature of brown root-rot is presented as a working hypothesis, awaiting further work for its substantiation or disproval.

A. B. BEAUMONT

NAMES OF AND SYMBOLS FOR THE ARTIFICIALLY RADIOACTIVE ELEMENTS

THE system of naming the artificially radioactive elements, and especially the symbols used for these names, have apparently been adapted from the systematic naming of the naturally radioactive elements which follow radon in the radium family. Due to the number of elements known to be artificially radioactive, a possibility of ambiguity in the symbols has arisen. The symbols for radium F and for radiofluorine are Ra F and RaF; the symbol for radiosulfur and the formula of the compound radium sulfide are both RaS. The first pair of symbols is similar; in the second case, the symbol of an element and the formula of a compound are identical. In the event of a compound composed of two or more artificially radioactive elements, the formula may become very confusing: radiophosphorus radiotriiodide, RaPRaI, is an example. This could readily be confused with the formula of a molecular compound.

The systematic naming of the later members of the radium family is quite distinct from the naming of the artificially radioactive elements: the first are radium A, etc., the second, radiophosphorus, etc. One is a member of the radium family of elements; the other is not. To emphasize this difference, it is proposed that the artificially radioactive elements be named in such a way that the element is considered to be of primary importance, and the radioactivity secondary: the radioactivity is indicated by the hyphenated italicized prefix radio-; e.g., radio-phosphorus. This nomenclature is entirely analogous to that of isomeric organic compounds, such as para-nitrophenol, and the emphasis is placed on the element rather than on one of its attributes. The symbol ra-P contains only one capital letter, and so can not lead to the confusion now possible in chemical formulae. The symbols mentioned in the first paragraph would become Ra F and ra-F, ra-S and RaS, and ra-P-ra-I_a. The subscript and superscript now used for the atomic number and the atomic weight may be included in the usual manner. No confusion is anticipated between the symbol ra- for radio- and the r- (or occasionally ra-) which is sometimes used for racemic compounds.

BRIDGEVILLE, PA.

GERALD M. PETTY

THE WEATHERING OF FLINT ARTIFACTS

I WANT to get some information about the weathering or patination of flint flakes and artifacts. Under the direction of the Smithsonian Institution Dr. A. R. Kelly has been doing a systematic study of the Indian mounds so numerous in the vicinity of Macon, Georgia. This is said to be the largest enterprise of the kind in the eastern United States, employing several hundred men for nearly two years. I had noted in a deep trench cut through at least four former levels of Indian civilization that there was an increasing amount of patination of the flints at the lower horizons. The study of several lots of 250 to 1,000 artifacts taken from this and other places of study seems to reveal an average definite increase for apparent age. The larger number, perhaps all, of these flints are derived from the near-by residuum of the Ocala limestone, one of the local upper members of the Eccene period. This limestone is associated with the unconsolidated sediments of the Coastal Plain, which have not been subjected to metamorphism.

The amount of patination found varies from zero to more than four millimeters, and the effects are found in all types of flints. The more attractive jaspers are undoubtedly affected by actinic rays, but a loosening of their silicic binding material occasionally exceeds the bleaching. Information from Dana, Merrill and others throws little light on the subject. Eoliths from Kent, England, show a different type of patination, which would prove to be affected easily by frost and other weathering factors.

No flints secured from localities of metamorphic