



such that they might well be concerned with light reception, but carefully executed experiments will be necessary in order to make this matter certain.

JOHN H. WELSH

BIOLOGICAL LABORATORIES
HARVARD UNIVERSITY

THE DISCOVERY AND IDENTIFICATION OF A NEW PURINE ALKALOID IN TEA

ALL the N-methyl derivatives of 2,6,8-trioxypurine I theoretically possible have been prepared synthetically, and we have to-day a very complete knowledge of their chemistry. Emil Fischer and Heinrich Biltz, with the collaboration of many coworkers, are the two investigators who have contributed the most to our present knowledge of the chemistry of these purines and their derivatives.



The occurrence of 2,6,8-trioxypurine as a product of purine catabolism in both the animal and plant kingdoms has been demonstrated conclusively, but, so far as the writer is aware, no N-methyl derivative of the purine I has, thus far, been shown to occur in nature. The author now presents this short note to report that the *tetramethyl-2,6,8-trioxypurine* represented by Formula II occurs in the mixture of purine alkaloids extracted from tea. It has been separated in a pure condition from such extracts, and has been shown to be identical with *1,3,7,9-tetramethyl-2,6,8-trioxypurine II* (tetramethyluric acid), which was first described by Emil Fischer¹ in 1884. Just as soon as proper and sufficient experimental material becomes available for the continuation of our plant extract researches, it is the intention of the author to search for this alkaloid and other N-methylated purines in the purine extracts of coffee and other plants. The results of this research program will be discussed in future papers to be presented for publication in the *Journal* of the American Chemical Society.

TREAT B. JOHNSON

YALE UNIVERSITY

SCIENTIFIC APPARATUS AND LABORATORY METHODS

A CONDENSER DISCHARGE STIMULATOR FOR PHYSIOLOGICAL PURPOSES

THE stimulator described in this report has been designed for the determination of the adequate shape and duration of current pulses used as stimuli on the cerebral motor cortex.¹ It has been found a useful device for general stimulation experiments wherever an attempt is made to gain more information about the excitable structures responsible for a certain effect. It also provides for a selective stimulation² in mixed peripheral nerves or in mixed tracts or centers within the central nervous system.

The set-up is based on the principle of condenser discharges adapted to the relatively low resistance of the tissue to be stimulated through a single-stage power amplifier.³ It allows stimulation with alternating single or double condenser discharges whose duration, *i.e.*, time constant, can be changed over a wide time range (from .01 to 100 or 1000 milliseconds) without any change at all in the amplitude (peak intensity) of the discharges. The stimulating voltage, up to 10 or 20 volts, is led off from a potentiometer of 2,000 ohms maximum resistance. Any influence of the stimulating circuit upon the time constant of the condenser system is excluded.

¹ O. A. M. Wyss and S. Obrador, *Am. Jour. Physiol.*, in press (1937).

² O. A. M. Wyss, *Schweiz. Arch. f. Neur. u. Psych.*, 28: 210, 1932.

From a source of potential A two condensers C and C^1 of different capacities are charged by make of K over two identical resistances R and R^1 to the same voltage, and they discharge over the same two resistances and a common resistance S , low in comparison with R and R^1 , when K is opened. The resulting potential wave between x and y has its shape and direction determined by the ratio of one capacity to the other. It represents an ordinary condenser discharge if one condenser is disconnected, and a double condenser discharge (see³) if both condensers are placed in the circuit. Any such potential wave between x and y causes in the plate circuit of the amplifier tube AT (Cunningham 2A3) a current wave of identical shape. The resting plate current of AT is compensated by another similar tube CT with adjustable heater resistance. Equilibrium between the two tubes, *i.e.*, absence of potential between the ends of the potentiometer p , is controlled by a high resistance galvanometer v . Both halves of the potentiometer are divided into twenty intervals of 50 ohms each. Provided that the stimulating current is always led off from two symmetrical steps on the corresponding halves of the potentiometer, the current in the stimulating circuit is only due to, and directly pro-

¹ Emil Fischer, *Ber.* 17: 1784 (1884); also *Ber.* 30: 3009 (1897).

³ *Idem.*, *Pflüger's Arch.*, 233: 754, 1934.




Fig. 1

portional to, changes in the plate current of AT ; its shape, therefore, is that of the potential wave between x and y .

For double condenser discharges the most convenient ratio between C and C^1 has been found to be 2 to 1 (see³). The resulting peak potential between x and y reaches then 25 per cent. of the applied voltage A , whereas for single condenser discharges the full potential appears between these two points. Therefore, from a dry cell battery used in A , 36 volts were picked out for double condenser discharges and 9 volts for single condenser discharges. The condensers C and C^1 are variable from .001 to 1, or very slow current waves, 10 microfarads.⁴ With two pairs of identical resistances for R and R^1 , of 10,000 ohms and 100,000 ohms, respectively, a range of time constants as wide as 1 to 10^5 can be covered. The resistance S has to be kept as low as possible; a few hundred ohms are negligible, even with 10,000 ohms in R and R^1 .

The duration of the single current pulse is determined (a) for single condenser discharges by the time constant: resistance $R \times$ capacity C , and (b) for double condenser discharges by the duration of the rising phase which is equal to $0.7 \times R \times C$, C being the greater capacity ($C = 2 C^1$). The time values are obtained in seconds if R and C are expressed in ohms and farads, respectively. If, therefore, for R and R^1 two pairs of resistances of either 14,500 or 145,000 ohms are used, the duration of the rising phase of the double condenser discharge is directly given by the number of microfarads of the greater capacity, one microfarad corresponding to 10 or 100 milliseconds,

⁴ Decade condensers of General Radio Company, Cambridge, Mass.

respectively. The full wave duration is approximately five times the rising phase, which has to be considered if repetitive stimulation is used. In the latter case the ordinary contact key K has to be replaced by an automatic commutator.

OSCAR A. M. WYSS

LABORATORY OF PHYSIOLOGY
YALE UNIVERSITY
SCHOOL OF MEDICINE

A METHOD FOR OBSERVING THE LOWER SURFACE OF SMALL OBJECTS

WHERE a simple device for examining the lower surface of small objects under the dissecting microscope is needed, a mirror may be used as a supplementary stage. The upper surface of the object can then be viewed by reflected light in the usual way, or by racking the microscope down twice the thickness of the mirror, the image of the lower surface of the object may be brought into focus. The object itself rarely causes any trouble while viewing the image; as, if the mirror is of adequate thickness, the object will be out of the line of vision and out of focus, often being entirely invisible. For the lower magnifications, where the objectives have a great depth of focus, the thickness of the mirror may need to be increased either by adding a supplementary glass plate or by raising the slide above the mirror on plasticene legs. This device, which no doubt has been repeatedly used, has several advantages, namely, the cheapness and availability of the material; the excellence of the illumination; and the speed and ease of changing views without the necessity of touching the slide or removing the hand from the focusing adjustment.

RALPH J. BAILEY

UNIVERSITY OF ARKANSAS
SCHOOL OF MEDICINE

BOOKS RECEIVED

- American Mathematical Society Colloquium Publications, Vol. XXI. *Differential Systems* by JOSEPH M. THOMAS. Pp. ix + 119. The Society.
- BLAIR, THOMAS A. *Weather Elements; A Text in Elementary Meteorology*. Pp. xv + 401. 107 figures. Prentice-Hall. \$5.00.
- EMERSON, ALFRED E. and ELEANOR FISH. *Termite City*. Pp. 127. Illustrated. Rand, McNally. \$1.50.
- Göteborgs Högskolas Årsskrift, XLII, 1936: 3. *Determinismus und Indeterminismus in der Modernen Physik; Historische und Systematische Studien zum Kausalproblem*. ERNST CASSIRER. Pp. xi + 265. Wettergren & Kerbers Förlag, Göteborg.
- HICKMAN, CLEVELAND P. *Physiological Hygiene*. Pp. xxvi + 493. 89 figures. Prentice-Hall. \$3.25.
- National Resources Committee, December, 1936. *Drainage Basin Problems and Programs*. Pp. viii + 540. U. S. Government Printing Office. \$1.50.
- Ölstra Gåtor Aktuella Problem. Pp. 302. Illustrated. Wahlström & Widstrand, Stockholm.
- Travaux et Memoires de l'Institut d'Ethnologie, XXIV. *Traité de langue amharique (Abyssinie)*. Marcel Cohen. Pp. xv + 444. Institut d'Ethnologie, Paris.