

trates furthermore the increased tendency of the body to retain water when part of the hydrogen has been replaced by deuterium. A decrease in the insensible water loss relative to the metabolism is described during the period when mice are being raised, either in two days or in five days, to the level of one fifth saturation with D_2O .

It is quite likely, too, that the reversible contraction of *Fundulus melanophores*, which we have also demonstrated,⁵ depends partly upon an osmotic factor.

So far as mice are concerned, then, we have yet to find evidence that D_2O "violently" dehydrates living cells. But a number of interesting manifestations of its lower vapor pressure are illustrated by its pharmacological action in mammals.

HENRY G. BARBOUR

YALE UNIVERSITY

THE NON-VOLATILE ACIDS OF THE FRUIT OF NYSSA OGECHÉ¹

THE "Ogeechee Lime" appears as a small fruit on the tree *Nyssa ogeche*, which grows in wet ground in the southern states. The fruit which came to the attention of the author was from along the Conochee River in Georgia. They were light green, olive-shaped fruits, 2.5 to 3.5 cm long with a large single central stone and a thin skin, with cell-walls connecting skin and stone which were firmly attached to the stone, running lengthwise. Forty to 50 per cent. by weight of juice can be expressed by squeezing the fruit. This juice is strongly acidic, but lacks any distinctive flavor. The juice showed about 6 per cent. of acid, calculated as citric. It is said that natives of the regions where they grow use them to make a "limeade" drink and a preserve.

The non-volatile acids were isolated from 100 fruits weighing 637 gm and converted to ethyl esters by the usual method of lead salt precipitation and esterification. A small portion (2 gm) boiled at 150–171° at 10 mm and the remainder boiled at 171–172° at 10 mm (12 gm). The hydrazides of the fractions were prepared. From the low-boiling fraction malic hydrazide, m. p. 176–177° (mixed m. p. 176–177°) was identified. The fraction boiling at 171–172° at 10 mm was citric ester, giving a hydrazide melting at 102–104° when allowed to form spontaneously (hydrated form) and melting at 149–150° when seeded with anhydrous citric hydrazide. It is evident that the principal acid of the Ogeechee lime is citric acid with a small amount of malic.

D. H. WHEELER

¹ H. G. Barbour and Lillie Rice, *Jour. Pharm. and Exp. Therap.*, 1937 (in press).

⁵ H. G. Barbour and S. B. Bogdanovitch, *Jour. Pharm. and Exp. Therap.*, 61: 148, 1937.

¹ Food Research Division Contribution No. 348, Bureau of Chemistry and Soils, U. S. Department of Agriculture.

IODOACETIC ACID AND SULFUR METABOLISM

IN view of the suggestions in the literature that iodoacetic acid exerts its inhibitory effect on certain body processes (*e.g.*, absorption, muscle metabolism) by combining with a sulfhydryl compound which may be essential for these processes, experiments have been designed in an attempt to demonstrate *in vivo* a combination of iodoacetic acid with sulfur-containing substances which are of biological importance. Using a technique developed in this laboratory,^{1, 2} it has been possible to markedly restrict the growth of young white rats, ingesting a relatively low protein diet, by incorporating suitable quantities of iodoacetic acid in the basal ration. Moreover, the addition of either l-cystine or dl-methionine to the basal diet already containing the iodoacetic acid results in an immediate growth response, and the animals continue to grow at a good rate. This response is striking and appears, to date, to be specific for cystine and for methionine. This study is being actively continued and expanded along the following lines of investigation: (1) To determine the mechanism by which iodoacetic acid appears capable of depleting the sulfur-containing amino acid reserves of the organism; (2) to determine the nature of the *in vivo* combination between iodoacetic acid and these sulfur compounds; (3) to determine the possible relationships of this type of combination to the processes of absorption from the intestine and muscle metabolism.

ABRAHAM WHITE

LABORATORY OF PHYSIOLOGICAL CHEMISTRY,
YALE UNIVERSITY SCHOOL OF MEDICINE

AN INTERESTING HOAX PERPETUATED IN AN EARLY SCIENTIFIC BULLETIN

READERS of *Time* for June 7, 1937, were no doubt amused by the article concerning the spider hoax perpetrated by Ralph D. Paine in the 1890's. The article reported that several scientific journals printed the hoax as fact. It may be of interest to know that one of these scientific journals was the *Bulletin of the U. S. Department of Agriculture, Division of Entomology*. The title of the bulletin, which was published in 1897, is "Some Miscellaneous Results of the Division of Entomology." Under "General Notes" on page 82 there is the following sub-head and account:

A NEW DIRECT BENEFIT FROM INSECTS

When Kirby and Spence wrote their chapter on "Direct benefits derived from insects" and recorded the use of insects for food, the use of honey from bees for the same purpose, the use in medicine and the arts and manufacturers of blister beetles, insect galls, Coccidae furnishing lac, wax insects and the silkworm, the time had hardly

¹ A. White and R. W. Jackson, *Jour. Biol. Chem.*, 111: 507, 1935.

² A. White, *Jour. Biol. Chem.*, 112: 503, 1936.