

is placed on the maze at the entrance, it is stimulated by the maze and the experimenter. Call this total stimulation S_E . S_E causes a response of running left at the first cul. Call this response R_L . This turn to the left and the approach to the next choice point produce more stimulation from inside and outside the maze as well as kinesthetic stimuli within the rat. We shall call all these sensory consequences of R_L by the symbol S_L . S_L causes the rat to turn right (R_R) at the next choice point; and R_R has S_R as its sensory consequences. S_R is then followed by a left turn, etc., through the maze. What the animal has had to learn, then, is to turn left for S_E and for S_R and to turn right after being stimulated by S_L . We shall leave aside the question whether the rat responds to S_L and S_R as individual stimuli or as interrelated patterns, since our major concern is with the sensory modality which is controlling the behavior.

If the maze were not rotated and the units were not interchanged, there would be cutaneous and olfactory stimuli from the various maze units as well as kinesthetic stimuli from the animal's own activity which, with stimuli from outside the maze, would enter as components of S_L and S_R as has been shown in the work of Hunter, Honzik and Wolfe. If, however, as under the present conditions, the maze is rotated and the units are interchanged, S_L and S_R reduce solely to the kinesthetic stimuli resulting from the rat's own activity in making left and right turns, in so far as differential stimuli are concerned. To be sure, the edges of the paths stimulate the rat and "show" him when a turn is to be made, but such stimuli can not determine the difference between a left and a right turn.

It is theoretically possible that some symbolic process, some symbolic equivalent of "go left-right-left-etc.," may be present to control a habit set up under the above conditions, as indeed it would be were the sequence of turns in double alternation *lrrllrr*.⁷ However, the presence of adequate differential kinesthetic stimulation makes such a hypothesis untenable on the basis of parsimony; and the great difficulty which the rat has with double alternation temporal problems as compared with the great ease with which the present maze habit was established renders the hypothesis of symbolic control still less tenable in the present case.

The significance of the present experiment lies not only in its demonstration of a habit built up and controlled by kinesthesia but also in the basis which is now laid for an extension of the neurological study of such intelligent acts. A repetition of Lashley and Ball's experiment on rats trained as here described (but in alley mazes because of the rat's motor inco-

ordinations) should show an interference with the habit if kinesthetic impulses are carried in definite tracts of the cord and if they can not shift to other tracts when their own are destroyed. Furthermore, cortical extirpations of the kinesthetic projection areas should reveal a localization of function in some rats as opposed to the equipotentiality which Lashley's work on the neural control of maze habits has hitherto supported. The expression "in some rats" is used advisedly because some rats can, with difficulty, develop the equivalent of symbolic controls. It remains to be determined whether or not localized brain lesions destroy the rat's capacity to perform the responses so controlled. (Jacobsen's⁸ work indicates that monkeys are unable to relearn the delayed reaction, which is also symbolically controlled, after bilateral frontal lobectomy.)

WALTER S. HUNTER

BROWN UNIVERSITY

A RADIOACTIVE ISOTOPE STUDY OF THE ABSORPTION OF PHOSPHORUS AND SODIUM BY CORN SEEDLINGS

THE artificial radioactive method is readily adapted to studies of the factors influencing the absorption of mineral matter by plants. The design of Geiger counter developed in this laboratory¹ is suited to studies of this type, since it has a low background of only two counts per minute, since it will detect rays with energies as soft as 50,000 volts and since it yields quantitative results when calibrated against known solutions.


The technique involved in a quantitative study of the absorption of phosphorus and sodium by corn seedlings is simple. The plants tested were germinated in sand and the seedlings transferred to water or culture solution; at the time of the experiment they were about 12 inches high. In making the tests a measured quantity of radioactive phosphorus in the form of tripotassium phosphate or of radioactive sodium as sodium chloride was added to the culture solution. At regular intervals thereafter the plant was placed so that the leaf to be examined was directly below the counter window and the number of counts per minute were recorded.

Effects of various factors influencing the assimilation of phosphorus and sodium are illustrated in Figs. 1, 2 and 3. Fig. 1 shows the total number of disintegrations counted per minute per unit area of leaf as a function of time for different total phosphorus concentrations in the solution. Total phosphorus (radioactive and non-radioactive) taken up

⁸ C. F. Jacobsen, *Comp. Psychol. Monog.*, 13: No. 63, 1936.

¹ A. K. Brewer and A. Bramley, *Jour. Applied Physics*, 9: 778, 1938; *Jour. Ind. Eng. Chem.*, 30: 893, 1938.

⁷ W. S. Hunter, *Jour. Genet. Psychol.*, 36: 512, 1929.



FIGS. 1-3

by the leaf and expressed in milligrams per sq. cm is computed directly from the number of counts per minute. The second leaf from the bottom, 2 inches from the base and 6 inches from the solution, was taken for study. Five minutes after radioactive phosphorus was added it could be detected in the leaves. Curves I, II and III show that for the concentrations tested the uptake increases with the amount supplied. Curve IV shows less uptake than Curve II, because in this case the plant was allowed to stand for 24 hours in a standard nutrient solution containing 0.0008 g of phosphorus per cc prior to the addition of radioactive phosphorus. Curve A in Fig. 3 is a corresponding uptake curve for sodium.

The distribution of phosphorus in the plant changes appreciably with time. During the initial stages of the experiment the number of disintegrations was from 25 to 50 per cent. more intense for the bottom than for the top leaf; it also decreased rapidly from the base to the tip of the individual leaves. The final distribution, in general, was quite uniform between the leaves as well as along each leaf except at the tip, which remained low.

Effect of the extent of root system on uptake was investigated in detail. For both sodium and phosphorus the rate and equilibrium value were found to be the same for intact roots, for plants with roots excised at different positions and for leaves removed at the base. Plants stunted in their early growth by a deficiency in their diet took up very little sodium or phosphorus due to an impairment which began in

the root system and gradually extended up the stalk; in all stunted plants tested normal uptake was observed when the impaired section was removed.

Effect of surrounding conditions on the uptake of phosphorus is shown in Fig. 2. In each instance the plant was kept in the condition designated for 15 hours before adding radioactive phosphorus. Unfortunately it was necessary to expose the plants to room conditions for about three minutes every hour while the observations were being made. Absorption was very low while plants were in the dark but increased rapidly upon exposure to light. Low temperatures and an atmosphere of carbon dioxide decreased the uptake.

Elimination of phosphorus and sodium from the plant presents a different picture. Curve B, Fig. 3, shows decrease of sodium in the leaf when the plant (root system intact) was transferred from a solution containing radioactive sodium to one containing sodium. Curve C shows the increase of sodium in the solution due to back diffusion from the plant. Analogous results have been obtained by Jenny, Overstreet and Ayers² on the depletion of sodium from barley plants. The phosphorus content of the leaf, in contrast, decreased only slightly with time, while the radioactivity of the solution increased slowly through loss of phosphorus from the roots and lower portion of the stalk.

The writers are indebted to Dr. L. A. DuBridge, of the University of Rochester, for the radioactive phosphorus and Dr. M. A. Tuve, of the Carnegie Institution of Washington, for the radioactive sodium.

A. KEITH BREWER

ARTHUR BRAMLEY

U. S. DEPARTMENT OF AGRICULTURE

A METHOD OF STUDYING THE AVAILABILITY OF VARIOUS SUBSTRATES FOR HUMAN BRAIN METABOLISM DURING THERAPEUTIC INSULIN SHOCK¹

PERFUSION of whole organs,² arterial and venous blood sampling from intact organs in the body³ and studies of surviving excised tissues in the manometric apparatus⁴ have been the chief methods by which the metabolism of various organs, particularly the brain, have thus far been studied. The method of blood sampling has been successfully applied to the study

² Jenny, Overstreet and Ayers, *Soil Science*, 48: 9, 1939.

¹ From the Division of Psychiatry, Bellevue Hospital, and the Department of Psychiatry, New York University Medical College, New York, N. Y.

² A. L. Chute and D. H. Smyth, *Quart. Jour. Exp. Med.*, 29: 379, 1939.

³ E. S. London, N. P. Kotscheref, A. M. Dubinsky and A. S. Katzwa, *Arch. ges. Physiol.*, 233: 160, 1933.

⁴ I. H. Page, "Chemistry of the Brain," Chas. C. Thomas, Baltimore, 1937.