

nicotinic acid by sheep on various regimens are average values for 3 consecutive 24-hour collections. While there is considerable variation in the amount of nicotinic acid excreted in the urine the level is not essentially different, irrespective of whether or not the diet is deficient. Supplementing the deficient diet with nicotinic acid augmented the excretion. The most probable explanation of the continued excretion of nicotinic acid by sheep on a diet deficient in this constituent is that this species can synthesize it either in its tissues or it is formed in the rumen by microorganisms, a process analogous to the synthesis of thiamin, riboflavin, B<sub>6</sub>, and pantothenic acid.<sup>4</sup>

A. H. WINEGAR  
P. B. PEARSON  
H. SCHMIDT

AGRICULTURAL EXPERIMENT STATION,  
AGRICULTURAL AND MECHANICAL  
COLLEGE OF TEXAS

#### BREAKDOWN OF SULFANILAMIDE MOLECULE BY ULTRA-VIOLET IRRADIATION OR CHEMICAL OXIDATION

In the course of some experiments with ultra-violet irradiation it was observed that dilute aqueous solutions of sulfanilamide became more acid (glass electrode or methyl red as pH indicator). This was taken as an indication of oxidation of the sulfonamide group with liberation of acid sulfur radicals. Proof of a breakdown of the molecule was obtained in that irradiated solutions showed the presence of ammonia by reacting strongly to Nessler's reagent, and of inorganic sulfur by giving a precipitate with barium chloride in acid solution. That the free amino group is also affected has been previously shown by Ottenberg and Fox<sup>1</sup> by a decrease in the diazo reaction under such conditions. By the use of light filters the most effective wave lengths were found to be below 270 mμ, the same region of the spectrum shown by Fox<sup>2</sup> to bring about the colored products.

The amount of sulfur split off was a function of the length of irradiation. When aqueous solutions of sulfanilamide containing 40 mg per cent. were irradi-

ated by a Hanovia Mercury Arc Lamp at 20 cm distance for 2, 4, 8 and 16 minutes the amounts of sulfur split off were 8.6, 16.1, 29.1 and 40.8 per cent., respectively, of the total amounts present. Varying the concentration of sulfanilamide from 20 to 100 mg per cent. caused only slight variations in the total amounts of inorganic sulfur recovered after a uniform exposure of 10 minutes.

Irradiation of the ortho and meta isomers of sulfanilamide, of acetyl sulfanilamide and of sulfanilic acid for 10 minutes under the above condition did not bring about similar changes, except for the liberation of some ammonia with sulfanilic acid.

Oxidation of sulfanilamide by chemical agents has been carried out by Shaffer<sup>3</sup> whose attention was directed to the amino group. Preliminary experiments have shown that the oxidation of dilute aqueous solutions by ferric chloride and hydrogen peroxide is also attended by the liberation of ammonia and inorganic sulfur. The amount of sulfur split off was likewise dependent upon the amount of iron added, indicating that the process is not a catalytic reaction.

After standing at room temperature for 18 hours, 400 cc of an aqueous solution containing 100 mg of sulfanilamide, 2.2 mg FeCl<sub>3</sub> and 0.5 cc 3 per cent. H<sub>2</sub>O<sub>2</sub> showed 15.7 per cent. of sulfur split off. Under similar conditions the ortho isomer showed 9.7 per cent., the meta isomer 4.8 per cent. and sulfanilic acid (neutralized) 18.3 per cent. of sulfur split off. These results are not comparable to those with irradiation, which was applied for only 10 minutes.

Because of the irreversible nature of these changes it is obvious that potentiometric studies of such solutions are not valid. Whether or not the body is capable of splitting off any of the sulfonamide group remains to be demonstrated. The demonstration by James<sup>4</sup> of p-aminophenol in the urine following sulfanilamide therapy is evidence to this effect.

SANFORD M. ROSENTHAL  
HUGO BAUER

NATIONAL INSTITUTE OF HEALTH,  
WASHINGTON, D. C.

## SCIENTIFIC APPARATUS AND LABORATORY METHODS

### A SPEEDIER AND LESS COSTLY METHOD OF CONCENTRATION IN NITRO-CELLULOSE IMBEDDING

In the hot nitrocellulose method of imbedding histological and cytological material as developed by

Jeffrey,<sup>1</sup> and somewhat elaborated by Wetmore,<sup>2</sup> chips of celloidin in small metal troughs are used to absorb the ether and alcohol and thus concentrate the liquid in which the material is being imbedded. Around any laboratory using this process a certain amount of used celloidin soon accumulates. It often contains particles of imbedded material as well as dirt from containers

<sup>4</sup> L. W. McElroy and H. Goss, *Jour. Biol. Chem.*, 130: 437, 1939.

<sup>1</sup> R. Ottenberg and C. L. Fox, Jr., *Proc. Soc. Exper. Biol. and Med.*, 38: 479, 1938.

<sup>2</sup> C. L. Fox, Jr., J. E. Cline and R. Ottenberg, *Jour. Pharm. and Exper. Therap.*, 66: 99, 1939.

<sup>3</sup> P. A. Shaffer, *SCIENCE*, 89: 547, 1939.

<sup>4</sup> G. V. James, *Biochem. Jour.*, 33: 1688, 1939.

<sup>1</sup> E. C. Jeffrey, *Bot. Gaz.*, 86: 456-467.

<sup>2</sup> R. H. Wetmore, *Stain Tech.*, 7: 37-62.

and tables. The author has found that when this is redissolved in the usual ether-alcohol solvent and allowed to stand for a week or so, almost all of the refuse will sink to the bottom of the jar. The nitrocellulose may then be poured into petri dishes to a depth of a quarter of an inch. A little chloroform poured on top will aid in the solidification which should proceed until the celloidin is dry enough to cut with the shears or a razor. Cut into squares of approximately an inch, gather these on a No. 1 or longer pin, and allow to dry completely. These may then be used in place of the troughs in the position as shown in Fig. 1. The use of a pin diminishes the chance of pro-




FIG. 1.

ducing a leaky cork after several changes and with the increase of absorbing surface the time required to concentrate the 10 per cent. celloidin is shortened by several days. This is also dependent upon the number of times the pins are changed each day; usually twice being sufficient. If mounted material is placed in the bottles with the cardboard side up and if the oven shelf is raised a bit in front, the hazard of finding some of the material sticking out of the celloidin is much reduced. For most cytological material the nitrocellulose series may also be cut down to changes of 2, 6 and 10 per cent. without any apparent damage to the material. The above recommendations reduce the time factor which some investigators find objectionable in a method of imbedding which is otherwise quite superior to paraffin.

HENRY WILHELM JENSEN

ASHVILLE FARM SCHOOL,  
SWANNANOVA, N. C.

### AN INEXPENSIVE MICROPHOTOGRAPHIC CAMERA

A "UNIVEX" Camera, Model A F, which costs but \$1.00, can be adapted for the filming of microscopic objects with little trouble and less mechanical skill. This camera is provided with instantaneous and time settings on its shutter. In order to use it in conjunc-

tion with a microscope, one has merely to remove the lens of the camera. This is done by setting the shutter for "time" and opening it. A nail is held against the lens and is then struck a smart blow with a small hammer. The lens is easily shattered, and the pieces can be shaken out.

The wire frame that is normally used as a finder is drawn out and the middle (horizontal) section is removed by bending the angles a few times. The two side pieces are then bent down along the front of the lens board and then curved slightly to receive the eye-piece of the microscope. The two wires are held together by means of a loop of wire passed around them just above the eye-piece.




FIG. 1

An eye-piece is clipped to the camera and then placed in the microscope. In order to focus the camera, a piece of ground glass or tracing paper 33 × 45 mm is placed over the opening of the camera, with the back removed. The roll of film, which costs but 10 cents, is placed in the camera and six exposures are made with varying times. At least one of the exposures will give a sharp negative which can be printed directly or enlarged many times since the grain is very fine.

No rules for exposure can be given since that depends so much on the illumination.

E. M. ABRAHAMSON

THE JEWISH HOSPITAL OF BROOKLYN

### BOOKS RECEIVED

- American Philosophical Society. *Year Book, 1939*. Pp. 494. The Society, Philadelphia.  
 DRESDEN, ARNOLD. *Introduction to the Calculus*. Pp. xii + 428. Holt. \$3.40.  
 ELLIS, CARLETON. *Printing Inks; Their Chemistry and Technology*. Pp. 560. 96 figures. Reinhold. \$7.00.  
 KLINEBERG, OTTO. *Social Psychology*. Pp. xii + 570. Holt. \$3.00.  
 MORGAN, ALFRED. *Things a Boy Can Do with Chemistry*. Pp. xvii + 288. Illustrated. Appleton-Century. \$2.50.  
 National Research Council. *Twelfth Report of the Committee on Catalysis*. Pp. x + 388. Wiley. \$5.00.  
 WARDEN, CARL J., THOMAS N. JENKINS and LUCIEN H. WARNER. *Comparative Psychology; Vol. II, Plants and Invertebrates*. Pp. xiii + 1070. 168 figures. Ronald Press. \$6.00.  
 WENIGER, WILLIBALD. *Fundamentals of College Physics*. Pp. viii + 694. 340 figures. American Book Co. \$3.75.