

and the uncontrollable escape of fibers. It is gradually transformed into perineural connective tissue and after several months can no longer be identified. Regenerating fibers traverse the wound in straight parallel courses instead of in the usual confusion following nerve suture.

It has recently been described⁵ that good binding between severed nerve trunks can be obtained by applying clotting blood plasma to the apposed cut ends, and that regenerating fibers under these conditions take fairly straight courses in crossing the scar.⁶ Whether this method provides the experimenter with as definite an insurance against undesirable stray fibers as tubulation does, remains to be seen.

PAUL WEISS

THE UNIVERSITY OF CHICAGO

A METHOD FOR DETERMINING AND SPECIFYING LOCALITY BY COLLECTORS

WHILE engaged in studying the flora of Maryland the author devised a system for determining and specifying locality which may be of interest to collectors. The system employs the topographical maps published by the United States Geological Survey which are scaled one inch to the mile. These quadrangle maps are already divided by the coordinates of longitude and latitude into nine nearly equal and approximately rectangular divisions. The longitudinal meridians, of course, converge to the north so that the northern boundary of the quadrangle and the northern boundary of every primary division is, in each case, slightly smaller than the opposite or southern boundary. The nine primary divisions of each quadrangle are numbered .1 to .9 counting from left to right: top row, .1, .2, .3; middle row, .4, .5, .6, and bottom row, .7, .8, .9.

A transparent celluloid over-layer is cut the size of an average primary division for the area to be studied. Since no over-layer can be made to fit all the quadrangular divisions for an area as large as the United States, due to the northward convergence of the meridians of longitude, it is necessary to take the average primary division in a given area as the size of the over-layer to be used.

The over-layer may be prepared from either of two kinds of celluloid: (a) with one side frosted so that it may be ruled with India ink, or, (b) with both sides smooth so that lines may be etched or cut into one side and subsequently filled with colored wax or India ink. This over-layer is divided by coordinately ruled lines into nine equal and approximately rectangular divisions each of which represents a secondary division of the quadrangle.

These secondary divisions are numbered .01 to .09, and in the manner indicated above for the primary divisions. Each secondary division is, in a similar way, divided into nine tertiary rectangles and numbered .001 to .009. Finally, each tertiary subdivision is divided into nine rectangles which are numbered .0001 to .0009. Thus, the preparation of the over-layer in the manner indicated provides four sizes of rectangular divisions.

By placing the celluloid over-layer on any one of the primary rectangular divisions of the quadrangle map, any point on the division can be read as a four point decimal. The fifth decimal can be estimated if desired and will locate the point to within approximately a 250 foot radius. Since each quadrangle already bears a name (printed on the map), this name precedes the decimal code in specifying locations, as, for example: "Ellicott .99887."

For field use, the author has found it convenient to cut each topographical map into its nine primary divisions and paste these into a loose-leaf notebook of appropriate size where each is labeled with the name of the quadrangle from which it was cut and the proper primary division number, as, for example: "Ellicott .9." Through the use of these maps in the field, specific locations are determined and recorded at the time of collecting each specimen.

Fortunately, the over-layer described above can be used on the soil maps which are published by the Bureau of Chemistry and Soils, United States Department of Agriculture, and which are scaled one inch to the mile. For field use, as well as for other kinds of uses, it is convenient to mount the soil maps in the same loose-leaf form referred to above, and with each map opposing the page on which the corresponding topographic map is mounted. The simultaneous use of both maps often brings to light very interesting ecological features.

CLYDE F. REED

MARYLAND ACADEMY OF SCIENCES,
BALTIMORE

BOOKS RECEIVED

- Japanese Journal of Astronomy and Geophysics: Transactions.* Pp. 331-565. Illustrated. National Research Council of Japan, Tokyo.
- Journal of Science of the Hiroshima University.* Series A, Mathematics, Physics, Chemistry, Vol. 10, No. 3. Pp. 117-269. Series B, Zoology, Vol. 8. Pp. 164. Illustrated. The University, Hiroshima, Japan.
- JOHANNSEN, OSKAR A. and FERDINAND H. BUTT. *Embryology of Insects and Myriapods.* Pp. xi + 462. 370 figures. McGraw-Hill. \$5.00.
- RORIMER, IRENE T. *A Field Key to Our Common Birds.* Pp. 160. 27 figures. 18 plates. Cleveland Museum of Natural History. \$1.50.
- University of Missouri Studies, October, 1940. Secret Societies; A Cultural Study of Fraternalism in the United States.* NOEL P. GIST. Pp. 184. The University, Columbia. \$1.25.

⁵ J. Z. Young and P. B. Medawar, *Lancet*, 1940: 126.

⁶ These experiments essentially reproduce conditions first established in tissue culture by the author (P. Weiss, *Jour. Exp. Zool.*, 68: 393, 1934) and confirm the guiding effect of an oriented fibrin matrix on nerve fiber growth.