

remembered that in 1936 a mathematical periodical was started in Germany under the title *Deutsche Mathematik*, which has since then been widely supported by German mathematical writers even if the contents exhibit the fact that mathematics is an international subject which contains many evidences of the cooperation of writers of many lands. In recent years the German contributions have been extensive, but they were largely based on the earlier contributions of writers in other lands including the Greeks whose achievements the noted prize may help to clarify and to whom the entire mathematical world has often acknowledged itself indebted notwithstanding the growing credit to earlier civilizations.

This prize and the recently reported appointment of a professor of the history of mathematics in the University of Berlin seem to imply that this history is now receiving relatively much attention in Germany, notwithstanding the fact that the present disrupted condition in scientific work makes the unbiased study of this subject very difficult. The articles in the periodical noted in the preceding paragraph also indicate an emphasis on the history in recent German mathematical writings. In so far as these efforts are directed towards learning the actual situations they naturally receive the approval of all and should be especially appreciated in America in view of the relatively small amount of such work in our rapidly expanding mathematical activities of recent years. According to Felix Klein the thinking through of old problems by new methods is the source of pure mathematics.

G. A. MILLER

UNIVERSITY OF ILLINOIS

## CROSS-FERTILIZATION OF ECHINODERMS

It has long been known that the sea urchins *Strongylocentrotus purpuratus* and *S. franciscanus*, will cross-fertilize, reciprocally, but every investigator has noted that the percentage of cross-fertilizations is extremely variable. During a recent stay at the Hopkins Marine Station at Pacific Grove, Calif., it was found that the variability was largely due to the method of preparing the eggs for experimental use. The usual procedure of preparing sea urchin eggs is as follows:

The ovaries are removed from the cut animal and placed in a fairly large quantity of sea water; this is filtered through cheese-cloth to remove the debris from the exuded eggs; some investigators advocate several washings of the eggs with fresh sea-water; then a small quantity of the eggs are pipetted off into a Syracuse watch glass containing fresh sea water, and the sperm added. Using eggs prepared in this manner, it was found that when crossed with the sperm of the other species, very few eggs were fertilized, though 100 per cent. fertilizations occurred with the sperm of its own

species. If, however, the eggs were taken directly from the ovary of the cut animal and left crowded together in sea water in a Syracuse watch glass and immediately fertilized with the sperm of the other species, quantities of the eggs were fertilized. In the cross, *Strongylocentrotus purpuratus* ♀ × *S. franciscanus* ♂, the percentage, in one experiment, was 1 per cent. fertilizations with eggs prepared in the usual way (well washed and separated), and 80 per cent. with eggs direct from the ovary and crowded. The same batch of eggs, and sperm from the same male, were used in the experiment, and the counts were made of the blastulae just before swimming. The experiment was repeated with many different batches of eggs with the same general result; the reciprocal cross gave similar results, but the difference was not so great. In all cases, a large quantity of sperm was used, as it has long been known that over-insemination increases cross-fertilizations.

When the eggs of the sea urchin, *Strongylocentrotus purpuratus*, were crossed with the sperm of an entirely different genus, *Dendraster excentricus*, a sand-dollar, not a single fertilization was observed when the eggs were prepared in the usual way. But when the eggs from the same female were taken directly from the ovary and left crowded together and crossed with the sperm from the same *Dendraster* male, 10 per cent. of the eggs were fertilized.

Loeb has shown that increased alkalinity of the sea water favors cross-fertilizations. In the present case, we should expect an increased acidity due to the accumulation of CO<sub>2</sub> around the unwashed eggs. However, bubbling CO<sub>2</sub> through the sea water did not increase the percentage of cross-fertilizations; possibly the optimum CO<sub>2</sub> tension was not attained. Keeping the eggs for several hours slightly increased the percentage of cross-fertilizations. Sea water in which unfertilized eggs had been kept for 4 to 24 hours (at about 8° C. for the longer periods) gave a slightly higher percentage of cross-fertilizations for fresh eggs than did fresh sea water. It would certainly seem that some substance diffuses from the eggs which favors cross fertilizations and that this is present in effective quantity when the eggs are unwashed and crowded.

ETHEL BROWNE HARVEY

PRINCETON UNIVERSITY

## PHOSPHORESCENCE OF HUMAN TEETH

THE fluorescence of teeth is usually intense white with an occasional yellowish or greenish tinge. In older persons the fluorescence shifts into the longer wave-lengths, becoming reddish. Reddish fluorescence is also noted in the teeth of diseased persons.<sup>1</sup> A. H.

<sup>1</sup> J. A. Radley and J. Grant, "Fluorescence Analysis in Ultra-Violet Light." New York: D. van Nostrand Company, Inc., 1939.