

stance is harmless when ingested by animals before it can be used in food or feed. Since such information is not yet available for diphenylamine, it is not safe to make any practical application of the results of this study.

KENNETH T. WILLIAMS

EMANUEL BICKOFF

WALTER VAN SANDT

WESTERN REGIONAL RESEARCH LABORATORY,  
BUREAU OF AGRICULTURAL CHEMISTRY AND  
ENGINEERING, AGRICULTURAL RESEARCH  
ADMINISTRATION, U. S. DEPARTMENT OF  
AGRICULTURE, ALBANY, CALIF.

### EDEMA IN VITAMIN E DEFICIENT CHICKS<sup>1</sup>

INCREASING the concentration of soluble salts in the diet has been stated by Dam and Glavind<sup>2</sup> to be an effective means of increasing the severity of the edema which develops in chicks fed a diet low in vitamin E. Experimental results confirming this statement have been obtained in this laboratory and were reported at the Ninth Informal Poultry Nutrition Conference held in Boston on March 31, 1942. For example, in one experiment 100 per cent. mortality resulted in a group of 9 three-week-old chicks given daily doses of NaCl solution (20 gms/100 ml) during a six-day period. Eight of the 9 chicks showed some form of edema. One of 8 control chicks given no NaCl died during the same period but showed no evidence of edema. The group given NaCl solution received 0.017 ml/gm body weight on the first day of treatment and the dose was gradually increased, reaching 0.021 ml/gm body weight on the sixth day. All chicks were fed a vitamin E-low diet consisting of dextrinized corn starch 59, dried skimmilk 20, casein 13, butyl fermentation residue 3, oyster shell flour 1, calcium phosphate 2, sodium chloride 1 and cod liver oil 1. To each kg were added 120 mg  $\text{MnSO}_4 \cdot 4\text{H}_2\text{O}$ , 2 mg thiamine, 2 mg 2-methyl-1-4-naphthoquinone and 1 gm choline hydrochloride.

Administration of NaCl solution was first tried because chicks fed diets similar to that given above did not develop edema as readily as did chicks fed the diet previously described,<sup>3</sup> which contained 54 per cent. of dried skimmilk. It appears that disturbance of the osmotic equilibrium is necessary in addition to vitamin E deficiency to cause generalized edema.

Supplementation of the diet outlined above with 20 per cent. of a practical chick mash protected chicks against edema even when NaCl solution was administered. Lower levels of the practical mash were

not effective. These results were interpreted as an indication of the vitamin E content of this mash, but some doubt is cast on this interpretation by the report of Dam and Glavind that inositol and lipocaeic also protect against edema.

Experiments in this laboratory on the specificity of the protective effect of vitamin E were directed toward compounds having similar antioxidant properties. Toluhydroquinone, p-xyloquinone and trimethylhydroquinone were inactive when fed at four times the effective level of alpha-tocopherol. They are also ineffective when fed to rats.<sup>4</sup> Beta-tocopherol<sup>5</sup> was about half as potent as alpha-tocopherol, and this also conforms to the results of rat experiments.<sup>6</sup>

Pappenheimer, *et al.*,<sup>7</sup> in their extensive studies on encephalomalacia in chicks showed that the primary changes in this manifestation of vitamin E deficiency were edema and hemorrhage resulting from alteration of the capillary walls in the brain. It now appears that such alteration occurs in other parts of the vitamin E deficient organism but is not usually sufficient to permit exudation unless the normal osmotic relationships are disturbed. The question may well be raised as to the possible relationship of capillary changes to the various manifestations of vitamin E deficiency in other species.

H. R. BIRD

UNIVERSITY OF MARYLAND

### ISOMORPHISM AND ISOTYPISM AMONG SILICATES AND PHOSPHATES

RECENT advances in the study of isomorphism and isotypism may be illustrated through examples taken from compounds related to two mineral groups, the garnets and the apatites. Investigations by numerous chemists and mineralogists have revealed new isomorphic and isotypic relationships among silicates and phosphates, such as the isotypism of  $\text{AlPO}_4$  and  $\text{SiO}_2$  reported by Huttenlocher. However, a complete discussion is beyond the scope of these brief comments and complete references to the diverse sources of the data are purposely omitted.

Isomorphism, in the sense of the mineralogist, is characteristic of those compounds which exhibit stereochemical miscibility (so-called mixed crystals). Limited isomorphism obtains when the stereochemical miscibility is limited, whereas complete isomorphism exists when numerous intermediate members have been proven.

<sup>4</sup> C. Golumbic and H. A. Mattill, *Jour. Biol. Chem.*, 134: 535, 1940.

<sup>5</sup> Alpha and beta-tocopherol and trimethylhydroquinone were furnished by Merck & Co., Inc., Rahway, N. J., through the courtesy of Dr. D. F. Green.

<sup>6</sup> P. Karrer and H. Fritzsche, *Helv. Chim. Acta*, 22: 260, 1939.

<sup>7</sup> A. M. Pappenheimer, M. Goettsch and E. Jungherr, *Conn. Agr. Exp. Sta. Bul.*, 229, 1939.

<sup>1</sup> Scientific paper No. A36a, Contribution No. 1857 of the Maryland Agricultural Experiment Station (Department of Poultry Husbandry).

<sup>2</sup> H. Dam and J. Glavind, *SCIENCE*, 96: 235, 1942.

<sup>3</sup> H. R. Bird and T. G. Culton, *Proc. Soc. Exp. Biol. and Med.*, 44: 543, 1940.