

the heat increment is also shown by five series of experiments conducted at this laboratory (Forbes, *et al.*, *Jour. Nutr.*, 10 (1935), 461; 15 (1938), 285; 18 (1939), 47; 20 (1940), 47), with mature as well as with growing rats as subjects, in which the heat production of animals receiving equicaloric diets differ-

ing in protein content decreased slightly in the order of the increase in protein.

E. B. FORBES

R. W. SWIFT

INSTITUTE OF ANIMAL NUTRITION,
PENNSYLVANIA STATE COLLEGE

SCIENTIFIC APPARATUS AND LABORATORY METHODS

THE CHEMICAL CONTROL OF BERMUDA GRASS AND OF CROWFOOT GRASS

SODIUM CHLORATE^{1, 2} and cyanamide³ have been suggested for an eradication of Bermuda grass (*Cynodon dactylon* Pers.). The compounds have been used with a varying degree of success.⁴ Experiments with other herbicides seemed therefore indicated. As laboratory experiments⁵ had shown that ammonium sulfamate has a high toxicity for Bermuda grass, tests at roadside plots and at a tennis court were started with this salt as well as with calcium thiocyanate, as this compound had been used successfully in the eradication of nut grass.⁶

Series of experiments were started in October, January, February, July, August and September, so as to compare the influence of the various seasons. 0.5 to 3 liters of the solution were sprinkled per square meter, which contained from 200 to 1,000 plants. The calcium thiocyanate has always been used as a solution 1.25 molar in CNS⁻, while the concentration of the ammonium sulfamate varied from 0.5 to 2 molar. We are obliged to the American Cyanamide Company for the supply with calcium thiocyanate solution and to E. I. du Pont de Nemours and Company, Inc., for the ammonium sulfamate.

The control of the Bermuda grass has been completed within a week with as small an amount as 0.6 l. 1.25 m CNS⁻ per square meter if there was no rain in the first four days after the treatment. The plots remained free of weeds for three to six months after the treatment. However, the success of the treatment depends largely on the season. In dry weather the control was complete. If there was about 0.5 inch rain in the first few days after the application of the solution, complete control of the weed could still be reached by the use of 1 liter 1.25 m CNS⁻ per square meter. Heavier rains limited the eradication and could not be counteracted by higher doses of the herbicide. Also the length of time for which a complete control lasted was influenced by the rain, immigration of weeds, especially crowfoot grass (*Eleusine indica*)

from neighboring plots occurred much quicker in the rainy than in dry season.

Ammonium sulfamate killed the Bermuda grass in doses of 0.6–1 liter of molar solution completely in the dry season. The control lasted from 3 to 5 months. Weaker solutions were only partially effective. The rain affected its action more than that of the calcium thiocyanate; 0.5 inch rain in the first 4 days already reduced the control to about 90 per cent., stronger rains made it rather incomplete.

Some of the plots treated with calcium thiocyanate contained also a large number of crowfoot grass (*Eleusine indica*). Ada Georgia⁷ reports that carbolic acid can be used for its eradication, but otherwise little seems to be known about its chemical control. Its eradication by 1.25 molar CNS⁻ seemed much more difficult than that of Bermuda grass. 1 to 1.5 liter per square meter gave only a 50 to 80 per cent. control. The picture changed, however, when the treatment was preceded by a cutting of the grass. Then, 1 liter of 1.25 m CNS⁻ eradicated 90 to 100 per cent. of the grass within the first week, also heavy rains (more than 3 inches in the first 4 days) did not seriously interfere with the herbicidal action of the calcium thiocyanate.

Hence, it can be said that 1.25 m CNS⁻ or m ammonium sulfamate control Bermuda grass effectively if applied at the ratio of 0.6 to 1 liter per square meter in the dry season. 1.25 m CNS⁻ at the ratio of 1 liter per square meter will eradicate crowfoot grass only if the grass was previously cut.

F. FROMM

IRMA M. VIDAL

POLYTECHNIC INSTITUTE OF PUERTO RICO,
SAN GERMÁN, PUERTO RICO

⁷ A. Georgia, "Manual of Weeds," p. 53, Macmillan, 1930.

BOOKS RECEIVED

- BROWNE, CHARLES A. *A Source Book of Agricultural Chemistry*. Illustrated. Pp. 290. Chronica Botanica Co. \$5.00.
- CAIN, STANLEY A. *Foundations of Plant Geography*. Illustrated. Pp. xiv + 556. Harper and Brothers. \$5.00.
- SANDELL, E. B. *Colorimetric Determination of Traces of Metals*. Illustrated. Pp. xvi + 487. Interscience Publishers. \$7.00.
- ¹ Fred V. Grau, *C.A.*, 30: 6123³, 1936.
- ² *Agr. Jour. (Barbados)*, 7: 13, 1938.
- ³ Sturkie, *C.A.*, 32: 714⁷, 1938.
- ⁴ Robbin, Crafts and Warner, "Weed Control," p. 458, McGraw-Hill, 1942.
- ⁵ Fromm, *Ciencia y Técnica*, 1: 69, 1943.
- ⁶ Fromm, *SCIENCE*, 96: 337, 1942.