

study, using a ration in which 6 per cent. liver extract (1:20 paste) was included, rats fed butterfat made better, but insignificantly better (Fisher's "t" test)⁶ gains over those fed corn oil. The average results of both these studies for the male rats are summarized in Table 2. The results on the females were quite similar in significance.

These studies indicate that, apart from differences in vitamin content, corn oil and butterfat are essentially equal in growth-promoting value for the rat.

L. P. ZIALCITA, JR.

H. H. MITCHELL

DIVISION OF ANIMAL NUTRITION,
UNIVERSITY OF ILLINOIS

SCIENTIFIC APPARATUS AND LABORATORY METHODS

A TECHNIQUE FOR MOUNTING FREE-LIVING PROTOZOA

THE mounting of free-living protozoa on to microscope slides has always been a source of trouble to the protozoologist. The method of centrifuging after each stage in the processes of staining, dehydration, etc., suffers from several disadvantages. Firstly, the control of differentiation is difficult; for if over- or under-staining occurs, all the specimens in the tube are affected, and the whole must be re-treated. Secondly, since the organisms have to be centrifuged a considerable number of times in their passage through alcohols, stains, etc., it means that the cells are frequently distorted and true cytological pictures are not obtained. This is especially the case in dividing protozoa, where the protoplasm is less viscid than usual.¹

Several methods of fixing protozoa to slides have been suggested,² but are unsatisfactory for one reason or another. The following method, recently developed by the writer, fixes the protozoa very securely to the slide and is simple and effective to use in practice.

The organisms are fixed in Schaudinn, and brought through 70 per cent. and 90 per cent. into absolute alcohol by gentle centrifuging. A small drop of albumen is placed on a clean slide, and a very thin film produced by smearing it with the edge of another slide—exactly as in the preparation of a blood film. A drop of the concentrated organisms is allowed to fall on to the film of albumen from a fine pipette held about an inch above the slide. The combined action of the dropping force and the rapid coagulation of the albumen by the alcohol, immediately causes the organisms to be fixed securely to the slide. These slides are then placed in absolute alcohol, and treated as ordinary sections.

This method avoids the difficulties mentioned above; the small amount of centrifuging necessary in the preliminary concentrating never being sufficient to damage the cells. The film of albumen, too, is so thin that it causes no interference with the staining reactions of the protozoa.

J. D. SMYTH

⁶ F. E. Croxton and D. J. Cowden, "Applied General Statistics." New York: Prentice-Hall, Inc. 1941.

¹ J. B. Gatenby and J. D. Smyth, *Quart. Jour. Micr. Sci.*, 81, 1940.

ON QUIETING PARAMECIUM WITH METHYL CELLULOSE

MARSLAND¹ has suggested an excellent method of quieting *Paramecia* for study by the elementary student, using Dow "Methocel," or methyl cellulose in 10 per cent. aqueous solution.

We have found in our laboratory a slight modification of Marsland's method to be even more satisfactory for our purposes. Since 10 per cent. was a little too viscous, we tried 5 per cent., and we suggest this procedure: Make a small ring of 5 per cent. Methocel, slightly smaller than the cover glass to be used. Into the center of the ring place a small drop of medium containing *Paramecia*. Add a cover glass. Practice teaches one how much of each to use, but less than a full drop of each is often satisfactory with small cover glasses.

This enables the student to observe normal movement for a few minutes before diffusion of the methyl cellulose has slowed him down, and then progressively increasing viscosity gradually slows him to a completely stationary position. At this point he may be placed under an oil immersion objective, and ciliary motion studied in detail. Eventually even this slows down until the cilia appear to beat with great effort.

The Dow Company was very generous in furnishing us with the methyl cellulose.

RELIS B. BROWN

WESLEYAN COLLEGE,
MACON, GA.

² J. B. Gatenby and T. S. Painter, "Microtometist's Vade Mecum," London, 1937.

¹ Douglas A. Marsland, *SCIENCE*, 98: 2549, 414, November 5, 1943.

BOOKS RECEIVED

- LILLIE, FRANK R. *The Woods Hole Marine Biological Laboratory*. Pp. ix + 284. University of Chicago Press. \$4.00.
- OSBORNE, STAFFORD L. and HOLMQUEST, HAROLD J. *Technic of Electrotherapy*. Pp. xix + 780. Charles C Thomas, Publisher. \$7.50.
- RIEGLER, EMIL RAYMOND. *Chemical Machinery*. Pp. ix + 583. Reinhold Publishing Corporation. \$5.00.
- WEIMER, BERNAL R. and CORE, EARL L. *A New Manual for the Biology Laboratory*. Pp. vii + 213. John Wiley and Sons, Inc. \$2.00.