

TECHNICAL PAPERS

Synthetic Caffeine

CHARLES C. SCOTT, E. BROWN ROBBINS,
and K. K. CHEN

*Lilly Research Laboratories, Eli Lilly and Company,
Indianapolis, Indiana*

The identity of a synthetic compound with a natural product is usually made by the determination of physical constants such as the melting point, mixed melting point, and elementary analysis. However, with the many pharmacologically active substances, such as hormones and vitamins, a biological assay has been employed to supplement chemical identification. This appears important and necessary, because sometimes the physical constants do not completely establish the absolute purity of the product. Recently, for example, there was encountered in this laboratory a sample of synthetic pyridoxine which answered the specifications of *New and nonofficial remedies* (4), but which proved to be much more toxic than the standard.

Synthesis of caffeine and theobromine was initiated by Fischer and his co-workers (1), and newer methods of synthesis, partial or complete, have been achieved by other investigators (5). Previously, we studied the synthetic alkyl derivatives of theobromine (3). As an extension of the above work, it occurred to us that pharmacological proof of identity of totally synthetic caffeine and commercial caffeine made from a natural intermediate was desirable. The latter is usually prepared by methylation of theobromine. Samples of both were generously supplied by L. P. Kyrides, research director of the Monsanto Chemical Company, St. Louis, Missouri. The melting point of commercial caffeine ranged between 234.5 and 237° C.; that of synthetic caffeine, between 234.5 and 237.5° C.; and the mixed melting point, between 235 and 237° C.

Two types of experiments were carried out—one, the measurement of the stimulation of the central nervous system in rats; the other, the comparison of toxicity in mice by intravenous injection. The central stimulating action was evaluated in spring-suspended cages according to the method of Schulte and co-workers (2). Ten rats, weighing between 230 and 280 grams, were employed. A fixed dose of 20 mg./kg. of each drug in 0.5 per cent solution was injected subcutaneously. The observation period was exactly 4 hours, counting from the time of injection. Five animals received synthetic caffeine, and the other 5, the commercial. On the following day a cross-over test was made—those receiving the synthetic caffeine the previous day received the commercial, and vice versa.

For the toxicity study, mice weighing from 14.6 to 19.3 grams were used. The concentration of the solution was the same as for rats (0.5 per cent). Both samples were run on the same day.

The results are unequivocal. The average number of revolutions recorded from the cages by the 10 rats for commercial

caffeine was 34.8; for synthetic caffeine, 34.1; and that of the control test (without medication), 5.1. The median lethal dose (LD₅₀) by intravenous injection of commercial caffeine in 30 mice was found to be 79.36 ± 6.94 mg./kg. The LD₅₀ of synthetic caffeine in 30 other mice was determined to be 75.51 ± 5.39 mg./kg. There is no significant difference between the two figures statistically.

Pharmacological results indicate, therefore, that commercial caffeine and synthetic caffeine have the same degree of central stimulation and toxicity.

References

1. FISCHER, E. *Ber. Dtsch. Chem. Gesellsch.*, 1882, 15, 453; 1897, 30, 2400; 1899, 32, 435; FISCHER, E., and BROMBERG, O. *Ibid.*, 1897, 30, 219; FISCHER, E., and ACH, F. *Ibid.*, 1898, 31, 1980.
2. SCHULTE, J. W., REIF, E. C., BACHER, J. A., JR., LAWRENCE, W. S., and TANTER, M. L. *J. Pharm. exp. Therap.*, 1941, 71, 62.
3. SCOTT, C. C., and CHEN, K. K. *J. Pharm. exp. Therap.*, 1944, 82, 89; SCOTT, C. C., ANDERSON, R. C., and CHEN, K. K. *Ibid.*, 1946, 86, 113.
4. SMITH, A. *New and nonofficial remedies*. Chicago: American Medical Association, 1945. P. 620.
5. TRAUBE, W. *Ber. Dtsch. Chem. Gesellsch.*, 1900, 33, 3035; SCHMIDT, E., and PRESSLER, H. *Liebig's Ann. Chem.*, 1883, 217, 287; ENGELMANN, M. *Ber. Dtsch. Chem. Gesellsch.*, 1909, 42, 177; FIORE, G. *Boll. chim. farm.*, 1915, 54, 609; BLITZ, H., and DAMM, P. *Liebig's Ann. Chem.*, 1917, 413, 186; ROSENEMUND, K. W., and ZETZSCHE, F. *Ber. Dtsch. Chem. Gesellsch.*, 1918, 51, 578.

Visibility of the Deer Fly in Flight¹

ELEK LUDVIGH

*Howe Laboratory of Ophthalmology,
Harvard University Medical School*

Visual acuity while viewing a moving object is a matter which has received little attention. In fact, the only reference bearing remotely on this subject which I have been able to find is a note by Irving Langmuir (1). Dr. Langmuir has ingeniously demonstrated that the speed of 818 miles/hour attributed to the deer botfly by the *Illustrated London News* (4) is excessive. As Dr. Langmuir expresses the matter, the insect must consume "1.5 times his own weight of food each second" in order to maintain this velocity.

To establish the approximate speed of the fly, Dr. Langmuir conducted an experiment which he describes as follows:

"... I took a short piece of solder about 1 cm. long and 0.5 cm. diameter and tied it about its middle to one end of a light silk thread, holding the other end in my hand. With lengths of thread of from 1 to 3 feet it was easily possible to swing the weight in a circle in a vertical plane at the rate of 3 to 5 rotations per second (timed with a telechron clock).

¹ This investigation was supported in part by a grant from the American Optical Company.