

The conclusion is drawn that the methionine can be metabolized to α -aminobutyric acid and that it is probably the main source of the latter, since it does not, of course, occur in the protein of the diet. The fate of the carbon chain of the former now seems clear. A reinvestigation of the possible role of α -aminobutyric acid in the body is indicated. If its presence is essential, methionine may have to be wasted in order to produce it, and, in that case, giving it to animals on a low methionine diet may exert a methionine-sparing action.

Methionine sulfoxide may be of significance in oxidation-reduction potential. It can oxidize cysteine to cystine *in vitro* (2). Methionine, on standing under various conditions, readily changes largely into the sulfoxide and may therefore act as an oxygen carrier. In view of this easy oxidizability, however, its presence in the urine after methionine feeding needs more careful checking. It may have been formed by aerial oxidation. The validity of procedures for the determination of methionine in urine by oxidation reactions (H_2O_2 , etc.) also arises.

Further details of these findings, which arose out of an investigation into the aminoaciduria in Fanconi syndrome, have been submitted to the *Biochemical Journal*.

References

1. DENT, C. E. *Biochem. J.*, 1946, **40**, xlv.
2. TOENNIES, G., and KOLB, J. J. *J. biol. Chem.*, 1939, **128**, 399.

Failure to Produce Neoplasms in Rats by Feeding Heated Wheat-Germ Oil

CHARLES HOFFMAN and GASTON DALBY

Ward Baking Company, New York City

K. GEORGE FALK and GRACE MCGUIRE

Laboratory of Industrial Hygiene, New York City

The announcement by Rowntree (2) in 1937 of the occurrence of neoplasms in Wistar rats following the ingestion of crude wheat-germ oil created considerable interest. The suggested relationship to diet as well as the production in experimental animals of malignant lesions of the gastrointestinal tract were factors of scientific importance. Attempts to duplicate these results were made in many laboratories, unfortunately without success (1). Since one of us (G. D.) had prepared some of the original wheat-germ oil used by Rowntree in his successful experiments, an attempt was made to duplicate his work, using oil prepared by the same operator and under the same laboratory conditions. Rowntree's diet was followed in detail, and Wistar rats were fed for almost a year. No neoplasms were produced (unreported work).

Some recent evidence has suggested that heated fats may contain carcinogenic factors possibly due to the conversion of sterols to carcinogens by heat (3). Wheat-germ oil is rich in sterols, and Rowntree, in the preparation of his diet, actually might have changed some of the sterols in the wheat-germ oil to carcinogens, since the oil was usually heated again before use to insure that no trace of ether smell remained.

In order to test this possibility, crude wheat-germ oil was prepared exactly as it had been previously. It was then

heated at 275° C. for two hours. Three liters of heated oil were mixed with 10 kg. of basic diet in accordance with Rowntree's directions. Thirteen female Wistar rats with an average weight of 64 grams and 14 male Wistar rats with an average weight of 65.5 grams were started on the diet in November 1945. The animals grew somewhat slowly, compared to colony rats, but the growth rate was steady and the animals appeared healthy. There were a number of deaths due to colds and pneumonia. The surviving animals were killed and autopsied in July 1946. Seven males averaging 253 grams and 5 females averaging 213 grams survived. The condition of all animals was excellent, with no suggestion of any malignant lesions.

This evidence would seem to indicate that the sterols in wheat-germ oil are not converted to carcinogens by heat. The heat conversion of sterols to carcinogens was apparently not a factor in Rowntree's production of gastrointestinal lesions in the rat.

References

1. BRUES, A. M., MARBLE, B. B., and RIEGEL, B. *Cancer Res.*, 1941, **1**, 815-817.
2. ROWNTREE, L. G., STEINBERG, A., DORRANCE, G. M., and CICCONE, E. *F. Amer. J. Cancer*, 1937, **31**, 359-372.
3. ———. *Nutr. Rev.*, 1945, **3**, 42-44. (Unsigned review.)

The Occurrence of Two Fertile Florets in the Spikelets of *Chloris*

MILDRED P. MAULDIN

Southwestern Seed Service, Waco, Texas

During the writer's experience in seed analysis Rhodes grass (*Chloris Gayana* Kunth) was frequently received for purity analysis and for germination test. Early examination of spikelets of this species disclosed that many contained a fertile pedicellate floret in addition to the fertile basal and sessile floret. The presence of caryopses was the criterion employed.

Thus far no confirmation of this observation has been found in the available grass and general floras examined. The seed analyst is perhaps the person best situated to make such an observation, since each of his many routine analyses requires the examination of thousands of individual seeds or seed units, but his usually elementary knowledge of taxonomy might lead him to overlook the significance of such a fact.

Descriptions of the genus *Chloris*, as far as the writer has been able to determine, appear to be unanimously in agreement on the occurrence of one fertile floret in the spikelet, and this the basal, sessile floret. For example, Hitchcock (1) states: "Spikelets with 1 perfect floret, sessile, . . . the rachilla disarticulating above the glumes, produced beyond the perfect floret and bearing 1 to several reduced florets consisting of empty lemmas. . . ." In no instance was there more than one floret in the spikelet described as fertile in the literature available. Silveus' description (3) is worded identically, and those in other floras are either essentially similar or identical.

As the average purity, or seed set of *C. Gayana* by weight, was found to be approximately 25 per cent (37 lots) (2), it may be assumed that the number of pedicellate florets matur-