

The Inhibition of Virus Increase by Malachite Green

WILLIAM N. TAKAHASHI

Division of Plant Pathology,
University of California, Berkeley

Since the reproduction of plant viruses has not been demonstrated outside of the living cell and purified plant viruses have not been proved to possess enzymic activity, it would appear that the processes involved in the reproduction of viruses are intimately associated with the metabolism of the host cell. Physical conditions and chemical compounds that affect host metabolism are likely to influence virus formation. There is a possibility that, by the use of enzyme blocks or inhibitors, enzyme systems indispensable for virus formation may be identified. Since the inhibition must take place *in vivo*, an inhibitor must not only possess a certain degree of specificity but also be active at a concentration nonlethal to the living tissue. Malachite green, a diaminotriphenylmethane dye, seems to possess these requirements. The inhibition of dehydrogenase activity (oxidation of glucose, lactate, succinate, and formate) of *B. coli* by malachite green was noted by Quastel and Wheatley (4); the ability of malachite green to penetrate living cells was shown by Albach (1); and Caldwell and Meiklejohn (2) found that this dye affected the respiration of tobacco stem tissue up to a dilution of 3.3 ppm.

By using a detached leaf tissue technic (Takahashi, 5), the effect of low concentrations of malachite green on the increase of tobacco mosaic virus in leaf tissue has been studied. In these experiments 1-gm strips from infected half-leaf pairs were used, one set floated on a solution of malachite green and the other on distilled water. The experiments were conducted in the dark and terminated after 8 days of culture. In order to determine the extent of virus formation under these conditions, the tissues were homogenized, diluted 1 part in 20 with distilled water, and inoculated on 20 opposite halves of *Nicotiana glutinosa* leaves (Holmes, 3).

Considerable inhibition of virus increase in this cultured leaf tissue was shown at concentrations of malachite green as low as 2 ppm, and more at 4 ppm. The total number of local lesions on 20 half-leaves of *Nicotiana glutinosa* for leaf tissue on 2 ppm was 70, and for the control on distilled water, 312. At 4 ppm the totals were 46 for treated and 1,068 for control. At the concentrations used in the above experiments malachite green has practically no effect on the extracted virus *in vitro*. A treated suspension of virus (4 ppm of malachite green) gave 1,890 local lesions, and the control, 2,060.

The inhibition of virus increase in living cells by low concentrations of malachite green appears to be due to its action as a block in enzyme reactions leading to virus formation. By the use of a series of more specific inhibitors there is a possibility of revealing some of the steps involved in virus formation. Meanwhile, the possi-

bility that malachite green may be of some value in the chemotherapy of virus diseases must not be overlooked.

References

1. ALBACH, W. *Z. Wiss. Mikrosk.*, 1927, **44**, 333-334.
2. CALDWELL, J., and MEIKLEJOHN, JANE. *Ann. Bot.*, 1937, **3**, 477-498.
3. HOLMES, FRANCIS O. *Bot. Gaz.*, 1929, **87**, 39-55.
4. QUASTEL, J. H., and WHEATLEY, A. H. M. *Biochem. J.*, 1931, **25**, 629-638.
5. TAKAHASHI, W. N. *Amer. J. Bot.*, in press.

Stability of Solutions of Pure Ascorbic Acid and of Dehydroascorbic Acid

LOUISE P. GUILD, ERNEST E. LOCKHART,
and ROBERT S. HARRIS

Nutritional Biochemistry Laboratories,
Massachusetts Institute of Technology

Results obtained in our laboratories, as well as those of other investigators (4, 7), have indicated a lack of agreement between the 2,6-dichlorophenol-indophenol method of Bessey, *et al.* (1, 6) and the 2,4-dinitrophenylhydrazine method of Roe, *et al.* (9, 10) in the estimation of ascorbic acid in foods. Therefore, these two methods have been compared as to their ability to measure the ascorbic or dehydroascorbic acid content of freshly prepared and aged solutions. A study has also been made of the effects of oxalic and metaphosphoric acids on the stability of ascorbic and dehydroascorbic acids in solution.

TABLE 1
STABILITY OF ASCORBIC ACID AND DEHYDROASCORBIC ACID IN
0.5% OXALIC ACID AS MEASURED BY THE ROE
AND BESSEY METHODS

Days stored at 4° C	Ascorbic acid solution μg/ml		Dehydroascorbic acid solution μg/ml	
	Roe	Bessey	Roe	Bessey
0	5.0	5.0	5.0	5.0
2			5.0	3.5
4	4.9		5.1	
6				2.4
8	5.1		5.4	
10	4.8		5.2	1.9
14		4.9		1.2
15	4.6		5.2	
22				0.8
24			5.1	
32		4.8		0.8

A stock solution containing 1,000 μg of pure ascorbic acid/ml of 0.5% oxalic acid was prepared. A working standard solution containing 100 μg of ascorbic acid/ml was obtained by diluting 10 ml of this solution with 10 ml of acetic acid (as required by the Roe method, 9) and making up to a volume of 100 ml with 0.5% oxalic acid. A portion of this solution was treated with Norite to oxidize the ascorbic acid to dehydroascorbic acid (9). Both this dehydroascorbic acid solution and the remaining untreated ascorbic acid solution were stored at 4° C.