

The observed pH of Schiff's reagent prepared with thionyl chloride is 1.24; that of Schiff's reagent prepared in the usual manner is 1.38. This slight difference in acidity does not appear to be significant.

Schiff's reagent prepared with thionyl chloride has been successfully used as a nuclear stain on tissue sections of human thymus gland, kidney, liver, and spleen, as well as on the fungi, *Blastomyces dermatitidis* and *Saccharomyces cerevisiae*.

References

1. COLEMAN, L. C. *Stain Technol.*, 1938, **13**, 123-124.
2. FEULGEN, R., and ROSSENBECK, H. *Z. physiol. Chem.*, 1924, **135**, 203-248.
3. SHRINER, R. L., and FUSON, R. C. *The systematic identification of organic compounds*. (2nd ed.) New York: Wiley, 1940. P. 62.
4. WIELAND, HEINRICH, and SCHEUING, GEORG. *Ber. deutsch. chem. Ges.*, 1921, **54**, 2527-2555.

Plant Virus for Electron Microscopy

JAMES JOHNSON¹

*Departments of Horticulture and Plant Physiology,
University of Wisconsin*

The uniformity of size of an infectious unit has a significant bearing on the purity of a culture when the causative agent of a disease is to be established. Considerable variability in the size, particularly in the length, of tobacco mosaic virus particles, as seen in the electron microscope, has been reported. Oster and Stanley (5) have supported earlier conclusions with new measurements of virus obtained directly from leaf hairs of infected plants. Sigurgeirsson and Stanley (6) have also measured the virus particles in freshly expressed juice and found them more uniform than those in older or in centrifuged material. Later results (3, 4) show that the greatest number of virus particles (20-40%) fall into a group size of $15 \times 280 \mu\mu$, and that less than 5% are extraordinarily short or long. It is emphasized by these workers that much of the variation in length is attributable to aggregative or disruptive forces occurring during preparation of the virus concentrate for examination under the electron microscope.

In connection with certain virus studies made in this laboratory, it has been found that the application of direct water pressure to plants systemically infected with mosaic yields clear exudation drops containing a sufficient virus concentration to be observed when applied to the electron microscope screens. The approximate concentration of individual droplets may be checked by wiping drops on a local lesion host with the tip of a finger or a glass spatula. This exudate contains some extraneous material; but, with the gold shadow-casting process, the virus particles were easily found in considerable numbers. In the case of plants with hydathodes,

¹ Aided by a grant from the Wisconsin Alumni Research Foundation. The writer is also indebted to Luther Preuss for cooperation in operating the electron microscope and the gold shadow-casting equipment.

the virus may be exuded from these natural openings, but with other plants the droplets were collected chiefly from the cut ends of veins.

This method of obtaining a relatively pure form of virus was described from this laboratory in 1924 (1). The method has also been illustrated in other papers in connection with water-congestion studies (2). Where larger quantities of virus are required, as for purposes of concentration by differential centrifugation, a more recent modification has consisted of collecting the exuded drops in a 14" metal side-arm funnel properly placed under the plant foliage. In this manner 25-50 cc may be collected in a few hours from suitable plants, especially if the leaf blades are cut so as to expose the ends of numerous veins. The virus concentration of this exudate is not usually as high as that secured by the usual procedure of expressed juice. However, practically all of the extraneous plant material that interferes with electron microscopy is eliminated. With the water-pressure method, the exuded droplets may be applied directly to the collodion film of the electron microscope screen. Drying in a desiccator, followed by gold shadow-casting, completes the shortened preparation for electron microscope examination. Several strains of the tobacco mosaic virus particles have been collected in this manner, and different host plants, especially tobacco, tomato, *Eryngium aquaticum* L., and *Digitalis lanata* L. (a previously unreported host species), have been used. The method is applicable to glaucous plants such as *E. aquaticum* and *D. lanata*, where the leaf-hair method would not be possible.

The pressure exudate method of preparation of the virus for electron microscopy should not cause aggregation or disruption of particles of the infectious unit, especially when the first drops are used, since the actual pressure at the point of exudation is very low. Variation from the normal virus length in the droplets, if such occurs, would necessarily seem to arise from the drying process on the screen. The latter supposition is discounted by those who have found the virus particle relatively uniform in length (3). Measurements of the virus particles obtained by the water-pressure method have not yet been made. The electron micrographs, however, show a high variability in the length of the particle, not approaching, for example, the uniformity exhibited by most cultures of microorganisms.

This method of obtaining virus may prove useful for other studies by means of electron microscopy.

References

1. JOHNSON, JAMES, and MULVANIA, M. *Science*, 1924, **60**, 19.
2. JOHNSON, JAMES. *Wis. agric. exp. Sta. Res. Bull.* **160**, 1947, 1-35.
3. KNIGHT, C. A., and OSTER, G. *Arch. Biochem.*, 1947, **15**, 289-294.
4. OSTER, G., KNIGHT, C. A., and STANLEY, W. M. *Arch. Biochem.*, 1947, **15**, 279-287.
5. OSTER, G., and STANLEY, W. M. *Brit. J. exp. Path.*, 1946, **27**, 261-265.
6. SIGURGEIRSSON, T., and STANLEY, W. M. *Phytopathology*, 1947, **37**, 26-38.