

main. The ultraviolet absorption spectrum (Fig. 2) closely resembled that of kynurenine, but it gave a negative Otani's kynurenine reaction (5). The ninhydrine, xanthoprotein, and Ehrlich's diazo reactions were positive. The alkaline solution was of a deep-yellowish color and gave a jasminelike odor when heated. Dilute  $\text{KMnO}_4$  solution also changed the color to deep yellow (urochromogen test of Weiss). The Folin-Denis' uric acid reagent changed the solution to a greenish, and then to a bluish, color. The murexide and lumiflavin tests were negative.

Biological tests employing the mutants vermilion and cinnabar of *Drosophila* were positive.

**Addendum:** In a previous paper in this journal (*Science*, 1950, 111, 608) concerning the tryptophan pigments, which dealt with the epidermis pigments of the yellow mutant "lem" of *Bombyx mori*, we reported that the yellow pigment was xanthopterin, but, according to direct comparison with a sample of xanthopterin (kindly sent by E. L. Rickes, Merck & Co., Inc.) it has been found that, although this pigment resembles xanthopterin remarkably closely, it is slightly different. It has been named xanthopterin-B (B from *Bombyx*), and 2 other new pterins have also been isolated (leucop-terin-B and a 6-dehydroxyleucopterin derivative) (6).

#### References

1. BEADLE, G. W. Private communication.
2. HASKINS, F. A., and MITCHELL, H. K. *Proc. Nat. Acad. Sci.*, 1949, **35**, 500.
3. KIKKAWA, H. *Rep. Sericultural Exp. Sta.* (in Japanese), 1943, **11**, 311.
4. MITCHELL, H. K., and NYC, J. F. *Proc. Acad. Sci.*, 1948, **34**, 1.
5. OTANI, S., NISHINO, N., and IMAI, K. *Z. Physiol. Chem.*, 1941, **270**, 60.
6. *Bull. Chem. Soc. Japan* (in English). In press.

### Built-up Poultry-House Litter as a Growth-promoting Supplement for Chicks on an All-Vegetable Vitamin $\text{B}_{12}$ -deficient Diet

E. R. Halbrook,<sup>1</sup> A. R. Winter, and T. S. Sutton

Departments of Poultry Husbandry and Agricultural Biochemistry,  
The Ohio State University, Columbus

Extensive trials by Kennard and Chamberlin (1, 2) at the Ohio Agricultural Experiment Station demonstrated that poultry-house floor litter which has accumulated (built up) for a long period of time gives better chick growth on all-vegetable diets than litter which is completely changed at frequent intervals. Reports have also been made regarding improved hatchability of eggs from hens on such litter (3, 8). It has been found that poultry droppings contain the "animal protein factor" (APF) (?), and that incubation of hen feces for 72 hr at 30° C stimulates further synthesis of this factor (4).

In a series of studies dealing with the synthesis of vitamin  $\text{B}_{12}$  (now recognized as being an important part

<sup>1</sup> Present address: Montana State College, Bozeman.

of the animal protein factor) by microorganisms in poultry-house litter, marked stimulation of growth was secured by adding built-up litter to an all-vegetable vitamin  $\text{B}_{12}$ -deficient diet fed to S. C. White Leghorn chicks in a wire-floored battery brooder.

The basal diet used was one consisting principally of soybean oil meal and ground yellow corn supplemented with the known requirements for vitamins, minerals, and the amino acid methionine. The diet was used for studies of soybean oil meal and animal protein factor by workers at the Purdue Agricultural Experiment Station (5, 6).

Vitamin  $\text{B}_{12}$  was shown to be the principal deficiency in the basal diet by the marked stimulation of growth secured by intramuscular injection of a saline solution of crystalline vitamin  $\text{B}_{12}$  (Cobione)<sup>2</sup> given in the amount of 0.3  $\mu\text{g}$  per chick weekly. This injection was as effective in stimulating chick growth as feeding the crude APF supplement<sup>2</sup> containing the equivalent of 20  $\mu\text{g}$  of vitamin  $\text{B}_{12}$ /kg of diet.

As shown in Table 1, addition of 1%, 2.5%, and 5%

TABLE 1  
EFFECT ON CHICK GROWTH OF ADDING BUILT-UP POULTRY-HOUSE LITTER AND OTHER SUPPLEMENTS TO AN ALL-VEGETABLE VITAMIN  $\text{B}_{12}$ -DEFICIENT DIET

| Diet  | No. Trials | Total No. chicks at start | No. died | Avg 6-wk wt in g (all trials) | Feed per g gain |
|---|------------|---------------------------|----------|-------------------------------|-----------------|
| Basal vitamin $\text{B}_{12}$ -deficient diet | 3          | 53                        | 8        | 345                           | 3.78            |
| Basal + APF supplement*                       | 3          | 53                        | 4        | 429                           | 3.23            |
| Basal + vitamin $\text{B}_{12}$ injections†   | 1          | 15                        | 1        | 426                           | 3.02            |
| Basal + 2.5% ground corncobs                  | 1          | 15                        | 1        | 250                           | 5.16            |
| Basal + 5% ground corncobs                    | 1          | 15                        | 1        | 265                           | 4.85            |
| Basal + 1% unautoclaved built-up litter       | 1          | 18                        | 0        | 419                           | 3.04            |
| Basal + 1% autoclaved built-up litter         | 1          | 18                        | 1        | 394                           | 3.28            |
| Basal + 2.5% unautoclaved built-up litter     | 3          | 53                        | 3        | 390                           | 3.29            |
| Basal + 2.5% autoclaved built-up litter       | 2          | 38                        | 0        | 466                           | 2.82            |
| Basal + 5% unautoclaved built-up litter       | 2          | 27                        | 3        | 412                           | 3.34            |
| Basal + 5% autoclaved built-up litter         | 2          | 35                        | 0        | 493                           | 2.55            |

\* Twenty  $\mu\text{g}$  vitamin  $\text{B}_{12}$ /kg of diet.

† Crystalline vitamin  $\text{B}_{12}$  (0.3  $\mu\text{g}$ /chick weekly).

levels of screened unautoclaved built-up corncob litter (over 1 year old) to the basal diet gave increased chick growth to 6 weeks of 74, 45, and 67 g, respectively, as compared to 84 g for the added APF supplement fed at the rate of 20  $\mu\text{g}$  of vitamin  $\text{B}_{12}$ /kg of diet. The built-up litter used for chick feeding was accumulated from hens that had been fed an all-vegetable vitamin

<sup>2</sup> We are indebted to Merck & Co., Inc., Rahway, N. J., through the courtesy of D. F. Green, for the crystalline vitamin  $\text{B}_{12}$  (Cobione) and the APF supplement (crude vitamin  $\text{B}_{12}$ ) used in these studies.