

hormones containing the α,β unsaturated ketone structure also contain an enzyme that will oxidize the Δ^5 -3-ol structure to the conjugated form in the presence of DPN as a hydrogen acceptor. This enzyme would appear fundamental in the biological synthesis of these important hormones.

References

1. DAVID, K., et al. *Z. physiol. Chem.*, **233**, 281 (1935).
2. TAGMANN, E., PRELOG, V., and RUZICKA, L. *Helv. Chim. Acta*, **29**, 440 (1946).
3. ALLEN, W. M., and WINTERSTEINER, O. *Science*, **80**, 190 (1934).
4. BUTENANDT, A., WESTPHAL, U., and HOHLWEG, W. *Z. physiol. Chem.*, **227**, 84 (1934).
5. MASON, H. L., MYERS, C. S., and KENDALL, E. C. *J. Biol. Chem.*, **116**, 267 (1936).
6. REICHSTEIN, T. *Helv. Chim. Acta*, **20**, 953 (1937).

Morphology of the Ovary of *Oryza Sativa* Linn. Var. *Plena* ("Double Rice")

R. M. Datta and A. K. Paul¹

*Jute Agricultural Research Institute,
Hoogly, West Bengal, India*

The ovary of *Oryza sativa* Linn. is superior, one-celled and contains a single ovule. In the variety *Plena* of the same species Prain (1) has recorded that there are as many as 7 ovaries in some specimens, the usual number of ovaries found being 2-5. Neither he nor any other worker has mentioned the number of ovules present in each such ovary. S. K. Mukherjee, Curator of the Herbarium, Indian Botanic Gardens, Sibpur, Calcutta, writes from the notes available in the Herbarium thus:

After the stamens wither, it is most usual to find that only two ovaries continue to develop and then not infrequently one of these fails to grow as fast as the other; but very often both grow equally, and the result is the "Double Rice." In this case the inner faces of both the grains are flat with a whitish vertical central band, and on section, the embryo of each is found at the outer, or glumal, aspect of the base of the grain. In a few cases three grains are developed, and then instead of having flat faces, they meet in the centre at the white line already mentioned, this line being at the apex of an obtuse angle; the embryo is in each case at the outside, as before.

In the course of our study on this variety, we have observed in several cases the presence of 2 ovules in a single ovary, the number of ovaries being the same as mentioned by Prain (1). These ovules grow to maturity side by side. They have 2 integuments each. More than one archesporial cell has also been observed (Fig. 1). Morinaga and Fukushima (2) found 2 ovules in an ovary of a haploid rice plant (*O. sativa*) and several ovules with 2 embryo-sac mother cells. In *Inapostol*, a Philippine variety of *O. sativa*, Juliano and Aldama (3) also recorded a two-celled archesporium in addition to the normal one-celled. They, however, recorded that only one of them becomes

¹ We offer our sincere thanks to S. K. Mukherjee for supplying us with the notes at the disposal of the Gardens.




FIG. 1. Section of the ovary of "double rice" showing 2 ovules with two-celled archesporium in one of them. ($\times 475$.)

functional. Kuwada (4) also found a single instance of a two-celled archesporium in a Japanese variety of *O. sativa*.

Hence, "double rice" is not only due to the presence of 2 or more ovaries, each containing a functional ovule, but also due to the presence of more than one ovule, generally 2, in each such ovary—a fact not recorded before. Fuller details will be published later.

References

1. PRAIN, SIR D. *Bengal Plants*, Vol. II, 1184 (1903).
2. MORINAGA, T., and FUKUSHIMA, E. *Japan. J. Botany*, **7**, 73 (1934).
3. JULIANO, J. B., and ALDAMA, M. J. *Philippine Agr.*, **26**, 1 (1937).
4. KUWADA, Y. *Botan. Mag. Tokyo*, **24**, 267 (1910).

A Root Disease of Plants Caused by a Nematode of the Genus *Trichodorus*

J. R. Christie and V. G. Perry

*Bureau of Plant Industry, Soils and Agricultural
Engineering, USDA, and Florida Agricultural
Experiment Station, Sanford*

A root disease of plants that seriously hampers the growing of certain vegetable crops in Florida is caused by a nematode of the genus *Trichodorus* Cobb, 1913. Stubby-root and the stubby-root nematode are suggested as common names for this disease and the causal organism, respectively. Thorne (1) places the genus *Trichodorus* in the family Diphtherophoridae of the superfamily Dorylaimoidea. No other member of this superfamily is known to be a serious plant pest, the only ones generally regarded as probably feeding on and injuring the roots of plants being species of *Xiphinema*.

Trichodorus primitivus (deMan 1876) Micoletzky, 1922, was collected first in the Netherlands by deMan, who named and described it (2). Subsequently, nematodes that Thorne believed to be the same species were collected by him near Fort Collins, Colo., near San Luis Obispo, Calif., and at various points in Utah, and by Cobb (3) near Arlington, Va. The stubby-root nematode is closely related to *T. primitivus*, but because there appears to be doubt in the minds of some taxonomists regarding its specific identity it is herein designated *Trichodorus* sp., pending further study.

Stubby-root has been produced experimentally under conditions sufficiently well controlled to provide