

tain the desired humidity in the moist chamber, and the yarn may be reused after sterilization.

WILLIAM A. FEDER

Ornamentals Laboratory, Cornell University
Farmingdale, Long Island

Dissa and Data

FOR nearly 20 years I have recommended to students my opinion that (1) the word "data" should be related to a plural verb form, as "the data are;" and that (2) the word "disinterested" means impartial or unbiased.

If recent publications are evidence of common usage, "data" can be used with either singular or plural verb form. It appears that this is a result of "growing pains" of our language. Are we all agreed to accept?

"Disinterested" has been often used recently in place of "uninterested." To this I object—probably ineffectually. "Disinterested" was a useful word, and I do not like losing it.

Shall I continue my former practices of instruction, or shall I stop being a bigot and forget?

S. REID WARREN, JR.

The Moore School of Electrical Engineering
University of Pennsylvania

EDITORIAL NOTE: *The editors subscribe to Dr. Warren's bigotry and will continue to correct these errors whenever they are made in manuscripts. They also object to the use of "presently" for "now," and "while" for "whereas" or "although." They are also disturbed because of (not "due to") the adverbial misuse of the adjectival expression "due to."*



Book Reviews

Taxonomy of Vascular Plants. George H. M. Lawrence. New York: Macmillan, 1951. 823 pp. \$7.95.

Modern botanical research has repeatedly substantiated the old taxonomic practice of treating the vascular plants as a major unit. Although many manuals and floras contain descriptions of all vascular plants found in a given area, they are rarely treated together in textbooks of systematic botany. Actually, *Taxonomy of Vascular Plants* is the first modern textbook of this kind and is eloquent testimony of the present tendency toward dealing with all vascular plants under the name Tracheophyta. Yet the author deliberately adopted the last published version (1936) of the widely used Engler system of classification, with its obvious shortcomings, as the basis for the systematic section of the book (Part II, pp. 333-72), because it is still the most carefully elaborated system available.

Other conspicuous changes in comparison with existing texts of systematic botany are seen in the elimination of all floral diagrams and formulas, the replacement of chapters dealing with the organography of vascular plants by an illustrated glossary of taxonomic terms (Appendix II, pp. 737-75), the introduction and consistent use of the term "taxon" (taxa), and the consolidation in Part I (pp. 1-331) of 14 chapters on "Principles and Practices of Plant Taxonomy." Appendix I (pp. 733-36) represents a "Suggested Syllabus for an Elementary Course in Taxonomy" for those who wish to use the book as a text in a one-term course. This syllabus proves that it is much more than an elementary textbook in both scope and contents. Thus it is not only the most inclusive textbook of systematic botany in English (or any other language) but also a convenient and indispensable reference work for the advanced student. The

latter will find in it well-balanced discussions of all major controversial aspects of phylogeny, along with informative chapters on field and herbarium techniques and other important principles and practices of taxonomy currently in use. The same is true of the systematic part with its enumeration of 264 families of vascular plants "known to grow as indigens or exotics in North America north of Mexico." The account of each family includes a technical description, enumeration of important genera, distributional data, discussion of morphological characteristics and assumed phylogenetic relationships, key references, and representative figures, many from L. H. Bailey's *Manual of Cultivated Plants* (1949). Completely extinct groups like the Pteridospermae are excluded.

A few interesting details may be singled out for comment. Under Ginkgoaceae five of the seven references deal with the spelling of the generic name *Ginkgo*, which should be corrected to *Ginkyo*. The future alone will tell how soon and how widely this spelling will be accepted. It is regrettable, however, that so much attention is being given to a problem of nomenclature when this important taxon is in such dire need of a synoptic treatment of its fossil forms. The recently proposed family Sarcopodaceae, here provisionally listed under the Gnetales (p. 368), has been rescinded, now that the genus *Sarcopus* has been identified with *Exocarpus* (Santalaceae). The Compositae are regarded as the largest family, with 950 genera and 20,000 species, thus rivaling or exceeding the Orchidaceae, here credited with 450 genera and 10,000-15,000 species, but considered to be the largest family by other authorities. Most likely, both families are larger than the remainder, containing numerous species, many of which may prove to be referable to others once critical studies of large genera are carried out.

In the light of the contents, fine typography, and illustrations, the price of this scholarly book is moderate, enabling taxonomists to own copies. Botanists in need of taxonomic information will find it an invaluable guide and source.

THEODOR JUST

Department of Botany
Chicago Natural History Museum

Phase Transformations in Solids. Symposium held at Cornell University, August 23-26, 1948. Sponsored by the Committee on Solids, Division of Physical Sciences, the National Research Council. R. Smoluchowski, J. E. Mayer, and W. A. Weyl, Eds. New York: Wiley; London: Chapman & Hall, 1951. 660 pp. \$9.50.

This handsome, well-printed volume resulted from a symposium at Cornell University on the subject indicated in the title. The symposium was organized by the National Research Council for the purpose of bringing together physicists, mineralogists, and metallurgists, to discuss a topic of common interest. The topic is one that appears easy on casual approach, but becomes harder as the bewildering variety of facts is passed in review. It is not surprising, in view of this, that the three groups of scientists have developed a slightly different slant on the same problem. Among these points of view, the crystallographer's appears the most mature. The physicist proposes and tears down his theories somewhat too quickly to be taken seriously, and the material accumulated by the metallurgist is lacking in scientific precision. Indeed, one mineralogist remarked somewhat sarcastically during the conference that the only crystal planes the metallurgists seem to have heard of are the 100, the 110, and the 111 planes.

The first few articles deal with the more profound aspects of equilibrium theory, and a middle section deals with phase diagrams, but the greater part of the book deals with rate-determining mechanisms such as diffusion or nucleation. In the experimental work concerned with these questions it is hard to come to clear formulations; hence, the material accumulated tends to become bulky.

In spite of the number of contributing experts, it seems to this reviewer that the book does not quite correspond to the effort that went into it. In the first place, the book appeared late: almost three years elapsed between the symposium and its appearance. It is true that several papers were partially or completely rewritten to take care of recent developments. But this can only partly remedy an unhappy situation. A conference report is most useful in the period immediately following the conference, when the scientists working in the field are trying to orient their thinking. With a delay of three years, research workers in the field must have turned to other sources of inspiration.

When the book is considered as a reference work

rather than as a conference report the delay is obviously much less important. However, it suffers then from other defects. A record in book form of even a successful conference does not make a first-rate handbook. Some speakers have justifiably taken a personal approach, others have written a survey article of recent work, giving all viewpoints. A person consulting such a volume in a library would probably be best served by a handbook-type article—that is, a survey limited to the work judged good by the author, but with a tolerant viewpoint and without discrimination between the old and the new. Such books would need centralized editorship, with an index at the end and clear division between the sections. The need for them becomes more urgent every year. The great German physics handbooks are now 20 to 30 years old and need replacement badly. If the National Research Council could sponsor such books it would earn the gratitude of every scientist.

GREGORY H. WANNIER

Bell Telephone Laboratories
Murray Hill, New Jersey

Finite Deformation of an Elastic Solid. Francis D. Murnaghan. New York: Wiley; London: Chapman & Hall, 1951. 140 pp. \$4.00.

This book is apparently intended to serve as a text, rather than as a treatise for research workers. The author regards the treatment as elementary. Matrix methods are used throughout, and the pages are dense with calculations.

After some classical results on the general theory of elasticity the author expands the strain energy in a power series, which he truncates after the cubic terms; the remainder of the book (with one exception presently to be noted) is devoted to this second approximation theory. The special forms assumed by the cubic terms for the various types of crystals are determined. There follow treatments of simple shear both for isotropic and for a certain aeolotropic material, simple tension, compression of a spherical shell and cylindrical tube, and torsion of a circular cylinder. The exception to the method of power series expansion is the treatment of hydrostatic pressure, where the author obtains what he calls an integrated linear theory by assuming that the ordinary linear elastic coefficients are linear functions of pressure.

Although the author's various approximate formulas may be useful in certain applications concerning moderate strains, it is unfortunate that his book makes no mention of the more fundamental recent researches in finite elasticity theory by Rivlin and others, where the form of the strain energy is left arbitrary and results, directly and successfully comparable with experiments on very large strain of rubber, are calculated in full generality.

C. A. TRUESDELL

Institute for Applied Mathematics
Indiana University