

Technical Papers

Curarization by Rectal Suppository

Jean Delay and Jean E. Thuillier

Faculté de Médecine de Paris, Laboratoire de Biologie
de la Clinique des Maladies Mentales
et de l'Encéphale, Paris, France

We have found that the introduction into the rectum of a rabbit of a suppository containing a sufficient quantity of curare or curarelike compounds produces the same effect as that caused by intravenous injection. Curarization of the rabbit follows the same pattern in both cases: first, the ears drop; then there is a softening of the muscles of the neck; the head drops; and finally a complete quadriplegia is observed. In the case of an overdose this is followed by paralysis of the diaphragm with cessation of breathing; and, last, death is caused by heart failure.

Curarization by rectal suppository is less abrupt than by intravenous injection, taking 4–6 min, and it lasts longer. Relaxation continues for 3–5 hr. Physostigmine and eserine keep their effect as antidotes, permitting one to stop the curarization at will or in case of an overdose.

In order to demonstrate more objectively the effect of rectal curarization, we have made a neuromuscular preparation. A rabbit is slightly anesthetized by Nembutal injection. The sciatic nerve is isolated, and the inner branch receives current through the electrodes of a battery-operated excitor and is regularly stimulated to a constant rhythm. The gastrocnemius muscle of the animal is then prepared and a sheaf from the tendon is attached to a myograph which registers the contractions provoked by the excitation of the sciatic nerve. One thus obtains a graph of the muscular contraction, with constant amplitude and rhythm. If one introduces into the rectum of the rabbit

a suppository containing 0.025 g/kg of 336 HC (iodoethylate of N,N bis-piperidyl-ethyl-piperazine) or 0.005 g/kg of D-tubocurarine or 0.005 g/kg of Flaxedil, there is a variable delay, constant for a given animal, after which one observes that the amplitude of the contractions decreases until there is no contraction at all, and the animal is completely curarized. Graphs A and B in Fig. 1 show the difference of effect between the suppository and the intravenous injection; C shows the antagonistic action of physostigmine.

These differences may be summarized as follows: Rectal administration by suppository permits (a) slower curarization; (b) a more persistent effect; and (c) a slower recuperation of muscular contractility.

This experimental procedure demonstrates for the first time, as far as we know, the use of a rectal suppository for the curarization of an animal. In clinical investigations, we have obtained interesting results in neurological and psychiatric syndromes by this procedure (1).

Reference

1. DELAY, J., et al. *Presse méd.*, **60**, (63), 1341 (1952).

Manuscript received June 24, 1952.

Invertase in *Palamnaeus bengalensis*

M. B. Lal and M. S. Kanungo

Department of Zoology,
The University, Lucknow, India

As far as the writers are aware, knowledge of the physiology of digestion in scorpions is largely due to the extensive investigations of Pavlovsky and Zarin (1) on *Buthus*, but unfortunately these authors had little fresh material at their disposal and based their work on stored glycerin extracts. No invertase was, however, detected by these workers in the scorpion *Buthus*.

Our investigations were conducted on *Palamnaeus bengalensis*, the common large Indian scorpion, which lives in dry, shady, uncultivated high platforms of land and feeds on various types of insects, myriapods, and even on its own kind. Large quantities of fresh extracts of the liver were prepared from a number of well-fed specimens that had been lightly chloroformed and immediately dissected. Weighed quantities of the glands were ground with quartz sand and measured quantities of distilled water and filtered. The solution so obtained gave a pH of 7.1. Estimations of the enzyme were conducted, using the methods of Cole (2), Hawk et al. (3), Sumner and Somers (4), and Smith (5). All experiments were performed with parallel controls and blank solutions. Toluene was used as an antiseptic in each case.

For the detection of invertase buffered extracts




FIG. 1. Rabbit ♂ 3 kg neuromuscular preparation. (A) Contractions of the gastrocnemius muscle; curarization by rectal suppository. (B) Contractions of the gastrocnemius muscle; curarization by intravenous injection. (C) Decurarizing action by intravenous injection of physostigmine.