

Reg. U. S.

GBI

Pat. Off.

SPECIAL RESEARCH BIOCHEMICALS

Allantoin
 Alloxan
 Arterenol (nor-adrenaline)
 Bilirubin
 Blood Fibrin
 L-Borneol
 Cadaverine
 Calcium Phytate
 Cephalin
 Cholesterol U.S.P.
 2,4 Dichloroanisole
 2,4 Dinitrophenol (M.P. 112-113° C.)
 Epichlorohydrin
 Fibroin (Silk)
 Flavianic Acid (2,4, Dinitro-1-Naphthol-7-Sulfonic Acid)
 Fumaric Acid 90% (Pract.)
 Glutaric Acid
 Glutathione (Reduced)
 Glutathione Sodium Salt (Reduced)
 Glutathione (Oxidized)
 Glycogen
 Histamine (Free Base)
 Histamine C.P. Dihydrochloride
 Histamine Diphosphate, U.S.P.
 Indole-3-Acetic Acid
 Indole Butyric Acid
 Indole-3-Propionic Acid
 Inosinic Acid (barium salt)
 Inositol Hexaphosphoric Acid

See Science October 9 issue
for additional listings

A READY REFERENCE
That will Save Time for You

Use this catalog as a "one stop" source of Amino Acids, Vitamins, Carbohydrates, Adenylates, Nucleates, Purines, Pyrimidines, Tetrazolium Salts, Enzymes, Microbiological and Bacteriological Media, Complete Animal Test Diets and Ingredients for investigational use.

WRITE FOR
YOUR COPY
TODAY

GENERAL BIOCHEMICALS, INC.
60 LABORATORY PARK • CHAGRIN FALLS, OHIO

A MIGHTY LIGHT FROM A SMALL SOURCE

"POINTOLITE" LAMPS

Laboratory workers rely on Pointolite's dependable, steady light that requires no attention after starting. Illumination is an intensely brilliant, remarkably white, "point of light". Designed to fit scientific optical instruments for photographic and projection purposes. Lamps in various candle-power sizes and necessary auxiliary equipment are described and listed in Bulletin S.



JAMES G. BIDDLE CO.

ELECTRICAL & SCIENTIFIC INSTRUMENTS
1316 ARCH STREET • PHILADELPHIA 7, PENNA.

LaMOTTE STANDARDS FOR ANALYSIS

TIRON

(disodium-1, 2-dihydroxybenzene-3, 5-disulfonate)

For the Determination of Iron

A new sensitive, stable, and widely applicable reagent, developed by John H. Yoe and A. Letcher Jones, University of Virginia (Reference—Industrial and Engineering Chemistry, Analytical Edition, Vol. 16, p. 111-115, 1944), for the colorimetric determination of ferric iron. The reagent may be used in either acid or alkaline medium. In alkaline medium it is sensitive to one part of iron in 100,000,000 parts of solution. In acid solution the sensitivity is one part in 30,000,000.

TIRON—YOE & JONES IRON REAGENT

1 g	5 g	10 g	25 g
\$0.50	\$1.50	\$2.50	\$5.00

For the Determination of TITANIUM (Method of Yoe & Armstrong)

TIRON is an extremely sensitive reagent for the determination of TITANIUM (indicated sensitivity, 1 part of TITANIUM in 100,000,000 parts of solution). Only a few ions interfere. Furthermore, TIRON may be used to determine both Titanium and Iron in same solution. TIRON REAGENT available from stock.

LaMotte Chemical Products Company
Dept. "H" Towson 4, Md.