

Technical Papers

One-Step Preparation of C^{14} -Cyanide from Barium Carbonate- C^{14}

J. K. Jeanes

Radiochemical Department,
Texas Research Foundation, Renner

The importance of cyanide in the synthesis of organic compounds labeled with isotopic carbon requires that it be made available in high yields by simple and inexpensive procedures starting with barium carbonate. Methods (1-5) have been described for the conversion of alkali carbonates to cyanide but each has its own limitation. A procedure has been developed in this laboratory based on the findings of Hood and Salamon (6) and J. A. McCarter (7) whereby barium carbonate can be converted to sodium cyanide in one step. This was accomplished by heating barium carbonate, zinc dust, and metallic sodium in a stream of anhydrous ammonia which had passed over hot iron. Iron seemed to have an important role in that it catalyzed the decomposition of ammonia to give nitrogen gas and other "active" forms of nitrogen which were essential in the reaction. The yields in this laboratory were quantitative as described in the following experimental procedure.

Powdered anhydrous C^{14} barium carbonate (1.0 millimole) was mixed with 1.0 g of zinc dust and 0.2000 g of metallic sodium in pea size chunks, and this mixture was transferred to a porcelain combustion boat of suitable size. The boat was placed in a Vycor combustion tube (600 mm in length and 19 mm inside diameter) in an atmosphere of anhydrous ammonia and containing a 5.0-g ball of iron wire or 5.0 g of powdered iron distributed throughout a plug of Pyrex glass wool occupying the mid-portion of the tube. The boat was pushed into the tube until it touched, or almost touched, the iron plug.

The end of the tube nearest the iron was connected to a cylinder of anhydrous ammonia. A stream of ammonia was allowed to flow through a gas bubbler at the rate of 3 bubbles/second in and through the tube containing the mixture. The portion of the tube containing the mixture and iron was brought to and maintained at approximately 650°C for a period of 4 hr. The flow of ammonia was continued until the tube was cool.

The boat and all contents of the tube, except the iron ball, were washed into a 250-ml flask attached to a Kjeldahl distillation head for subsequent distillation of hydrogen cyanide. The solution in the flask was acidified with dilute (2 *N*) sulfuric acid and 20-30 ml of distillate were collected in an Erlenmeyer flask containing a 20% excess of the theoretical amount of 1 *N* sodium or potassium hydroxide.

Analysis by the argentimetric method (8) of the distillate obtained in three experiments showed a yield

of cyanide which was quantitative. The specific activity of the labeled cyanide was unchanged from that of the barium carbonate used as the starting material.

References

1. CRAMER, R. D., and KISTIAKOWSKY, G. B. *J. Biol. Chem.*, **137**, 547 (1941).
2. ADAMSON, A. W. *J. Am. Chem. Soc.*, **69**, 2564 (1947).
3. ABRAMS, R. *Ibid.*, **71**, 3835 (1949).
4. BELLEAU, B., and HEARD, R. D. H. *Ibid.*, **72**, 4268 (1950).
5. LOFFFIELD, R. B. *Nucleonics*, **1**, 54 (1947).
6. HOOD, J. J., and SALAMON, A. G. *Chem. Zentr.*, **66**, I, 670 (1895).
7. MCCARTER, J. A. *J. Am. Chem. Soc.*, **73**, 483 (1951).
8. KOLTHOFF, I. M., and SANDELL, E. B. *Textbook of Quantitative Inorganic Analysis*, Rev. Ed. New York: Macmillan, 1943, p. 574.

Manuscript received October 8, 1953.

Evaluation of Oral Temperature Readings

Francis L. Harmon

Department of Psychology,
Saint Louis University, Saint Louis, Missouri

Many studies have dealt with the variability of body temperature as reflected in inter-individual differences, day-to-day changes, diurnal variations, and other comparisons (1). Such studies, while yielding considerable information as to the extent of variability to be expected under different conditions, have failed to provide any systematic method for evaluating the significance of a given temperature reading, or a difference between two readings taken under specified circumstances. One possible solution to this problem is presented in the following paragraphs; it involves an approach which has been used in connection with a similar question regarding metabolism measurements (2).

The data for this analysis were taken from 29 ostensibly normal male university students, ranging in age from 17 to 27 years. Twelve readings of oral temperature were made upon each subject, the complete series consisting of records taken at 8:00 A.M. (before breakfast), 12:00 M., 6:00 P.M., and 10:00 P.M. on each of 3 different days, approximately a week apart. Only 1 subject was observed on any given day. Every session was preceded by a 30-min rest, the subject lying upon a cot throughout this period and during the actual observation of his temperature. All readings were made with the same thermometer, a MICH 3 Oral (Serial J 1273).¹ The thermometer always was inserted under the subject's tongue and left in place a full 3 min.

The 348 temperature readings thus obtained ranged from 95.4° to 99.1° , and averaged 98.0°F , a value very close to Ivy's mean of 98.1° for his 276 medical students, tested in class between 8:00 and 9:00 A.M. (3). Individual means for our 29 subjects ranged

¹ Manufactured by E. Kessling, New York, N. Y.